

Transfer profile of intramuscularly administered tetrodotoxin to non-toxic cultured specimens of the pufferfish *Takifugu rubripes*

Koichi Ikeda^a, Yumi Murakami^a, Yu Emoto^b, Laymithuna Ngy^a, Shigeto Taniyama^a, Motoaki Yagi^c, Tomohiro Takatani^b, Osamu Arakawa^{b,*}

^a*Graduate School of Science and Technology, Nagasaki University, Nagasaki 852-8521, Japan*

^b*Faculty of Fisheries, Nagasaki University, Nagasaki 852-8521, Japan*

^c*Nagasaki Municipal Fisheries Center, Nagasaki 852-0114, Japan*

*Corresponding author. Tel./fax: +81 95 819 2844.

E-mail address: arakawa@nagasaki-u.ac.jp

Abstract

Tetrodotoxin (TTX) was intramuscularly administered to non-toxic cultured specimens of the pufferfish *Takifugu rubripes* to investigate TTX transfer/accumulation profiles in the pufferfish body. In two groups of test fish administered either 50 MU/individual of TTX standard (purified TTX; PTTX) or crude extract of toxic pufferfish ovary (crude TTX; CTTX), TTX rapidly transferred from the muscle via the blood to other organs. The toxin transfer profiles differed between groups, however, from 4 to 72 h. In the PTTX group, little TTX was retained in the liver, and most (> 96%) of the toxin remaining in the body transferred/accumulated in the skin after 12 h, whereas in the CTTX group, a considerable amount of toxin (15%-23% of the administered toxin or 28%-58% of the remaining toxin) was transferred/retained in the liver for up to 24 h, despite the fact that 89% of the remaining toxin transferred/accumulated in the skin at the end of rearing period (168 h). The total amount of toxin remaining in the entire body at 1 to 4 h was approximately 60% of the administered toxin in both groups, which decreased at 8 to 12 h, and then increased again to approximately 60% to 80% at 24 to 168 h. Immunohistochemical observation revealed that the toxin accumulated in the skin was localized at the basal cells of the epidermal layer.

Keywords: Tetrodotoxin; pufferfish; *Takifugu rubripes*; intramuscular administration; immunohistochemical observation

1. Introduction

The pufferfish *Takifugu rubripes*, as well as many marine pufferfish of the family Tetraodontidae, possess a potent neurotoxin, tetrodotoxin (TTX). In wild adult *T. rubripes*, the liver and ovary usually have strong toxicity, whereas the muscle, skin, and testes are non-toxic and are safe for human consumption (Noguchi and Arakawa, 2008). TTX is originally produced by marine bacteria, and distributed over a wide variety of animals other than pufferfish, including gobies, blue-ringed octopuses, carnivorous gastropods, starfish, toxic crabs, horseshoe crabs, flat worms, and ribbon worms (Miyazawa and Noguchi, 2001). The facts that pufferfish become non-toxic when fed non-toxic diets in an environment in which the invasion of TTX-bearing organisms has been eliminated (Matsui et al., 1982; Saito et al., 1984; Noguchi et al., 2006), and that such non-toxic pufferfish become toxic when orally administered TTX (Matsui et al., 1981, Yamamori et al., 2004, Honda et al., 2005, Kono et al., 2008a), indicate that TTX is exogenous in pufferfish and is derived from the food chain that starts from bacteria (Noguchi and Arakawa, 2008). The transfer, accumulation, and elimination mechanisms of TTX taken up into the pufferfish body via food organisms remain unclear. In our studies to clarify this point, we investigated the short-term transfer and accumulation profiles of TTX intramuscularly administered to non-toxic cultured specimens of *T. rubripes*. In oral administration experiments, long-term toxin accumulation is observed, but these experiments are not suitable for tracing short-term inter-tissue toxin transfer, because it is difficult to accurately administer a single large dose of toxin (Honda et al., 2005). To overcome this problem in the present study, we administered the TTX intramuscularly. Matsui et al. (1981) reported that when non-toxic cultured specimens of *T. rubripes* are fed diets containing crystalline TTX or crude toxic pufferfish ovary extract, only the test fish fed the crude extract of toxic pufferfish ovary accumulated TTX in their liver. Based on this information, we administered two types of toxins, ‘purified TTX’ and ‘crude TTX’, to evaluate whether the transfer profiles differed after entering the pufferfish body.

2. Materials and methods

2.1. Pufferfish specimens

Non-toxic cultured specimens of *T. rubripes* (approximately 4 months old; body weight, 13.2 ± 3.4 g; body length, 7.1 ± 0.6 cm; $n = 80$) (Noguchi et al., 2006) were purchased from a culture farm in Toishi, Nagasaki Prefecture, Japan. The specimens were acclimatized in aerated tanks for several days before administration of the toxin.

2.2. Preparation of toxin solutions

Toxic ovaries of the pufferfish *Takifugu vermicularis* were extracted with 1% acetic acid in 80% methanol, and the extract was defatted with dichloromethane and evaporated to make a condensed toxin solution (designated crude TTX). The toxicity of the crude TTX was evaluated using a mouse bioassay according to the official guidelines of the Japan Food Hygiene Association (2005). Lethal potency was expressed in mouse units (MU), where 1 MU was defined as the amount of toxin required to kill a 20-g male ddY strain mouse within 30 min after intraperitoneal administration. Liquid chromatography/mass spectrometry (LC/MS) analysis (Nakashima et al., 2004) revealed that the crude TTX was composed mainly of TTX and its analogues, such as 4-*epi*TTX and 4,9-anhydroTTX; TTX alone accounted for more than 90% of the total toxicity (data not shown).

Both TTX standards, purchased from Wako (purity > 90%; designated purified TTX) and crude TTX, were dissolved or diluted individually with a physiologic saline solution containing 1.35% NaCl, 0.06% KCl, 0.025% CaCl₂, 0.035% MgCl₂, and 0.02% NaHCO₃ at a concentration of 500 MU/ml and used in the following toxin administration experiments.

2.3. Toxin administration experiments

The acclimatized pufferfish specimens were divided into two groups of 40 individuals; one group was administered purified TTX (PTTX group) and the other was administered crude TTX (CTTX group). The groups were then maintained separately in two aerated 90-l tanks. Each fish was intramuscularly administered 0.1 ml (50 MU) of either purified or crude TTX solution and immediately returned to the tank (total handling time <30 s/individual to minimize stress to the fish). Then, 5 fish from each group were randomly collected at 1, 4, 8, 12, 24, 72, 120, and 168 h after toxin administration and toxin quantification was performed as described below.

2.4. Toxin quantification

Using a syringe precoated with sodium heparin, blood was withdrawn from the portal vein of each fish and centrifuged at 4200 g for 10 min. As TTX is partially binding to the TTX/PSP-binding protein in pufferfish blood plasma (Matsui et al., 2000; Yotsu-Yamashita et al., 2001), the supernatant (blood plasma) obtained was added with acetic acid at a final concentration of 0.1% to cut the binding, ultrafiltered through an Ultrafree-MC 5000 NMWL (Millipore Corp., Bedford, MA), and then submitted to enzyme-linked immunosorbent assay (ELISA) for TTX. After blood collection, all specimens were dissected into different

anatomic tissues (liver, skin, and muscle), which were extracted with 0.1% acetic acid (Japan Food Hygiene Association, 2005). Each tissue extract was filtered through a USY-1 membrane (0.45 μm ; Toyo Roshi Co., Ltd, Japan) and submitted to ELISA.

ELISA was performed according to the previously reported method (Ngy et al., 2008) using a monoclonal anti-TTX antibody developed by Kawatsu et al. (1997). The amount of TTX (in ng) determined by ELISA was converted to MU based on the specific toxicity of TTX (220 ng/MU). In a preliminary experiment using crude liver extracts ($n = 10$) of the pufferfish *Takifugu poecilonotus*, a significant and positive correlation (Pearson's test; $r = 0.9641$, $p < 0.01$) was observed between the TTX amounts determined by ELISA and those calculated from the TTX peak areas in LC/MS (Nakashima et al., 2004) (Figure 1). The regression line, $y = 0.9874 x + 7.301$ ($r^2 = 0.9295$) indicated that TTX was selectively quantified by ELISA in the presence of some TTX analogs including 4-*epi*TTX, 4,9-anhydroTTX, and deoxyTTXs (Yotsu-Yamashita, 2001) that were detectable in the extracts by LC/MS (data not shown).

2.5. Immunohistochemical observation

A part of the skin of each fish collected at 120 h after toxin administration was submitted to the immunohistochemical observation under light microscope according to the previously reported method (Tanu et al., 2002; Mahmud et al., 2003a,b) using the anti-TTX antibody.

3. Results

Changes in the toxin content (MU/g or MU/ml) of each pufferfish tissue during the rearing period are shown in Figure 2. Changes in the toxin content of the liver differed between the PTTX and CTTX groups. In both groups, the toxin content was 12 MU/g at 1 h after administration. In the PTTX group, the toxin content gradually decreased until only a small amount (0.5-1.5 MU/g) remained after 12 h. In the CTTX group, however, the toxin content increased, reaching a maximum (around 21 MU/g) at 8 to 12 h; thereafter, the toxin content decreased, but 6.4 MU/g remained at the end of the rearing period (168 h). Changes in the skin toxin content also differed between groups. In both groups, the toxin content increased remarkably between 1 and 72 h and then remained at about 15 MU/g. The onset of the increase in the PTTX group occurred earlier than that in the CTTX group (after 12 h). The toxin content of the muscle, which was the site of administration, rapidly decreased and after 8 h was below the detection limit (0.01 MU/g) in both groups. The toxin content of the blood

plasma was highest at 1 h (5.9 MU/ml in the PTTX group, 9.8 MU/ml in the CTTX group), and rapidly decreased thereafter.

Changes in the anatomic distribution of TTX, demonstrated by the relative amount of toxin retained in each tissue [% of the administered toxin (50 MU/individual)], are shown in Figure 3. The total amount of toxin remaining in the whole body at 1 to 4 h was around 60% of the administered toxin (50 MU/individual) in both groups. The amount first began to decrease at 8 to 12 h, and then increased to approximately 60% to 80% at 24 to 168 h. Changes in the amount of toxin in the liver tissues differed between the PTTX and CTTX groups. In the PTTX group, the amount of toxin in the liver rapidly decreased from 1 to 8 h, becoming less than 1.6% of the administered toxin (< 3.2% of the remaining toxin) after 12 h. In the CTTX group, the amount of toxin in the liver did not decrease for up to 24 h and accounted for 15% to 23% of the administered toxin (28%-58% of the remaining toxin). The amount of toxin in the skin gradually increased during the rearing period in both groups, and accounted for most (PTTX group, 98%; CTTX group, 89%) of the remaining toxin at 168 h.

Light micrographs of representative skin sections at 120 h after toxin administration are shown in Figure 4. The epidermal layer of the skin was comprised of two distinct cell types, basal cells and succiform cells (Tanu et al., 2002; Mahmud et al., 2003a,b), and no gland or gland-like structure was observed. In both the PTTX and CTTX groups, positive reactions for TTX (brown color) were localized at basal cells along the basement membrane. No positive reaction was observed in the succiform cells, or in the skin sections of negative control, i.e., the fish without toxin administration (data not shown).

4. Discussion

TTX intramuscularly administered to non-toxic cultured *T. rubripes* rapidly transferred to other body tissues, and the toxin content of the liver and skin exceeded that of muscle within as little as 1 h after administration. At 1 h after intramuscular administration, a high concentration of TTX was present in the blood plasma, indicating that TTX transferred mainly via the bloodstream. The fact that muscles in toxic wild specimens of *T. rubripes* are not toxic indicates that the muscles of this species either do not retain and accumulate TTX or have a mechanism for eliminating TTX.

The toxin transfer profiles from 4 to 72 h differed between the PTTX and CTTX groups. In the PTTX group, little TTX was retained in the liver, and most of the toxin transferred to, and accumulated in, the skin after 12 h, whereas in the CTTX group, a considerable amount of toxin was transferred to, and retained in, the liver for up to 24 h. Although most of the

toxin transferred to, and accumulated in, the skin thereafter, some toxin remained in the liver even at 168 h. Matsui et al. (1981) reported that when non-toxic cultured specimens of *T. rubripes* are fed diets containing crystalline TTX or crude extract of toxic pufferfish ovary, only the test fish fed with the toxic pufferfish ovary accumulate TTX in their liver. The liver tissue of *T. rubripes* is equipped with a specific TTX uptake mechanism (Nagashima et al., 2003; Matsumoto et al., 2005, 2007), suggesting that some substance(s) coexisting in the crude TTX might enhance the uptake mechanism or change TTX molecules into a form that is more easily processed by this mechanism, resulting in the above mentioned difference. Kono et al. (2008b) recently reported that 40% of purified TTX intramuscularly administered to the cultured pufferfish *Takifugu (Fugu) niphobles* was transformed to 4,9-anhydroTTX within 4 days after administration. It is unclear, however, whether such transformation between TTX and its analog(s) was involved in the concerned difference. Further studies are needed to clarify these points.

In both the PTTX and CTTX groups, most of the toxin that remained in the body was eventually transferred/accumulated in the skin. Wild adult specimens of *T. rubripes* generally possess no toxin in the skin, but toxicity of several tens MU is occasionally detected in juveniles (unpublished data). The test fish used in the present study were 4 months old and therefore considered to be in the juvenile stage. Therefore, these specimens are assumed to have an immature skin TTX-excreting ability or liver and ovary TTX-accumulating ability. Our previous immunohistochemical investigations (Tanu et al., 2002; Mahmud et al., 2003a, b) revealed that in the pufferfish having toxic skin, such as *T. vermicularis*, *Chelonodon patoca*, *Tetraodon steindachneri*, and *Tetraodon nigroviridis*, TTX is mainly found in the secretory glands or secretory cells (succiform cells) of their skin. The succiform cells of the present test fish, however, showed no positive reaction for TTX, suggesting that TTX, in *T. rubripes* juveniles, remains at basal cells and does not easily reach the succiform cells, which provably excrete TTX in adult fish.

When non-toxic cultured specimens of *Tetraodon turgidus*, the freshwater pufferfish that possesses paralytic shellfish toxin (PST) in the skin, are intramuscularly administered PST, a toxin similar to that used in the present study, the PST rapidly transfers from muscle to other tissues and accumulates mostly in the skin at the end of rearing period (Ngy et al., 2008). Interestingly, when *T. turgidus* specimens were administered the same dosage of TTX, all died within 3 to 4 h, and more than half of the TTX administered remained in the muscle in the dead specimens. It is therefore inferred that marine pufferfish that ingest TTX are thus endowed with a mechanism by which they transport TTX specifically and actively, and

freshwater pufferfish that ingest PST are endowed with a mechanism that processes PST. TTX/PST-binding proteins have been isolated from the blood plasma of marine pufferfish (Matsui et al., 2000; Yotsu-Yamashita et al., 2001), and may be involved in the transportation mechanism.

In both the PTTX and CTTX groups, a temporary decrease in the total amount of toxin remaining in the whole body was observed at 8 to 12 h. A similar decrease was observed when *T. turgidus* was administered PST (Ngy et al., 2008). Watabe et al. (1987) reported that when *T. rubripes* was intraperitoneally administered [³H]-TTX, the amount of TTX in the gallbladder was greatly increased 6 days after administration. The temporary decrease may be due to the temporary storage of a large amount of TTX in a particular organ or tissue other than muscle, liver, and skin. This point, along with the properties of the TTX/PST-binding proteins and specific toxin transportation/accumulation mechanisms, remain to be elucidated. Further studies are in progress.

Acknowledgements

We would like to express sincere thanks to Dr. Kentaro Kawatsu and Dr. Yonekazu Hamano of Osaka Prefectural Institute of Public Health, Japan, for providing the anti-TTX antibody. This work was partly supported by a Grant-in-aid from the Ministry of Education, Culture, Sports, Science, and Technology, Japan.

Conflict of Interest statement

The authors declare that there are no conflicts of interest.

References

- Honda, S., Arakawa, O., Takatani, T., Tachibana, K., Yagi, M., Tanigawa, A., Noguchi, T., 2005. Toxification of cultured puffer fish *Takifugu rubripes* by feeding on tetrodotoxin-containing diet. *Nippon Suisan Gakkaishi* 71, 815–820.
- Japan Food Hygiene Association, 2005. Puffer toxin. In: Environmental Health Bureau, Ministry of Health and Welfare, (Ed.), *Shokuhin Eisei Kensa Shishin (Manual for Methods for Food Sanitation Testing)*, Tokyo, pp. 661–673.
- Kawatsu, K., Hamano, Y., Yoda, T., Terano, Y., Shibata, T., 1997. Rapid and highly sensitive enzyme immunoassay for quantitative determination of tetrodotoxin. *Jpn. J. Med. Sci. Biol.* 50, 133–150.
- Kono, M., Matsui, T., Furukawa, K., Yotsu-Yamashita, M., Yamamori, K., 2008a. Accumulation of tetrodotoxin and 4,9-anhydrotetrodotoxin in cultured juvenile kusafugu *Fugu niphobles* by dietary administration of natural toxic komonfugu *Fugu poecilonotus* liver. *Toxicon* 51, 1269–1273.
- Kono, M., Matsui, T., Furukawa, K., Takase, T., Yamamori, K., Kaneda, H., Aoki, D., Jang, J.H., Yotsu-Yamashita, M., 2008b. Examination of transformation among tetrodotoxin and its analogs in the living cultured juvenile puffer fish, kusafugu, *Fugu niphobles* by intramuscular administration. *Toxicon*, in press.
- Mahmud, Y., Okada, K., Takatani, T., Kawatsu, K., Hamano, Y., Arakawa, O., Noguchi, T., 2003a. Intra-tissue distribution of tetrodotoxin in two marine puffers *Takifugu vermicularis* and *Chelonodon patoca*. *Toxicon* 41, 13-18.
- Mahmud, Y., Arakawa, O., Ichinose, A., Tanu, M.B., Takatani, T., Tsuruda, K., Kawatsu, K., Hamano, Y., Noguchi, T., 2003b. Intracellular visualization of tetrodotoxin (TTX) in the skin of a puffer *Tetraodon nigroviridis* by immunoenzymatic technique. *Toxicon* 41, 605-611.
- Matsui, T., Hamada, S., Konosu, S., 1981. Difference in accumulation of puffer fish toxin and crystalline tetrodotoxin in the puffer fish, *Fugu rubripes rubripes*. *Bull. Jpn. Soc. Sci. Fish.* 47, 535–537.
- Matsui, T., Sato, H., Hamada, S., Shimizu, C., 1982. Comparison of toxicity of the cultured and wild puffer fish *Fugu niphobles*. *Bull. Jpn. Soc. Sci. Fish.* 48, 253.
- Matsui, T., Yamamori, K., Furukawa, K., Kono, M., 2000. Purification and some properties of a tetrodotoxin binding protein from the blood plasma of kusafugu, *Takifugu niphobles*. *Toxicon* 38, 463–468.

- Matsumoto, T., Nagashima, Y., Takayama, K., Shimakura, K., Shiomi, K., 2005. Difference between tetrodotoxin and saxitoxins in accumulation in puffer fish *Takifugu rubripes* liver tissue slices. *Fish Physiol. Biochem.* 31, 95-100.
- Matsumoto, T., Nagashima, Y., Kusuhara, H., Sugiyama, Y., Ishizaki, S., Shimakura, K., Shiomi, K., 2007. Involvement of carrier-mediated transport system in uptake of tetrodotoxin into liver tissue slices of puffer fish *Takifugu rubripes*. *Toxicon* 50, 173–179.
- Miyazawa, K., Noguchi, T., 2001. Distribution and origin of tetrodotoxin. *J. Toxicol.-Toxin Rev.* 20, 11–33.
- Nagashima, Y., Toyoda, M., Hasobe, M., Shimakura, K., Shiomi, K., 2003. In vitro accumulation of tetrodotoxin in pufferfish liver tissue slices. *Toxicon* 41, 569-547.
- Nakashima, K., Arakawa, O., Taniyama, S., Nonaka, M., Takatani, T., Yamamori, K., Fuchi, Y., Noguchi, T., 2004. Occurrence of saxitoxins as a major toxin in the ovary of a marine puffer *Arothron firmamentum*. *Toxicon* 43, 207–212.
- Ngy, L., Tada, K., Yu, C.-F., Takatani, T., Arakawa, O., 2008. Occurrence of paralytic shellfish toxins in Cambodian Mekong pufferfish *Tetraodon turgidus*: selective toxin accumulation in the skin. *Toxicon* 51, 280–288.
- Noguchi, T., Arakawa, O., 2008. Tetrodotoxin – Distribution and accumulation in aquatic organisms, and cases of human intoxication. *Marine Drugs* 6, 220–242.
- Noguchi, T., Arakawa, O., Takatani, T., 2006. Toxicity of pufferfish *Takifugu rubripes* cultured in netcages at sea or aquaria on land. *Comp. Biochem. Physiol. Part D* 1, 153–157.
- Saito, T., Maruyama, J., Kanoh, S., Jeon, J.K., Noguchi, T., Harada, T., Murata, O., Hashimoto, K., 1984. Toxicity of the cultured pufferfish *Fugu rubripes rubripes* along with their resistibility against tetrodotoxin. *Bull. Jpn. Soc. Sci. Fish.* 50, 1573–1575.
- Tanu, M.B., Mahmud, Y., Takatani, T., Kawatsu, K., Hamano, Y., Arakawa, O., Noguchi, T., 2002. Localization of tetrodotoxin in the skin of a brackishwater puffer *Tetraodon steindachneri* on the basis of immunohistological study. *Toxicon* 40, 103-106.
- Watabe, S., Sato, Y., Nakaya, M., Nogawa, N., Oohashi, K., Noguchi, T., Morikawa, N., Hashimoto, K., 1987. Distribution of tritiated tetrodotoxin administered intraperitoneally to pufferfish. *Toxicon* 25, 1283–1289.
- Yamamori, K., Kono, M., Furukawa, K., Matsui, T., 2004. The toxification of juvenile cultured kusafugu *Takifugu niphobles* by oral administration of crystalline tetrodotoxin. *J. Food Hyg. Soc. Jpn.* 45, 73–75.

- Yotsu-Yamashita, M., 2001. Chemistry of puffer fish toxin. *J. Toxicol.-Toxin Rev.* 20, 51-66.
- Yotsu-Yamashita, M., Sugimoto, A., Terakawa, T., Shoji, Y., Miyazawa, T., Yasumoto, T., 2001. Purification, characterization, and cDNA cloning of a novel soluble saxitoxin and tetrodotoxin binding protein from plasma of the puffer fish, *Fugu pardalis*. *Eur. J. Biochem.* 268, 5937–5946.

Figure Captions

Fig. 1. Comparison of TTX amounts determined by LC/MS and ELISA.

Fig. 2. Changes in the content of TTX (MU/g or MU/ml) retained in each tissue of the *T. rubripes* specimens during the rearing period after toxin administration.

▣: PTTX group; ●: CTTX group.

The toxin content in blood plasma was determined using a combined sample of 5 individuals for each point.

Fig. 3. Changes in the relative amount of TTX [% of the administered amount (50 MU/individual)] retained in each tissue of the *T. rubripes* specimens during the rearing period after toxin administration.

A: PTTX group; B: CTTX group.

Fig. 4. Light micrographs of representative skin sections (x 100) of the test fish collected at 120 h after toxin administration, showing TTX-positive basal cells (arrow heads) and TTX-negative succiform cells (arrows). A: PTTX group; B: CTTX group.




Fig. 1








Fig. 2

(A) PTTX group



(B) CTTX group




Fig. 3

(A) PTTX group



(B) CTTX group



Fig. 4