

Conformational studies on peptides having chiral five-membered ring
amino acid with two azido or triazole functional groups within the
sequence of Aib residues

Makoto Oba^{a*}, Naomi Kawabe^b, Hiroomi Takazaki^b, Yosuke Demizu^c, Mitsunobu Doi^d, Masaaki
Kurihara^c, Hiroshi Suemune^b, Masakazu Tanaka^{a*}

^a*Graduate School of Biomedical Sciences, Nagasaki University, Nagasaki 852-8521, Japan*

^b*Graduate School of Pharmaceutical Sciences, Kyushu University, Fukuoka 812-8582, Japan*

^c*Division of Organic Chemistry, National Institute of Health Sciences, Tokyo 158-8501, Japan*

^d*Osaka University of Pharmaceutical Sciences, Osaka 569-1094, Japan*




Fig. S1. CD spectra of a) pentapeptides (R = azide: **1**; R = triazole: **2**), b) pentapeptides (R = azide: **3**; R = triazole: **4**), c) Boc-[(*R,R*)-Ac₅c^{dN₃}-(Aib)₂]_n-OEt (n = 1, 3 mer: **5**; n = 2, 6 mer: **6**; n = 3, 9 mer: **7**), d) azide compounds (pentapeptide **1**; pentapeptide **3**; amino acid **8**), and d) triazole compounds (pentapeptide **2**; pentapeptide **4**; amino acid **9**) in TFE solution.