

1

2

3

4 **Protective role of interferon regulatory factor3-mediated signaling**  
5 **against prion infection**

6

7

8

9 Running title: Role of IRF3 in prion infection

10

11 Daisuke Ishibashi<sup>1\*</sup>, Ryuichiro Atarashi<sup>1,3</sup>, Takayuki Fuse<sup>1</sup>, Takehiro Nakagaki<sup>1</sup>,  
12 Naohiro Yamaguchi<sup>1</sup>, Katsuya Satoh<sup>1</sup>, Kenya Honda<sup>2</sup> and Noriyuki Nishida<sup>1,4</sup>

13

14 1. Department of Molecular Microbiology and Immunology, Nagasaki University Graduate  
15 School of Biomedical Sciences, 1-12-4 Sakamoto, Nagasaki 852-8523, Japan.

16 2. Department of Immunology, Graduate School of Medicine and Faculty of Medicine,  
17 University of Tokyo, Hongo 7-3-1, Bunkyo-ku, Tokyo 113-0033, Japan.

18 3. Nagasaki University Research Centre for Genomic Instability and Carcinogenesis, Nagasaki,  
19 Japan.

20 4. Global Centers of Excellence Program, Nagasaki University, Nagasaki, Japan.

21 \*: Corresponding author: Daisuke Ishibashi,

22 Department of Molecular Microbiology and Immunology, Nagasaki University Graduate School  
23 of Biomedical Sciences, 1-12-4 Sakamoto, Nagasaki 852-8523, Japan.

24 Tel.: +81-95-819-7059

25 Fax: +81-95-819-7060

26 E-mail: dishi@nagasaki-u.ac.jp

27

28 Keywords: prion, innate immunity, interferon regulatory factor 3 (IRF3)

29 **ABSTRACT**

30 Abnormal prion protein (PrP<sup>Sc</sup>) generated from the cellular isoform of PrP (PrP<sup>C</sup>) is  
31 assumed to be the main or sole component of the pathogen, called prion, of  
32 transmissible spongiform encephalopathies (TSE). Because PrP is a host-encoded  
33 protein, acquired immune responses are not induced in TSE. Meanwhile, activation of  
34 the innate immune system has been suggested to partially block the progression of TSE;  
35 however, the mechanism is not well understood. To further elucidate the role of the  
36 innate immune system in prion infection, we investigated the function of interferon  
37 regulatory factor 3 (IRF3), a key transcription factor of the MyD88-independent type I  
38 interferon (IFN) production pathway. We found that IRF3-deficient mice exhibited  
39 significantly earlier onset with three murine TSE strains, namely, 22L, FK-1, and mBSE  
40 following intraperitoneal transmission, when compared with wild-type controls.  
41 Moreover, overexpression of IRF3 attenuated prion infection in the cell culture system,  
42 while PrP<sup>Sc</sup> was increased in prion-infected cells treated with small interfering RNAs  
43 (siRNA) against IRF3, suggesting that IRF3 negatively regulates PrP<sup>Sc</sup> formation. Our  
44 findings provide new insight into the role of the host innate immune system in the  
45 pathogenesis of prion diseases.

46

47

48 **INTRODUCTION**

49 Transmissible spongiform encephalopathies (TSE) are fatal zoonoses, and  
50 include Creutzfeldt–Jakob disease (CJD) in humans, and scrapie and bovine spongiform  
51 encephalopathy (BSE) in animals. All exhibit the three major histopathological features  
52 of spongiform change, neuronal loss and gliosis in the central nervous system (CNS)  
53 (30). The infectious agent, prion, is considered not to possess its own genome and to be  
54 composed mainly of the proteinase K (PK)-resistant and  $\beta$ -sheet-rich abnormal isoform  
55 of prion protein, designated PrP<sup>Sc</sup>, which is generated by conformational conversion of  
56 the normal form of PrP (PrP<sup>C</sup>) (43). In contrast to conventional pathogens, such as  
57 bacteria and virus, acquired immunity against prion infection is not elicited, probably  
58 because PrP is a host-encoded protein, resulting in immunotolerance to PrP<sup>Sc</sup> (1).

59 Prior to activation of acquired immune responses, the invasion of pathogens,  
60 including bacteria and viruses, is first recognized by the innate immune system, with the  
61 switching on of the cellular defense system leading to the production of cytokines and  
62 interferons (IFNs). The innate immune responses are initiated through both toll-like  
63 receptors (TLRs) (2) and intracellular sensor molecules such as retinoic acid inducible  
64 gene-I (RIG-I) and melanoma differentiation associated gene-5 (MDA5), each of which  
65 recognizes specific components of foreign pathogens, namely pathogen-associated  
66 molecular patterns (PAMPs) (20). In addition, the innate immunity is the main system  
67 contributing to inflammation caused by microbial infection or tissue damage (3, 8).  
68 Since gliosis, a major characteristic of TSE, is thought to be a kind of inflammatory  
69 response, it is reasonable to assume that innate immunity may play a role in the  
70 pathogenesis of TSE. Indeed, it was reported that pretreatment with complete Freund's  
71 adjuvant (CFA) (39) or unmethylated CpG DNA (35), both of which activate innate  
72 immunity through TLRs, delays the onset of TSE in mice inoculated with  
73 mouse-adapted scrapie prion, suggesting that activation of innate immunity is protective  
74 against prion infection. In contrast, deletion of MyD88 gene, which is an essential  
75 intracellular signal transducer in all TLRs except for TLR3, has been shown not to  
76 significantly affect incubation time in the same mouse scrapie model (29). Thus,  
77 MyD88-dependent signaling pathways are unlikely to be implicated in prion infection in  
78 the absence of forced activation of innate immune responses by conventional PAMPs (2,  
79 20). On the other hand, mice that possess a non-functioning mutation of TLR4, which  
80 activates not only MyD88-dependent but also MyD88-independent (also called

81 TRIF-dependent) pathway, develop scrapie earlier than control mice (36). Accordingly,  
82 it is suggested that blockade of TLR4 signaling pathway accelerates the progression of  
83 TSE. Nonetheless, the effects of the innate immune system on prion infection remain  
84 controversial and have not been fully clarified.

85 We focus on interferon regulatory factor 3 (IRF3), which is a key transcription  
86 factor of the MyD88-independent pathway that has an essential role in the type I IFN  
87 response to microbial infection, and whose deficiency in mice leads to susceptibility to  
88 many viruses (19). In this study, we investigated the role of IRF3 in prion infection  
89 using IRF3-deficient mice and prion-susceptible cell lines.

90

## 91 **RESULTS**

### 92 **Prion infection is accelerated in IRF3-deficient mice.**

93 To clarify the significance of IRF3-dependent signaling pathways in prion infection *in*  
94 *vivo*, we peripherally inoculated 22L strain into IRF3-gene knockout mice (IRF3<sup>-/-</sup>)  
95 (33) and control wild-type (C57BL/6) mice. When the mice were challenged with 10<sup>-3</sup>  
96 dilution of 22L inoculum by the intraperitoneal (i.p.) route, the IRF3<sup>-/-</sup> mice showed  
97 significantly abbreviated survival periods (257 ± 8 days, p < 0.001) compared with  
98 those of the control mice (281 ± 15 days) (Table 1 and Fig. 1A). To further investigate  
99 the protective effect of IRF3 on prion infection, the mice were challenged with 10<sup>-3</sup>  
100 dilution of mouse-adapted BSE (mBSE) or Fukuoka 1 (FK-1) strain by the i.p. route.  
101 IRF3<sup>-/-</sup> mice showed shorter survival periods of mBSE (335 ± 37 versus 380 ± 33 days,  
102 p < 0.05), and FK-1 (251 ± 31 versus 321 ± 24 days, p < 0.01) compared with control  
103 mice (Table 1). The shortening of survival periods in the IRF3<sup>-/-</sup> mice is unlikely to be  
104 due to developmental defects in the brains or lymphoreticular organs, because IRF3<sup>-/-</sup>  
105 mice were shown to have normal lymphocyte populations in the thymus and spleen (33).  
106 Moreover, immunostaining with anti-follicular dendritic cell (FDC) antibody (CNA.42)  
107 (31) was performed to compare the FDC population in the spleen between IRF3<sup>-/-</sup> and  
108 wild-type mice. Staining reactions were similar in the two groups (Fig. 4B).

109 We examined for the presence of PrP<sup>Sc</sup> in the brain tissues of terminal-stage  
110 infected with 22L prion strain by Western blotting. The levels of PrP<sup>Sc</sup> in IRF3<sup>-/-</sup> mice  
111 at 32 weeks post-infection (w.p.i.) were equivalent to those of wild-type mice at 40 w.p.i.  
112 (Fig. 1B). Moreover, no significant differences were observed between wild-type mice  
113 and IRF3<sup>-/-</sup> mice in the accumulation of PrP<sup>Sc</sup> in the lesion profiles of

114 PrP-immunostaining (Fig. 1C). Because spongiform changes and gliosis are common  
115 characteristics of prion diseases, brain sections including the cerebral cortex (Cx),  
116 hippocampus (Hi), thalamus (TH), cerebellum (CE), and pons (Po) from 22L-inoculated  
117 mice were examined by histologically and subjected to immunohistochemical analysis  
118 using anti-Iba-1 antibodies for microgliosis (Fig. 3A) or anti-GFAP for astrogliosis (Fig.  
119 3B). The severity and distribution of vacuolation and glial activation in the IRF3<sup>-/-</sup> mice  
120 at 32 w.p.i. were indistinguishable from those in the wild-type mice at 40 w.p.i. (Fig 2  
121 and 3), while IRF3<sup>-/-</sup> mice displayed significantly increased vacuolation (Fig. 2) and  
122 astrogliosis (Fig. 3B) in the Cx and CE at 25 w.p.i.. Collectively, these results suggest  
123 that the progression of TSE following i.p. transmission is accelerated in IRF3<sup>-/-</sup> mice,  
124 although genetic elimination of IRF3 does not affect the final neuropathological  
125 outcome. Furthermore, as shown in Table 2 and Fig. 4A, the deposition of PrP<sup>Sc</sup> in the  
126 white pulp region of the spleens from the IRF3<sup>-/-</sup> mice was detectable in 1/5 at 2 w.p.i.,  
127 4/5 at 5 w.p.i., and 5/5 at 8 w.p.i., whereas none of the wild-type mice was positive at  
128 the same timepoints. These observations indicate that the rate of accumulation of PrP<sup>Sc</sup>  
129 in the spleen was enhanced in the IRF3<sup>-/-</sup> mice.

130

### 131 **IRF3-dependent pathway is protective against prion infection in cell culture.**

132 We tested whether overexpression of IRF3 could affect the production of PrP<sup>Sc</sup> in the  
133 cell culture models. The level of PrP<sup>C</sup> was not affected by the transient expression of the  
134 genes in uninfected N2a58 cells (data not shown). PrP<sup>Sc</sup> was significantly decreased by  
135 overexpression of IRF3 in the 22L-N2a58 cells (Fig. 5A). We confirmed that the  
136 activated form of IRF3 (phosphorylated at Ser396 of IRF3) increases in a  
137 dose-dependent manner after transfection of the IRF3 gene in both 22L-N2a58 cells  
138 (Fig. 5A) and uninfected N2a58 cells (data not shown), indicating that the up-regulation  
139 of IRF3-phosphorylation seen in the Fig. 5A is most likely due to an increase in the level  
140 of IRF3 protein after transfection.

141 To investigate the effect of down-regulation of IRF3 in the 22L-N2a58 cells,  
142 we performed knockdown experiments using small interfering RNAs (siRNA). IRF3  
143 expression was significantly decreased by two types of siRNA against IRF3, whereas  
144  $\beta$ -actin expression, as the internal standard, was not changed (Fig. 5B). Application of  
145 siRNA did not influence the expression of PrP<sup>C</sup> in N2a58 cells (data not shown),  
146 whereas the level of PrP<sup>Sc</sup> was increased in 22L-N2a58 cells treated with siRNA against

147 IRF3 (Fig. 5B). These data suggest that IRF3 has an inhibitory effect on the production  
148 of PrP<sup>Sc</sup> in the 22L-N2a58 cells.

149 To further evaluate the protective effect of IRF3, we established cell clones  
150 stably expressing HA-tagged IRF3 using another mouse PrP-overexpressed N2a cell  
151 clone (designated N2a75). Several HA-IRF3-negative and -positive clones were isolated  
152 by selection for resistance to hygromycin. HA-tagged IRF3 was expressed in clones A4,  
153 C1, and H3, which were accompanied by an increase in total IRF3 protein, while clones  
154 A1 and E1 were negative (Fig. 5C). After incubation with 22L-infected BH, the cell  
155 clones were subcultured for 5 passages, and analyzed by Western blotting with anti-PrP  
156 antibodies. The levels of PrP<sup>Sc</sup>/PrP<sup>C</sup> ratio were inversely correlated with the levels of  
157 IRF3/beta-actin ratio (Fig. 5C), indicating that enhanced expression of IRF3 effectively  
158 blocks new prion infection.

159

## 160 **DISCUSSION**

161 In the present study, we found that a genetic deficiency of IRF3 accelerates the  
162 progression of TSE following i.p. transmission in mice and the accumulation rate of  
163 PrP<sup>Sc</sup> in the spleen is increased in the IRF3<sup>-/-</sup> mice. Furthermore, we demonstrated that  
164 IRF3 has the inhibitory effect on the PrP<sup>Sc</sup> accumulation and the levels of IRF3 are  
165 inversely correlated with resistance to prion infection in cell culture.

166 IRF-3 is known to be constitutively expressed in many tissues and cells (6, 22,  
167 45). Indeed, we confirmed the expression of IRF3 in brains (data not shown) and the  
168 N2a58 cells (Fig. 5). Furthermore, not only glial cells but also neurons express most  
169 innate immunity-related genes and produce type I IFN in response to virus infection  
170 (11). Although the role of IRF3 in prion propagation into the CNS is still unclear, we  
171 speculate that an absence of IRF-3 signaling leads to increased prion replication not  
172 only in peripheral tissues but also in the CNS. It would be of great value to examine this  
173 further using neuron-specific IRF3-disrupted mice or neuron-specific IRF3-expressing  
174 mice.

175 It was reported, in prion infection, that genetic disturbance of TLR4 (36) or  
176 IL-10 (41) leads to shorter incubation periods of prion infection. Since these,  
177 respectively, are an upstream and a downstream factor of IRF3-mediated pathway, the  
178 findings may be due in part to functional changes in the IRF3-mediated signaling.

179 Based on these results, two hypothetical models are proposed to explain the inhibitory

180 effect of IRF3 on the prion infection. The first is that MyD88-independent “pattern  
181 recognition receptors (PRRs)” such as TLR3, TLR4 or RIG-I/MDA5 might recognize  
182 prion, and the resulting activation of IRF3 could induce various IRF3-responsive genes  
183 that may participate in the protective effect. The fact that the *in vivo* administration of  
184 IFNs, a representative of the IRF3-responsive genes, previously failed to show  
185 inhibitory effects on TSE (13, 16) suggests that IRF3-responsive genes other than IFNs  
186 may be important for the inhibitory effect of IRF3 on prion infection. Of note, the  
187 protective effect of IRF3 against several viruses has been suggested to be largely  
188 independent of the production of type I-IFN, and probably responsible for the anti-viral  
189 actions of specific IRF3-responsive genes (10, 18, 21). Peritoneal macrophages from  
190 wild-type mice moderately induced TNF- $\alpha$  or IL-6 following exposure to  
191 PrP<sup>Sc</sup>-mimicking PrP peptides (PrP106–126 or PrP118–135), whereas TLR4  
192 signaling-mutant mice were impaired in their ability to produce these cytokines (36),  
193 supporting in part the hypothesis that some PRRs may sense PrP<sup>Sc</sup> as a sort of PAMP.  
194 On the other hand, it should be noted that the MyD88-independent pathway activates  
195 both NF- $\kappa$ B and IRF3. Although the induction of proinflammatory cytokines essentially  
196 depends upon NF- $\kappa$ B, it was unclear whether the activation of IRF3 was induced by  
197 these PrP peptides. In fact, the hallmarks of IRF3 activation, such as phosphorylation,  
198 dimerization and cytoplasm-to-nucleus translocation of IRF3 in 22L-N2a58 cells were  
199 not detected (data not shown). Moreover, it was previously reported that IFNs were not  
200 detected in the serum, spleens, or brains of mice infected with scrapie (44). In addition,  
201 IFN- $\beta$  mRNA does not increase in the brains of CJD patients (7) or mice infected with  
202 ME7 prion strain (14). Hence, these results argue against the notion that the  
203 IRF3-mediated signaling is activated by prion infection, but it remains to be determined  
204 whether transient and weak responses are evoked at an early phase in the infection. The  
205 question as to whether IRF3-mediated signaling directly suppresses the production of  
206 PrP<sup>Sc</sup> or increases its degradation also remains open.

207 Another explanation is that prion infection itself may have little effect on the  
208 pathway, but that the basal activity of IRF3 may have some degree of inhibitory effect  
209 on prion propagation. It has been reported that IRF3 can be activated not only by  
210 viruses but also by multiple activators such as cellular stress and DNA damage (24)  
211 (34). Accordingly, it is possible that constitutive activation of IRF3, albeit at a low level,  
212 occurs in the brain even in the absence of a pathogen. This notion is further supported

213 by the fact that constitutive, weak IFN-signaling in the absence of viral infection plays a  
214 role in modifying cellular responsiveness in the immune and other biological systems  
215 (38, 40). Accumulating evidence indicates that many viruses have evolved to evade the  
216 innate immune system, including IRF3-mediated signaling (15, 23). For instance, an  
217 active mutant of IRF3 has been reported to exert a markedly suppressive effect on  
218 cellular HIV-1 infection and administration of poly I:C potently inhibits HIV-1  
219 replication in microglia through a pathway requiring IRF3. Nonetheless, HIV-1 itself  
220 does not activate IRF3 but rather decreases IRF3 protein in HIV-1 infected cells (12, 37).  
221 Likewise, prion infection might disturb the activation of IRF3, even though prion is  
222 considered to be largely composed of PrP<sup>Sc</sup>. We are currently investigating this  
223 possibility. Furthermore, an analogy can be made between the role of IRF3 in prion  
224 infection and that of IL-10. The levels of IL-10 are not increased in the brains of  
225 scrapie-infected mice (14, 42), whereas IL-10 knockout mice are highly susceptible to  
226 the development of scrapie (41).

227 In conclusion, we have shown that IRF3, a key transcription factor of the  
228 MyD88-independent pathways, operates in the host defense machinery against prion  
229 infection. The findings provide new insight into understanding of the innate immunity  
230 to prion infection.

231

## 232 **MATERIALS AND METHODS**

233

### 234 **Reagents and Antibodies**

235 The anti-PrP polyclonal mouse antiserum (SS) has been described previously (5). M20  
236 is an affinity purified goat polyclonal antibody recognizing the C-terminus of mouse  
237 PrP (Santa Cruz Biotechnology, Inc., CA, USA). Anti-mouse IRF3 (Santa Cruz  
238 Biotechnology) and anti-mouse phospho-IRF3 (Ser396) (Cell Signaling Technology,  
239 Japan) were rabbit polyclonal antibodies, and anti-mouse  $\beta$ -actin (Sigma Aldrich, St.  
240 Louis, MO, USA) was a mouse monoclonal antibody. Horseradish peroxidase  
241 (HRP)-conjugated anti-goat immunoglobulin G antibody (Santa Cruz Biotechnology),  
242 anti-mouse and anti-rabbit IgG antibodies (Amersham Pharmacia Biotech AB, Uppsala,  
243 Sweden) were used for Western blotting.

244

### 245 **Cell cultures**



246 The mouse neuroblastoma cell line N2a was purchased from the American Type Culture  
247 Collection (ATCC CCL131), and N2a58 cells overexpressing mouse PrP prepared from  
248 N2a were transfected with a plasmid carrying wild-type mouse prnp cDNA (PrP-a  
249 genotype, codons 108L and 189T) (27). Prion infected cells, 22L-N2a58, were produced  
250 as previously described (27). After limiting dilution, several PrP<sup>Sc</sup>-positive clones were  
251 isolated. The cell clones producing the highest level of PrP<sup>Sc</sup> were used for subsequent  
252 studies. The 22L-N2a58 cells stably expressed PrP<sup>Sc</sup> for over 50 passages. The cells  
253 were cultured in DMEM (Sigma) containing 10% heat-inactivated fetal bovine serum  
254 and penicillin-streptomycin (Invitrogen, Carlsbad, CA, USA), and split every 3 days at  
255 a 1:10 ratio. All cultured cells were maintained at 37 °C in 5% CO<sub>2</sub> in the biohazard  
256 prevention area of the authors' institution.

257

#### 258 **Plasmid and siRNA**

259 The mammalian expression pUNO vector contains a strong and ubiquitous composite  
260 promoter designated EF1 $\alpha$ /HTLV. In this experiment, we inserted mouse IRF3 cDNA  
261 into multiple cloning sites of the pUNO vector (Invivogen, San Diego, CA, USA.). The  
262 HA-tagged mouse IRF3 cDNA was inserted into pcDNA3.1 vector (Invitrogen). The  
263 plasmids were introduced by Lipofectoamine LTX (Invitrogen) in the prion-infected  
264 cells and incubated for 48 h. The small interfering RNAs (siRNAs) were purchased  
265 from QIAGEN, Hilden, Germany. Two specific siRNA-targeted sequences (product  
266 IDs: SI00210770 and SI00210784) were used for IRF3 (according to GenBank  
267 accession No: NM016849), 5'- ACA GGT GGT GAT GGT TGG CAA -3' and 5'- GAC  
268 CCT TAT GAC CCT CAT AAA -3'. For the negative control, a siRNA (product ID:  
269 1022076) targeting 5'- AAT TCT CCG AAC GTG TCA CGT-3' was used. The siRNAs  
270 were introduced into cells using Fugene 6 (Roche Diagnostics, Mannheim, Germany)  
271 and incubated for 48 h.

272

#### 273 **Western blotting**

274 Samples were lysed with Triton-DOC lysis buffer (50 mM Tris-HCl [pH 7.5] containing  
275 150 mM NaCl, 0.5% Triton X-100, 0.5% sodium deoxycholate, 2 mM EDTA, and  
276 protease inhibitors (Nakarai Tesque, Inc., Japan), for 30 min at 4 °C. After 1 min of  
277 centrifugation at 5000 $\times$ g, the supernatant was collected and its total protein  
278 concentration was measured using the BCA protein assay kit (Pierce, Rockford, IL). To

279 detect PrP<sup>Sc</sup>, the protein concentration was adjusted to 10 mg/ml, and samples were  
280 digested with 20 µg/ml of proteinase K (PK, Sigma) at 37 °C for 30 min, and boiled for  
281 5 min with sodium dodecyl sulfate (SDS) loading buffer (50 mM Tris-HCl, pH 6.8,  
282 containing 5% glycerol, 1.6% SDS, and 100 mM dithiothreitol) and subjected to  
283 SDS–polyacrylamide gel electrophoresis. The proteins were transferred onto an  
284 Immobilon-P membrane (Millipore, MA, USA) in transfer buffer containing 15%  
285 methanol at 300 mA for 1 h, and the membrane was blocked with 5 % nonfat dry milk  
286 in TBST (10 mM Tris-HCl [pH 7.8], 100 mM NaCl, 0.1% Tween 20) for 1 h at room  
287 temperature and reacted with diluted primary antibodies. Immunoreactive bands were  
288 visualized by HRP-conjugated secondary antibodies, using an enhanced  
289 chemiluminescence system (Amersham). The detailed methods have previously been  
290 described (4). To quantify the signals, we measured the intensity of each band using the  
291 NIH image J software.

292

#### 293 **Establishment of stably IRF3-overexpressing cells and in vitro 22L scrapie** 294 **infection experiments**

295 To establish cell lines stably expressing IRF3, pcDNA3.1 plasmid containing  
296 HA-tagged IRF3 gene was transfected using Fugen 6 (Roche) into PrP<sup>C</sup>-overexpressing  
297 N2a cells (N2a 75 cells) and cultured for 48 h. After the cells were treated with 350  
298 µg/mL hygromycin (HygroGold, Invivogen) in culture medium for 4 days, the  
299 drug-resistant colonies were isolated. We used the HygroGold-selected N2a 75 cells  
300 transfected with empty vector as a negative control. Then, the cells were infected with  
301 22L scrapie strain (final concentration, 0.02 or 0.2% brain homogenate) in a 6-well  
302 culture plate for 48 h, and subsequently passaged into a 75 cm<sup>2</sup>-flask. Once confluent,  
303 the sub-cultures were diluted 10-fold (27). After treatment with 40 µg/mL PK,  
304 immunoblotting was done to detect PrP<sup>Sc</sup>.

305

#### 306 **In vivo scrapie infection experiments**

307 Four-week-old wild-type (+/+) and IRF3 knockout (-/-) male mice of the same  
308 C57BL/6-derived genetic background were intraperitoneally (i.p.) inoculated with 100  
309 µL of 10<sup>-3</sup> dilution of brain homogenate (BH) from mice terminally infected with 22L,  
310 Fukuoka-1 (FK-1) or mouse-adapted BSE (mBSE) strain (28). The IRF3<sup>-/-</sup> mice were  
311 obtained from Dr. T. Taniguchi's group (Tokyo University) (33). As a negative

312 (mock-infected) control, age- and strain-matched mice were inoculated i.p. with normal  
313 mouse BH. The spleens and brains of the mice were removed at 1, 2, 5, 8, 25 weeks  
314 post-infection (w.p.i.) and at the terminal stage of disease. Animals were cared for in  
315 accordance with the Guidelines for Animal Experimentation of Nagasaki University.

316

### 317 **Histopathology and Immunohistochemical Staining**

318 The spleen and brain tissues were fixed in 4% paraformaldehyde, and 5- $\mu$ m paraffin  
319 sections prepared on PLL coat slides with microtome. To measure vacuolation in brain,  
320 the tissue sections were stained with hematoxylin and eosin (HE). In PrP<sup>Sc</sup> staining,  
321 after deparaffinization and rehydration, the sections were pretreated by hydrated  
322 autoclaving at 121 °C for 15 min in 1 or 1.2 mM hydrochloric acid (17), followed by  
323 immersion in 90% formic acid for 5 min (25) to enhance PrP visualization, according to  
324 the protocol described by Brown et al (9). Endogenous peroxidase activity was inhibited  
325 with 0.3% hydrogen peroxidase in methanol for 30 min. Nonspecific binding sites of  
326 the primary antibody SAF32 (SPI-BIO, Montigny le Bretonneux, France) was blocked  
327 by preincubation in normal rabbit serum at 1:20 (Dako, Glostrup, Denmark) for 30 min,  
328 and then optimally titrated and diluted to 1:5000 and 1:500 (26, 32). The negative  
329 control sections were incubated with normal mouse IgG1 and IgG2b serum (DAKO)  
330 and then exposed to the primary antibodies overnight at room temperature. For  
331 determining follicular dendritic cell population in the spleen, we used anti-FDC  
332 antibody CNA.42 (DAKO) (31) and normal mouse IgM serum (DAKO) as a negative  
333 control for primary antibodies, and Histofine mouse stain kit (Nichirei biosciences, Inc.,  
334 Jpn) for the secondary antibody. In glial staining, primary antibodies of anti-GFAP (glial  
335 fibrillary acidic protein) (DAKO) for the detection of activated astrocyte, and anti-Iba-1  
336 (WAKO, Japan) for activated microglia were used. The secondary antibodies were  
337 Envision-mouse or rabbit HRP (DAKO) used at 1:200 for 1 h. Finally, the samples were  
338 stained with 0.025% 3,3' diaminobenzidine (DAB, Dojindo Lab, Japan) to visualize the  
339 reaction product, and counterstained with hematoxylin. The pattern of vacuolation,  
340 PrP<sup>Sc</sup> deposits and gliosis were examined in 5 areas, namely the cortex, hippocampus,  
341 thalamus, cerebellum, and Pons. In semi-quantitative evaluation of spongiosis and  
342 gliosis, lesion severity of vacuolation as spongiform degeneration was scored on a 0–5  
343 scale (non-detectable, a few, mild, moderate, severe and status spongiosus). PrP<sup>Sc</sup>  
344 deposit was scored on a 0–4 scale (non-detectable, a few, mild, moderate and severe).

345 Microgliosis and astrogliosis were scored on a 0–3 scale (non-detectable, mild,  
346 moderate and severe) and the values for each brain region were averaged.

347

#### 348 **Statistical analysis**

349 The Student's t-test and Mann-Whitney U-test were used for comparison between two  
350 groups, and the one-way ANOVA followed by the Tukey-Kramer test, for multiple  
351 comparisons. The log rank test was used to analyze mortality of prion-infected mice.  
352 The correlation between parameters was determined by simple regression analysis and  
353 Pearson's correlation coefficient test. Statistical analysis of all data was performed using  
354 the Statcel 2 on Excel and GraphPad Prism software.

355

#### 356 **Acknowledgments**

357 We thank Dr. Tadatsugu Taniguchi for the gift of IRF3 knockout mice; and Drs. Hitoki  
358 Yamanaka, Takehiro Matsubara, Kazunori Sano, for helpful discussions and critical  
359 assessment of the manuscript; and Mari Kudo, Ayumi Yamakawa and Atsuko Matsuo,  
360 for technical assistance. This work was supported in part by the global COE Program  
361 (F12); a Grant-in-Aid for Young Scientists (B) (No. 22790955) from the Ministry of  
362 Education, Culture, Sports, Science, and Technology of Japan; a grant for BSE research,  
363 and a grant-in-aid of the Research Committee of Prion disease and Slow Virus Infection,  
364 from the Ministry of Health, Labor and Welfare of Japan.

365

366 **FIGURE LEGENDS**

367 **Fig. 1.** Acceleration of prion pathogenesis in IRF3<sup>-/-</sup> mice.

368 (A) Survival curves in wild-type (+/+) (n=9, circle) and IRF3 knockout (-/-) (n=5,  
369 triangle) mice after i.p. inoculation of 10<sup>-3</sup> dilution of 22L-BH are plotted. The  
370 difference between the two groups is statistically significant (p < 0.05, Logrank test).  
371 (B) Accumulation of PrP<sup>Sc</sup> in the brain tissues from wild-type or IRF3<sup>-/-</sup> mice was  
372 analyzed by Western blotting. Molecular mass markers are indicated in kilodaltons  
373 (kDa) on the left side of each panel. (C) PrP<sup>Sc</sup> deposits were similarly stained in cortex  
374 area of wild-type (+/+) and IRF3 knockout (-/-) mice at the terminal stages (upper  
375 panels). There were no significant differences between wild-type and IRF3<sup>-/-</sup> mice in  
376 the staining levels of PrP<sup>Sc</sup> in the five brain regions: cortex (Cx), hippocampus (Hi),  
377 thalamus (TH), cerebellum (CE) and Pons (Po) (lower graph). Scale-bars equal 25 μm.  
378 All data are representative of at least three mice.

379

380 **Fig. 2.** Comparison of spongiform change between wild-type and IRF3<sup>-/-</sup> mice after  
381 22L prion infection.

382 (A) Sections of the cortex (Cx) and cerebellum (CE) stained with hematoxylin and eosin  
383 from wild-type (+/+) and IRF3 knockout (-/-) mice at 0, 25 w.p.i. and terminal stages  
384 after i.p. inoculation of 10<sup>-3</sup> dilution of 22L-BH are shown. Scale-bars equal 25 μm. (B)  
385 Vacuolation scores in the same five brain regions as Fig. 1B were compared between  
386 the prion-inoculated wild-type (circle) and IRF3<sup>-/-</sup> (triangle) mice at 25 w.p.i. or the  
387 terminal stages. Statistical significance was determined using a two-tailed Student's  
388 *t*-test. \*\*\*p < 0.001, \*p < 0.05 compared with wild-type mice. Error bars indicate SEM.  
389 These results are representative of at least three mice.

390

391 **Fig. 3.** Histopathological examination of gliosis in the brains of prion-infected mice.

392 (A) Immunohistochemical staining for Iba-I with hematoxylin counterstaining was  
393 performed. The sections of cortex from wild-type (+/+) and IRF3 knockout (-/-) mice at  
394 0, 25 w.p.i. and terminal stages are shown (upper panels). Lesion profiles of  
395 microgliosis in the same five brain regions as Fig. 1B were compared between the  
396 prion-inoculated wild-type (circle) and IRF3<sup>-/-</sup> (triangle) mice at the terminal stages  
397 (lower graph). No significant differences were observed between the two groups at 25

398 w.p.i. or at the terminal stage. (B) Immunostaining for GFAP with hematoxylin  
399 counterstaining (upper panels), and lesion profiles of astrogliosis (lower graph).  
400 Significant differences were observed between the two groups in Cx and CE regions at  
401 25 w.p.i., but not at the terminal stage. Statistical significance was determined using a  
402 two-tailed Student's *t*-test. \*\**p* < 0.01, \**p* < 0.05 compared with wild-type mice. Error  
403 bars indicate SEM. Scale-bars equal 25  $\mu$ m. All results are representative of at least  
404 three mice.

405

406 **Fig. 4.** Early detection of PrP<sup>Sc</sup> deposits in the spleen of prion-inoculated IRF3<sup>-/-</sup> mice.  
407 (A) Accumulation of PrP<sup>Sc</sup> in the spleens was analyzed by immunohistochemistry  
408 (hydrophobic autoclaving method) at 5, 8 w.p.i. and terminal stages after i.p. inoculation  
409 of 10<sup>-3</sup> dilution of 22L-BH. Scale-bars equal 25  $\mu$ m. (B) The sections of spleen from  
410 wild-type (+/+) and IRF3 knockout (-/-) 5-week-old mice were stained with anti-FDC  
411 antibody (CNA.42). IgM is a negative control for the primary antibody. Scale-bars equal  
412 20  $\mu$ m.

413

414 **Fig. 5.** Inhibitory effect of IRF3 on PrP<sup>Sc</sup> replication in cell culture.

415 (A) Plasmids (pUNO) containing IRF3 gene were transiently transfected into  
416 22L-N2a58 cells and incubated for 48 h. The panels on the left show PK-treated PrP as  
417 PrP<sup>Sc</sup> (upper), total IRF3 (middle) and phosphorylated IRF3 (lower) in the cells. Mock,  
418 plasmids without IRF3 gene. The numbers above the panels represent the amount ( $\mu$ g)  
419 of the plasmids used for transfection. C, untransfected cells for negative control. The  
420 levels of PrP<sup>Sc</sup> band intensity in the cells transfected with 4 $\mu$ g of the plasmids with or  
421 without IRF3 gene are expressed as a percentage compared with the control (right  
422 graph). The results in the graph are the mean  $\pm$  SD of three independent experiments.  
423 Asterisks indicate statistically significant differences (\*, *p* < 0.05). (B) The siRNAs of  
424 IRF3 were transfected into 22L-N2a58 cells and incubated for 48 h. The cells were  
425 subjected to Western blotting to detect PrP<sup>Sc</sup>,  $\beta$ -actin and IRF3. The band intensities of  
426 PrP<sup>Sc</sup> (middle graph) or IRF3/ $\beta$ -actin ratio (right graph) were quantified. Asterisks  
427 indicate statistically significant differences (\*, *p* < 0.05 and \*\*, *P* < 0.01). The data are  
428 representative of three independent experiments. (C) The effect of overexpression of  
429 HA-tagged IRF3 on new prion infection was analyzed using stable IRF3-overexpressing  
430 clones, A4, C1 and H3. A hygromycin-resistant but IRF3-negative clones, A1 and E1,

431 were used as the negative control. The top four panels show protein expression of PrP<sup>C</sup>,  
432 total IRF3, HA-tagged IRF3 and  $\beta$ -actin prior to infection in each clone. After  
433 incubation with 0.02 % of 22L-BH for 48 h and then 5 passages, PrP<sup>Sc</sup> levels in the  
434 clones were determined by Western blotting. The scatter diagram is indicative of a  
435 correlative relationship between the PrP<sup>Sc</sup>/PrP<sup>C</sup> ratio and the IRF3 expression ratio  
436 (right graph). A statistically significant ( $P < 0.001$ ) correlation ( $r = -0.8$ ) was observed  
437 between the PrP<sup>Sc</sup> and the IRF3/ $\beta$ -actin ratio. The coefficient of determination is shown  
438 as  $R^2$  values. All results are representative of at least three independent experiments,  
439 and each experiment was performed in triplicate.

440

441

#### 442 REFERENCES

443

- 444 1. **Aguzzi, A., and M. Polymenidou.** 2004. Mammalian prion biology: one  
445 century of evolving concepts. *Cell* **116**:313-327.
- 446 2. **Akira, S., and K. Takeda.** 2004. Toll-like receptor signalling. *Nat Rev*  
447 *Immunol* **4**:499-511.
- 448 3. **Akira, S., S. Uematsu, and O. Takeuchi.** 2006. Pathogen recognition  
449 and innate immunity. *Cell* **124**:783-801.
- 450 4. **Arima, K., N. Nishida, S. Sakaguchi, K. Shigematsu, R. Atarashi, N.**  
451 **Yamaguchi, D. Yoshikawa, J. Yoon, K. Watanabe, N. Kobayashi, S.**  
452 **Mouillet-Richard, S. Lehmann, and S. Katamine.** 2005. Biological and  
453 biochemical characteristics of prion strains conserved in persistently  
454 infected cell cultures. *J Virol* **79**:7104-7112.
- 455 5. **Atarashi, R., V. L. Sim, N. Nishida, B. Caughey, and S. Katamine.**  
456 2006. Prion strain-dependent differences in conversion of mutant  
457 prion proteins in cell culture. *J Virol* **80**:7854-7862.
- 458 6. **Au, W. C., P. A. Moore, W. Lowther, Y. T. Juang, and P. M. Pitha.** 1995.  
459 Identification of a member of the interferon regulatory factor family  
460 that binds to the interferon-stimulated response element and  
461 activates expression of interferon-induced genes. *Proceedings of the*  
462 *National Academy of Sciences of the United States of America*  
463 **92**:11657-11661.

- 464 7. **Baker, C. A., Z. Y. Lu, and L. Manuelidis.** 2004. Early induction of  
465 interferon-responsive mRNAs in Creutzfeldt-Jakob disease. *J*  
466 *Neurovirol* **10**:29-40.
- 467 8. **Beutler, B., Z. Jiang, P. Georgel, K. Crozat, B. Croker, S. Rutschmann,**  
468 **X. Du, and K. Hoebe.** 2006. Genetic analysis of host resistance:  
469 Toll-like receptor signaling and immunity at large. *Annu Rev*  
470 *Immunol* **24**:353-389.
- 471 9. **Brown, K. L., D. L. Ritchie, P. A. McBride, and M. E. Bruce.** 2000.  
472 Detection of PrP in extraneural tissues. *Microsc Res Tech* **50**:40-45.
- 473 10. **Daffis, S., M. A. Samuel, B. C. Keller, M. Gale, Jr., and M. S. Diamond.**  
474 2007. Cell-specific IRF-3 responses protect against West Nile virus  
475 infection by interferon-dependent and -independent mechanisms.  
476 *PLoS pathogens* **3**:e106.
- 477 11. **Delhaye, S., S. Paul, G. Blakqori, M. Minet, F. Weber, P. Staeheli, and**  
478 **T. Michiels.** 2006. Neurons produce type I interferon during viral  
479 encephalitis. *Proc Natl Acad Sci U S A* **103**:7835-7840.
- 480 12. **Doehle, B. P., F. Hladik, J. P. McNevin, M. J. McElrath, and M. Gale,**  
481 **Jr.** 2009. Human immunodeficiency virus type 1 mediates global  
482 disruption of innate antiviral signaling and immune defenses within  
483 infected cells. *J Virol* **83**:10395-10405.
- 484 13. **Field, E. J., G. Joyce, and A. Keith.** 1969. Failure of interferon to  
485 modify scrapie in the mouse. *J Gen Virol* **5**:149-150.
- 486 14. **Field, R., S. Champion, C. Warren, C. Murray, and C. Cunningham.**  
487 Systemic challenge with the TLR3 agonist poly I:C induces amplified  
488 IFNalpha/beta and IL-1beta responses in the diseased brain and  
489 exacerbates chronic neurodegeneration. *Brain Behav Immun*  
490 **24**:996-1007.
- 491 15. **Goodbourn, S., L. Didcock, and R. E. Randall.** 2000. Interferons: cell  
492 signalling, immune modulation, antiviral response and virus  
493 countermeasures. *J Gen Virol* **81**:2341-2364.
- 494 16. **Gresser, I., C. Maury, and R. L. Chandler.** 1983. Failure to modify  
495 scrapie in mice by administration of interferon or anti-interferon  
496 globulin. *J Gen Virol* **64 (Pt 6)**:1387-1389.




- 497 17. **Haritani, M., Y. I. Spencer, and G. A. Wells.** 1994. Hydrated autoclave  
498 pretreatment enhancement of prion protein immunoreactivity in  
499 formalin-fixed bovine spongiform encephalopathy-affected brain. *Acta*  
500 *Neuropathol* **87**:86-90.
- 501 18. **Hidmark, A. S., G. M. McInerney, E. K. Nordstrom, I. Douagi, K. M.**  
502 **Werner, P. Liljestrom, and G. B. Karlsson Hedestam.** 2005. Early  
503 alpha/beta interferon production by myeloid dendritic cells in response  
504 to UV-inactivated virus requires viral entry and interferon regulatory  
505 factor 3 but not MyD88. *Journal of virology* **79**:10376-10385.
- 506 19. **Hiscott, J.** 2007. Triggering the innate antiviral response through  
507 IRF-3 activation. *The Journal of biological chemistry*  
508 **282**:15325-15329.
- 509 20. **Honda, K., and T. Taniguchi.** 2006. IRFs: master regulators of  
510 signalling by Toll-like receptors and cytosolic pattern-recognition  
511 receptors. *Nat Rev Immunol* **6**:644-658.
- 512 21. **Honda, K., H. Yanai, H. Negishi, M. Asagiri, M. Sato, T. Mizutani, N.**  
513 **Shimada, Y. Ohba, A. Takaoka, N. Yoshida, and T. Taniguchi.** 2005.  
514 IRF-7 is the master regulator of type-I interferon-dependent immune  
515 responses. *Nature* **434**:772-777.
- 516 22. **Karpova, A. Y., P. M. Howley, and L. V. Ronco.** 2000. Dual utilization of  
517 an acceptor/donor splice site governs the alternative splicing of the  
518 IRF-3 gene. *Genes Dev* **14**:2813-2818.
- 519 23. **Katze, M. G., Y. He, and M. Gale, Jr.** 2002. Viruses and interferon: a  
520 fight for supremacy. *Nat Rev Immunol* **2**:675-687.
- 521 24. **Kim, T., T. Y. Kim, Y. H. Song, I. M. Min, J. Yim, and T. K. Kim.** 1999.  
522 Activation of interferon regulatory factor 3 in response to  
523 DNA-damaging agents. *The Journal of biological chemistry*  
524 **274**:30686-30689.
- 525 25. **Kitamoto, T., K. Ogomori, J. Tateishi, and S. B. Prusiner.** 1987. Formic  
526 acid pretreatment enhances immunostaining of cerebral and systemic  
527 amyloids. *Lab Invest* **57**:230-236.
- 528 26. **Liu, W. G., D. A. Brown, and J. R. Fraser.** 2003. Immunohistochemical  
529 comparison of anti-prion protein (PrP) antibodies in the CNS of mice

- 530 infected with scrapie. *J Histochem Cytochem* **51**:1065-1071.
- 531 27. **Nishida, N., D. A. Harris, D. Vilette, H. Laude, Y. Frobert, J. Grassi, D.**  
532 **Casanova, O. Milhavet, and S. Lehmann.** 2000. Successful  
533 transmission of three mouse-adapted scrapie strains to murine  
534 neuroblastoma cell lines overexpressing wild-type mouse prion protein.  
535 *J Virol* **74**:320-325.
- 536 28. **Nishida, N., S. Katamine, and L. Manuelidis.** 2005. Reciprocal  
537 interference between specific CJD and scrapie agents in neural cell  
538 cultures. *Science* **310**:493-496.
- 539 29. **Prinz, M., M. Heikenwalder, P. Schwarz, K. Takeda, S. Akira, and A.**  
540 **Aguzzi.** 2003. Prion pathogenesis in the absence of Toll-like receptor  
541 signalling. *EMBO Rep* **4**:195-199.
- 542 30. **Prusiner, S. B.** 1998. Prions. *Proc Natl Acad Sci U S A* **95**:13363-13383.
- 543 31. **Raymond, I., T. Al Saati, J. Tkaczuk, S. Chittal, and G. Delsol.** 1997.  
544 CNA.42, a new monoclonal antibody directed against a  
545 fixative-resistant antigen of follicular dendritic reticulum cells. *Am J*  
546 *Pathol* **151**:1577-1585.
- 547 32. **Rosicarelli, B., B. Serafini, M. Sbriccoli, M. Lu, F. Cardone, M.**  
548 **Pocchiari, and F. Aloisi.** 2005. Migration of dendritic cells into the  
549 brain in a mouse model of prion disease. *J Neuroimmunol*  
550 **165**:114-120.
- 551 33. **Sato, M., H. Suemori, N. Hata, M. Asagiri, K. Ogasawara, K. Nakao, T.**  
552 **Nakaya, M. Katsuki, S. Noguchi, N. Tanaka, and T. Taniguchi.** 2000.  
553 Distinct and essential roles of transcription factors IRF-3 and IRF-7 in  
554 response to viruses for IFN-alpha/beta gene induction. *Immunity*  
555 **13**:539-548.
- 556 34. **Servant, M. J., N. Grandvaux, and J. Hiscott.** 2002. Multiple signaling  
557 pathways leading to the activation of interferon regulatory factor 3.  
558 *Biochem Pharmacol* **64**:985-992.
- 559 35. **Sethi, S., G. Lipford, H. Wagner, and H. Kretzschmar.** 2002.  
560 Postexposure prophylaxis against prion disease with a stimulator of  
561 innate immunity. *Lancet* **360**:229-230.
- 562 36. **Spinner, D. S., I. S. Cho, S. Y. Park, J. I. Kim, H. C. Meeker, X. Ye, G.**


- 563        **Lafauci, D. J. Kerr, M. J. Flory, B. S. Kim, R. B. Kasczak, T.**  
564        **Wisniewski, W. R. Levis, G. B. Schuller-Levis, R. I. Carp, E. Park, and**  
565        **R. J. Kasczak.** 2008. Accelerated prion disease pathogenesis in  
566        Toll-like receptor 4 signaling-mutant mice. *J Virol* **82**:10701-10708.
- 567    37.    **Suh, H. S., M. L. Zhao, N. Choi, T. J. Belbin, C. F. Brosnan, and S. C.**  
568        **Lee.** 2009. TLR3 and TLR4 are innate antiviral immune receptors in  
569        human microglia: role of IRF3 in modulating antiviral and  
570        inflammatory response in the CNS. *Virology* **392**:246-259.
- 571    38.    **Takaoka, A., and T. Taniguchi.** 2003. New aspects of IFN-alpha/beta  
572        signalling in immunity, oncogenesis and bone metabolism. *Cancer Sci*  
573        **94**:405-411.
- 574    39.    **Tal, Y., L. Souan, I. R. Cohen, Z. Meiner, A. Taraboulos, and F. Mor.**  
575        2003. Complete Freund's adjuvant immunization prolongs survival in  
576        experimental prion disease in mice. *J Neurosci Res* **71**:286-290.
- 577    40.    **Taniguchi, T., and A. Takaoka.** 2001. A weak signal for strong  
578        responses: interferon-alpha/beta revisited. *Nat Rev Mol Cell Biol*  
579        **2**:378-386.
- 580    41.    **Thackray, A. M., A. N. McKenzie, M. A. Klein, A. Lauder, and R.**  
581        **Bujdoso.** 2004. Accelerated prion disease in the absence of  
582        interleukin-10. *J Virol* **78**:13697-13707.
- 583    42.    **Tribouillard-Tanvier, D., J. F. Striebel, K. E. Peterson, and B.**  
584        **Chesebro.** 2009. Analysis of protein levels of 24 cytokines in scrapie  
585        agent-infected brain and glial cell cultures from mice differing in prion  
586        protein expression levels. *J Virol* **83**:11244-11253.
- 587    43.    **Weissmann, C., M. Enari, P. C. Klohn, D. Rossi, and E. Flechsig.** 2002.  
588        Molecular biology of prions. *Acta Neurobiol Exp (Wars)* **62**:153-166.
- 589    44.    **Worthington, M.** 1972. Interferon system in mice infected with the  
590        scrapie agent. *Infect Immun* **6**:643-645.
- 591    45.    **Zhai, J., D. Gao, W. Liu, R. Hong, Y. Qin, H. Ouyang, Y. Kong, Y. Wang,**  
592        **Y. Xie, and J. Liu.** 2008. Characterization of a novel isoform of murine  
593        interferon regulatory factor 3. *Biochemical and biophysical research*  
594        *communications* **377**:384-388.
- 595

596



A




B





C











A



B














Table 1. Survival periods of prion-infected wild-type (+/+) and IRF3 knockout (-/-) mice.

Strains	Survival periods (days) <sup>a</sup>	
	+/+ (n) <sup>b</sup>	-/- (n) <sup>b</sup>
22L	281 ± 15 (9)	257 ± 8 (5) ***
BSE	380 ± 33 (7)	335 ± 37 (7) *
FK-1	321 ± 24 (4)	251 ± 31 (5) **

<sup>a</sup> Survival periods were shown as “ Average ± SD (days) ”.

<sup>b</sup> (n), number of mice.

Animals were intraperitoneally administrated with 10<sup>-3</sup> dilution of brain homogenate from prion-infected terminal mice. \*p < 0.05, \*\*p<0.01, \*\*\*p < 0.001 by student's t-test.

Table 2. PrP<sup>Sc</sup> positive rate in the spleens in mice with i.p. inoculation of 22L at the indicated time points.

Weeks post inoculation	PrP <sup>Sc</sup> positive/total number of mice <sup>a</sup>	
	+/+	-/-
1	0/5	0/1
2	0/5	1/5
5	0/5	4/5
8	0/5	5/5
Terminal	1/1	1/1

<sup>a</sup>The presence of PrP<sup>Sc</sup> accumulation in the spleen of prion-infected mice was determined by immunohistochemical staining.