

**Inhibitory Effect of Sulfated Polysaccharide Porphyran on Nitric Oxide Production
in Lipopolysaccharide (LPS)-Stimulated RAW264.7 Macrophages**

Zedong Jiang ¹, Yoichiro Hama ², Kenichi Yamaguchi ^{1,3}, and Tatsuya Oda ^{1,3,*}

¹*Graduate School of Science and Technology, Nagasaki University, 1-14 Bunkyo-machi, Nagasaki 852-8521,* ²*Applied Biochemistry and Food Sciences, Faculty of Agriculture, Saga University, 1 Honjo, Saga 840-8502, and* ³*Division of Biochemistry, Faculty of Fisheries, Nagasaki University, 1-14 Bunkyo-machi, Nagasaki 852-8521, Japan*

Running title: Inhibition of NO Production by Porphyran in RAW264.7 cells

Field: Biochemistry

Topic: Biochemistry General

*To whom correspondence should be addressed.

Fax: +81-95-819-2831, E-mail: t-oda@nagasaki-u.ac.jp

SUMMARY

Porphyran, extracted from an edible red alga (*Porphyra yezoensis*), is a sulfated polysaccharide with a wide variety of biological activities including anti-tumor, antioxidant, and immuno-modulating activities. In this study, we examined the effect of porphyran on nitric oxide (NO) production in mouse macrophage cell line RAW264.7 cells. Although no significant activity of porphyran to induce NO or tumor necrosis factor- α (TNF- α) production in RAW264.7 cells was observed at the concentration range tested (10-500 $\mu\text{g/ml}$), it was found for the first time that porphyran inhibited NO production and expression of inducible NO synthase (iNOS) in RAW264.7 cells stimulated with lipopolysaccharide (LPS). In the presence of 500 $\mu\text{g/ml}$ porphyran, NO production and expression of iNOS in LPS-treated RAW264.7 cells were completely suppressed. On other hand, porphyran showed only a marginal effect on the secretion of TNF- α from LPS-stimulated RAW264.7 cells. Electrophoretic mobility shift assay (EMSA) using infrared dye labeled oligonucleotide with nuclear factor-kappa B (NF- κB) consensus sequence suggested that porphyran inhibited the LPS-induced NF- κB activation. The LPS-inducible nuclear translocation of p65, and the phosphorylation and degradation of I κB - α were also inhibited by the pretreatment with porphyran. Our results obtained in *in vitro* analysis suggest that porphyran suppresses NO production in LPS-stimulated macrophages by the blocking of NF- κB activation.

Key words: Porphyran, Sulfated polysaccharide, Nitric oxide, iNOS, TNF- α , NF- κB

Abbreviations: LPS, lipopolysaccharide; MAP kinase, mitogen-activated protein kinase; DMEM, Dulbecco's modified Eagle's minimal essential medium; FBS, fetal bovine serum; PBS, phosphate-buffered saline; iNOS, inducible nitric oxide synthase; NO, nitric oxide; TNF- α , tumor necrosis factor- α ; NF- κ B, nuclear factor-kappa B; RT-PCR, reverse transcription-polymerase chain reaction; ELISA, enzyme-linked immunosorbent assay; EMSA, electrophoretic mobility shift assay.

INTRODUCTION

Many marine resources, especially marine algal polysaccharides such as alginate, fucoidan, carrageenan, and agarose have recently drawn a great attention from diverse research fields to develop new drugs and health foods or supplements. A number of studies have demonstrated that various polysaccharides from marine algae have various bioactivities including antitumor and immuno-modulatory activities. Among these polysaccharides, there have been extensive studies on the biological activities of fucoidan, a fucose-containing sulfated polysaccharide with particular focus on its immuno-modulatory actions. For instance, it has been reported that fucoidan induces NO production in RAW264.7 cells through p38 mitogen-activated protein kinase (MAP kinase) and NF- κ B-dependent signal transduction *via* macrophage scavenger receptors (1). However, another study found that blocking of macrophage scavenger receptors did not result in the inhibition of the fucoidan activity, suggesting that such receptor is not involved in fucoidan action (2). Recent studies have reported that fucoidan inhibits the release of NO from LPS-stimulated RAW264.7 cells (3, 4). Thus, the precise action mechanism of fucoidan in terms of NO production in stimulated macrophages remains a matter of debate.

Porphyra species are the important edible red algae abundantly cultivated in East and South-east Asia including Japan, and are commonly known as “nori” and traditionally used to prepare sushi. Porphyran, one of the main constituents of *Porphyra yezoensis*, related to agarose, is a linear sulfated polysaccharide comprising the hot-water

soluble portion of the cell wall and intracellular matrix (5-7). It is constituted of D-galactose, 3, 6-anhydro-L-galactose, 6-*O*-methyl-D-galactose, and L-galactose-6-sulfate. In addition to the health benefit as a major dietary fiber in “nori”, previous studies have reported that porphyran also has diverse physiological activities including antitumor, immuno-modulating, antioxidant, antihyperlipidemic, and hypercholesterolemic activities (8-12).

Although several studies on the bioactivities of porphyran have been conducted, it has not been studied yet whether porphyran affects the formation of NO in LPS-stimulated macrophages. Thus, in this study, we examined the effect of porphyran on NO production in LPS-stimulated RAW264.7 cells. Unexpectedly, we found that porphyran inhibited NO production and the expression of inducible nitric oxide synthase (iNOS) by blocking NF- κ B activation in LPS-stimulated RAW264.7 cells.

MATERIALS AND METHODS

Materials – LPS from *Escherichia coli* 0111: B4 (purified by phenol extraction), fucoidan (from *Fucus vesiculosus*), and custom oligonucleotides (iNOS and β -actin primers) were purchased from Sigma-Aldrich, Co. (St. Louis, MO, USA). Alamar blue cell counting reagent was purchased from Invitrogen, Ltd. (Paisley PA4 9RF, UK). Sulfanilic acid, *N*-1-naphthyl-ethylenediamine dihydrochloride, glacial acetic acid and 5-methylphenazium-methyl sulfate (PMS) were obtained from Wako Pure Chemical industries, Ltd. (Osaka, Japan). NF- κ B SN50 (cell-permeable inhibitor peptide) and

NF- κ B SN50M (cell-permeable inactive control peptide) were purchased from Calbiochem (La Jolla, CA, USA). PrimeScript[®] 1st strand cDNA synthesis kit for reverse transcription-polymerase chain reaction (RT-PCR) was purchased from TaKaRa Bio Inc. (Otsu, Shiga, Japan). GoTag Green Master Mix was purchased from Promega KK. (Tokyo, Japan). Rabbit anti-mouse iNOS antibody and goat anti-rabbit IgG-horseradish peroxidase conjugated secondary antibody were purchased from Upstate Biotechnology (Lake Placid, NY, USA). Rabbit anti-mouse β -actin antibody was obtained from Abcam Inc. (Cambridge, USA). Horseradish peroxidase-conjugated donkey anti-rabbit IgG and enhanced chemiluminescence western blotting detection reagents were obtained from Amersham Biosciences UK Ltd. (Buckinghamshire HP7 9NA, UK). BCA protein assay kit was purchased from Bio-Rad Lab. (Hercules, CA, USA). Anti-I κ B- α , anti-c-Rel (p65), and anti-phospho-I κ B- α antibodies were obtained from Cell Signaling Technology, Inc. (Beverly, MA, USA). Sepasol-RNA I Super, protease inhibitor cocktail, nitro blue tetrazolium (NBT), and nicotinamide adenine dinucleotide (NADH) were obtained from Nacalai tesque Co. (Kyoto, Japan). TNF- α capture antibody and anti-mouse TNF- α monoclonal antibody were purchased from Endogen, Inc. (Boston, MA, USA). NF- κ B consensus IRDye[®] 700 infrared dye labeled oligonucleotides probe was purchased from LI-COR Bioscience (Lincoln, NE USA). Other chemicals were of the highest grade commercially available.

Preparation of porphyrin – Porphyrin was prepared from nori (*Porphyra yezoensis*) as reported previously (13). In brief, dry sheets (3 g) of nori were homogenized with 500 ml

of 85% ethanol and heated at 75 °C with constant stirring for 1 h, and then filtered to remove 85% ethanol-soluble substances. This extraction was repeated again. The residue was washed with methanol and dried to obtain 2 g of partially decolorized powder. The decolorized powder was extracted with 1,000 ml of distilled water at 95 °C with constant stirring for 1.5 h and centrifuged at 8000 x g for 20 min, and then the supernatant was concentrated to 600 ml by using a rotary evaporator, and then digested with 50 units of nuclease P1 at 37°C for 24 h. After that, the mixture was heated for 5 min in the boiling water bath, and centrifuged to remove the precipitates. To the supernatant, sodium acetate and acetic acid were added to make a 0.5 M sodium acetate solution, pH 5.0, and then ethanol was added to the concentration of 42% (v/v). After 1 h, the precipitates in the solution were removed by centrifugation, and ethanol was added to the supernatant to the concentration of 60%. The precipitate was dissolved in distilled water, dialyzed against distilled water, and lyophilized to obtain 0.4 g of porphyran. Before use, porphyran solution was filtered through an endotoxin-removing filter (Zetapor Dispo filter) purchased from Wako Pure Chemicals Industries, Ltd. (Osaka, Japan).

Cell culture – RAW264.7 (mouse macrophage) cells were obtained from the American Type Culture Collection (Rockville, MD, USA), and cultured in CO₂ (5%)-incubator at 37°C in Dulbecco's modified Eagle's minimum essential medium (DMEM) supplemented with 10% fetal bovine serum (FBS), penicillin (100 IU/ml), and streptomycin (100 µg/ml), which was used as the growth medium throughout the experiments otherwise specified.

Cytotoxicity assay – Cytotoxicities of polysaccharide samples were measured by the Alamar blue assay as described previously (14). In brief, adherent RAW264.7 cells in 96-well plates (3×10^4 cells/well) were treated with varying concentrations of porphyran or fucoidan for 24 h in the growth medium, and then Alamar blue reagent was added to the cells at a final concentration of 10%. After 2 h incubation at 37°C, the absorbance of each well was measured at 570 nm and 600 nm using a multiwell scanning spectrophotometer (Thermo Electron Co., Yokohama, Japan).

Measurement of DNA fragmentation – Adherent RAW264.7 cells in 35 mm-diameter culture dishes (2×10^6 cells/dish) were treated with 500 µg/ml of porphyran or fucoidan in the growth medium at 37°C for 24 h. After removal of the medium, the cells were lysed in 300 µl of ice-cold lysis buffer containing 0.5% Triton X-100, 10 mM Tris-HCl, pH 8.0, and 20 mM EDTA. The cell lysates were centrifuged for 20 min at 15,000 x g to separate DNA fragments (supernatant) from intact DNA (pellet). The DNA contents in supernatant and pellet fractions were measured using the diphenylamine reagent, and the extent of DNA fragmentation was estimated as described previously (15).

Nitrite assay for the estimation of nitric oxide (NO) – Nitrite, a stable reaction product of NO with O₂, in the supernatants from each treated RAW264.7 cells, was determined by the method based on Griess assay as described previously (16). In brief, adherent RAW264.7 cells in 96-well plates (3×10^4 cells/well) were treated with varying

concentrations of porphyrin (0 to 500 µg/ml) for 1 h in the growth medium, and then LPS was added at the final concentration of 2 ng/ml. After 18 h incubation at 37°C, Griess reagent (100 µl: 3 mM sulfanilic acid and 30 µM *N*-1-naphthyl-ethylenediamine dihydrochloride, and 25% glacial acetic acid) was added to 50 µl of cultured supernatant. After 20 min incubation at room temperature, the optical density was measured at 540 nm using a multiwell scanning spectrophotometer (Thermo electron Co., Yokohama, Japan). Calibration curve was made with known concentration of NaNO₂ standard solution.

Isolation of RNA and reverse transcription-polymerase chain reaction (RT-PCR) for the detection of iNOS mRNA – Adherent RAW264.7 cells in 24-well plates (5 x 10⁵ cells/well) were pre-treated with porphyrin at the concentrations of 0, 250, and 500 µg/ml for 1 h in the growth medium, and then incubated with or without LPS at the final concentration of 2 ng/ml. After 4 h incubation, total RNA of each treated cells was isolated using Sepasol-RNA I Super. Total RNA (2.5 µg) was reverse transcribed with an oligo dT primer in a 10 µl using PrimeScript[®] 1st strand cDNA synthesis kit according to the manufacturer's instruction. PCR was performed with 1 cycle of 70 s at 95°C, 25 cycles of 55 s at 93°C, 45 s at 61°C, 40 s at 72°C and 1 cycle of 100 s at 72°C, in a 25 µl reaction mixture containing 12.5 µl of GoTag Green Master Mix, 0.5 µl of forward and reverse iNOS primers (1µM each) or β-actin primers (10 pM each), 0.5 µl of 1st strand cDNA and 11 µl nuclease-free water. The primer sequences were 5'-CAACCAGTATTATGGCTCCT-3' (forward) and 5'-GTGACAGCCCGGTCTTTCCA-3' (reverse) for mouse iNOS (17) and, 5'

-GGAGAAGATCTGGCACCACACC-3' (forward) and 5' -CCTGCTTGCTGATCCACATCTGCTGG-3' (reverse) for mouse β -actin (18). The β -actin primer was used as an internal control. Each PCR reaction (10 μ l) product was run on 2% agarose gels containing 0.1 μ g/ml ethidium bromide, and the amplified products (835 bp for iNOS and 840 bp for β -actin) were detected by a Light capture (ATTO Co., Tokyo, Japan).

Western blot analysis – Analysis of iNOS was performed on whole-cell extracts. Adherent RAW264.7 cells in 35 mm-diameter culture dishes (2×10^6 cells/dish) were pre-treated with 0, 250, or 500 μ g/ml porphyran for 1 h in the growth medium. After 5 h incubation with LPS (final 2 ng/ml), the cells were washed three times with ice-cold PBS and lysed with 100 μ l extraction buffer (10 mM HEPES, 150 mM NaCl, 1 mM EGTA, 1% CHAPS, and 1% Triton X-100) containing 1% (v/v) protease inhibitor cocktail. After shaking for 30 min at 4°C, the extracts were obtained from the supernatant after centrifugation at 15,000 $\times g$ for 10 min, and the protein concentrations were measured with the BCA protein assay (Sigma) using bovine serum albumin as a standard. The extract was mixed with equal volume of 2 \times SDS-sample buffer and incubated at 100°C for 5 min. Samples containing 20 μ g of protein were applied on 8% SDS-PAGE, and then electrically transferred to a polyvinylidene difluoride (PVDF) membrane. The membrane was blocked with 1% skim milk in TBS-0.1% Tween 20 (TBST). Immunostaining of the blot was performed with the specific antibody against mouse iNOS and β -actin. Horseradish peroxidase conjugated-goat anti-rabbit IgG was used as a secondary

antibody. The blots were detected using an enhanced chemiluminescence kit (Amersham, Arlington Heights, IL). The analysis of I κ B- α and phospho-I κ B- α in cytosolic extracts and NF- κ B p65 in nuclear extracts were performed, respectively. Adherent RAW264.7 cells (2×10^6 cells/dish) were incubated in serum-free DMEM for 2 h at 37°C, and then porphyrin (0, 250, or 500 μ g/ml) was added to the cells and incubated for 1 h. After 30 min incubation with LPS (final 2 ng/ml), the cells were washed three times with ice-cold PBS and incubated with 100 μ l ice-cold cytosol extraction buffer (10 mM HEPES, pH 7.9, 1.5 mM MgCl₂, 10 mM KCl, 0.2% Igepal CA-630, 1 mM dithiothreitol, 20 mM β -glycerophosphate, 1mM sodium orthoradate, 0.5 mM phenylmethsulfonyl fluoride, 1 μ g/ml leupeptin, and 1 μ g/ml aprotinin) for 25 min on ice. The cytosolic extracts were collected after centrifugation at 7000 x g for 5 min at 4°C. The nuclear pellets were re-suspended in 30 μ l ice-cold nuclear extraction buffer (20 mM HEPES, pH 7.9, 1.5 mM MgCl₂, 0.45 M NaCl, 25% glycerol, 0.2 mM EDTA, 1 mM dithiothreitol, 0.5 mM phenylmethsulfonyl fluoride, 1 μ g/ml leupeptin and 1 μ g/ml aprotinin) and incubated on ice for 25 min, and the nuclear extracts were obtained after centrifugation at 15000 x g for 10 min at 4°C. After measuring the protein concentrations, the extract was mixed with equal volume of 2 x SDS-sample buffer and incubated at 100°C for 5 min. Samples containing 20 μ g of protein were applied on 12.5% SDS-PAGE, and then electrically transferred to a polyvinylidene difluoride (PVDF) membrane. The western blot analyses of the cytosolic and nuclear extracts were conducted by the similar way as described above except using specific anti-NF- κ B p65, anti-I κ B- α , and anti-phospho-I κ B- α antibody.

Enzyme-linked immunosorbent assay (ELISA) – Adherent RAW264.7 cells in 24-well plates (5×10^5 cells/well) were treated with porphyrin at the concentrations of 0, 250, and 500 $\mu\text{g/ml}$ for 1 h at 37°C in the growth medium. After further 4 h incubation with LPS (final 2 ng/ml), the levels of TNF- α in culture supernatants of treated cells were measured by sandwich ELISA with two antibodies to two different epitopes on TNF- α molecule by similar method as described previously (19). The TNF- α concentrations were estimated from a reference to a standard curve for serial twofold dilution of murine recombinant TNF- α .

Electrophoretic mobility shift assay (EMSA) – Adherent RAW264.7 cells in 35 mm-diameter culture dishes (2×10^6 cells/dish) were incubated in serum-free DMEM for 2 h at 37°C , and then porphyrin (0, 250, or 500 $\mu\text{g/ml}$) was added to the cells and incubated for 1 h. After further 30 min incubation with LPS (final 2 ng/ml), the nuclear proteins were extracted from the cells as described (20). EMSA were carried out with Odyssey[®] IRDye[®] 700 infrared dye labeled double-stranded oligonucleotides coupled with the EMSA buffer kit (LI-COR Bioscience, Lincoln, NE USA) according to manufacturer's instructions. Briefly, 5 μg of nuclear extract was incubated with 1 μl of IRDye[®] 700 Infrared dye labeled double-stranded oligonucleotides, 2 μl of 10 x binding buffer, 2.5 mM DTT, 0.25% Tween-20, and 1 μg of poly (dI-dC) in a total volume of 20 μl for 20 min at room temperature in the dark. Sample proteins were separated on a 4% polyacrylamide gel in 0.25 x Tris-borate-EDTA running buffer for 60 min at 100 V. The gel was scanned by direct infrared fluorescence detection on the Odyssey[®] imaging

system (LI-COR Bioscience, Lincoln, NE USA). NF- κ B IRDye[®] 700 infrared dye labeled oligonucleotide's sequences of the double-stranded DNA probes used were as follows: 5' -AGTTGAGGGGACTTTCCCAGGC-3' and 3' -TCAACTCCCCTGAAAGGGTCCG-5' (21, 22). The underlined nucleotides indicate the binding sites for NF- κ B. The specificity of the binding was examined using competition experiments, where 100-fold excess of the unlabeled oligonucleotides were added to the reaction mixture before adding the infrared dye labeled oligonucleotide.

Superoxide radical scavenging activity – The superoxide radical scavenging activity of porphyrin was assessed by the method described previously (23). For the assay, 0.5 μ l of 30 mM nitro blue tetrazolium (NBT), and 0.5 μ l of 46.8 mM NADH in 16 mM Tris-HCl buffer (pH 8.0) were added to 100 μ l of sample solution in 16 mM Tris-HCl buffer (pH 8.0) (final 10-1000 μ g/ml). The reaction was initiated by adding 0.5 μ l of 6 mM 5-methylphenazinium-methyl sulfate (PMS) to the mixture. The reaction mixture was incubated at room temperature for 5 min and the absorbance was measured at 560 nm by a spectrophotometer. Decrease in absorbance of the reaction mixture reflected increased superoxide radical-scavenging activity. The superoxide radical-scavenging activity was calculated using following equation.

$$\text{O}_2^- \text{ scavenging activity (\%)} = (1 - A_{\text{sample}}/A_{\text{control}}) \times 100$$

Where A_{sample} is the absorbance in the presence of sample, and A_{control} is the absorbance of Tris-HCl buffer alone without sample.

Statistical analysis – All the experiments were repeated at least three times. Data were expressed as means \pm SD and analyzed using one-way analysis of variance (ANOVA) followed by a Dunnett's test to determine any significant differences. A value of $p < 0.05$ was considered statistically significant.

RESULTS

Cytotoxicities of porphyrin and fucoidan on RAW264.7 cells – Since it has been reported that porphyrin inhibits cell proliferation and induces apoptosis in AGS human gastric cancer cells (9), we first examined the cytotoxicity of porphyrin on RAW264.7 cells. After 24 h incubation with varying concentrations of porphyrin, the viabilities of the cells were determined by the Alamar blue assay. As shown in Fig.1A, porphyrin showed no cytotoxic effect on RAW264.7 cells up to 500 $\mu\text{g/ml}$. In contrast to the lack of significant cytotoxic effect of porphyrin to RAW264.7 cells, fucoidan, a sulfated polysaccharide, showed cytotoxicity to RAW264.7 cells in a concentration dependent manner at the same concentration range tested. DNA fragmentation (Fig. 1B) and morphological changes (Fig. 1C) were also induced in the cells treated with 500 $\mu\text{g/ml}$ fucoidan, suggesting that the cytotoxicity of fucoidan was accompanied by induction of apoptosis, while no such significant apoptosis-related changes were observed in porphyrin-treated RAW264.7 cells (Fig. 1). Therefore, it seems likely that porphyrin and fucoidan are quite different in terms of cytotoxicity to RAW264.7 cells. Hence, it is considered that cellular toxicity of porphyrin on RAW264.7 may not be a main reason for any observed changes in NO

production or related events in the following studies.

Effect of porphyrin on NO production in LPS-treated RAW264.7 cells – Because of the extremely short half-life of NO, we measured nitrite as a stable biomarker of NO production in LPS stimulated RAW264.7 cells by Griess assay. Our preliminary study demonstrated that LPS potently induced NO production in RAW264.7 cells, even at a very low concentration (1 ng/ml). Time course analysis showed that NO production was initiated at 6 h and reached a stable and maximum level at 18 h, and the level continued to 24 h after stimulation with LPS (data not shown). Thus, in subsequent experiments, we evaluated nitrite concentration at 18 h. As shown in Fig. 2, nitrite level was almost undetectable in RAW264.7 cells treated with 500 µg/ml porphyrin alone, and porphyrin did not induce significant levels of NO at any other concentrations (1-1000 µg/ml) tested in RAW264.7 cells (data not shown).

To examine whether porphyrin affects LPS-induced NO production in RAW264.7 cells, cells were pre-incubated with porphyrin (0-500 µg/ml) for 1 h in the growth medium, followed by stimulation with LPS (2 ng/ml). After 18 h incubation, the nitrite levels of the supernatants of the treated cells were measured by Griess assay. As shown in Fig. 2, porphyrin inhibited NO production in LPS-stimulated RAW264.7 cells in a concentration-dependent manner, with 500 µg/ml porphyrin completely blocking the LPS-inducible NO production. Considering that 500 µg/ml porphyrin alone did not affect NO production and was not cytotoxic to RAW264.7 cells (Fig. 1), it seems obvious that inhibition of LPS-induced NO production in RAW264.7 cells by porphyrin is unrelated

to its any cytotoxic effects.

Effect of porphyrin on expression of iNOS in LPS-treated RAW264.7 cells – To study whether the inhibition of NO production by porphyrin is due to the inhibition of expression of inducible NO synthase (iNOS), which catalyses the generation of NO from L-arginine, RT-PCR and immuno blot analysis were performed to detect the levels of iNOS mRNA and protein expression, respectively. As shown in Fig. 3, treatment of cells with 2 ng/ml LPS resulted in increased expression of both iNOS mRNA and protein, while iNOS expression was not detected in RAW264.7 cells treated with control normal medium or 500 µg/ml porphyrin alone. Pretreatment of cells with porphyrin, especially at 500 µg/ml, inhibited iNOS expression by nearly 100% at the transcription and translation levels in LPS-stimulated RAW264.7 cells.

Effect of porphyrin on TNF- α secretion in LPS-treated RAW264.7 cells – LPS activates various intracellular signaling cascades leading to the secretion of various cytokines in addition to the induction of NO production *via* iNOS expression. In fact, increased level of TNF- α was detected in LPS-treated RAW264.7 cells (Fig. 4). We next investigated whether porphyrin can also affect the secretion of TNF- α from LPS-stimulated RAW264.7 cells. As shown in Fig. 4, porphyrin showed only a marginal inhibitory effect on the TNF- α production in LPS-treated RAW264.7 cells even at 500 µg/ml, at which NO production was completely inhibited, though.

Effect of porphyrin on LPS-induced NF- κ B activation in RAW264.7 cells – NF- κ B is an essential transcription factor for the induction of several inflammatory mediators including iNOS (24, 25). Therefore, an EMSA on the nuclear extracts from LPS-stimulated RAW264.7 cells was performed using IRDye[®] 700 Infrared dye labeled oligonucleotide with NF- κ B consensus sequence to examine whether porphyrin-caused inhibition of iNOS induction is due to the suppression of NF- κ B activation. As shown in Fig. 5A, increase in band thickness of the slow migrating nucleoprotein complex was observed after the LPS treatment for 30 min, suggesting that NF- κ B was activated by LPS. Pretreatment with porphyrin (250 or 500 μ g/ml) resulted in the inhibition of the LPS-inducible NF- κ B DNA binding in a concentration-dependent manner. The addition of 100-fold excess of the unlabeled oligonucleotides with same sequences to the nuclear extract resulted in almost complete abolishment of the binding activity, which confirms that the binding is specific to NF- κ B (Fig. 5B). To examine whether porphyrin directly inhibits NF- κ B binding to DNA, the nuclear extracts prepared from the LPS-treated RAW264.7 cells were treated with porphyrin and labeled NF- κ B oligonucleotides. The *in vitro* exposure of the LPS-activated nuclear extracts to porphyrin caused no decrease in the NF- κ B DNA binding (Fig. 5C). Hence, it seems likely that porphyrin itself does not inhibit NF- κ B DNA binding activity.

The process of the activation of NF- κ B proceeds through phosphorylation and degradation of the I κ B- α inhibitory subunit and subsequent translocation of p65/p50 complex into the nucleus (26). Thus, the effect of porphyrin on nuclear translocation of p65 was also examined by immunoblotting. As shown in Fig. 6A, the nuclear levels of the

p65 protein increased in LPS-treated RAW264.7 cells, and the LPS-induced nuclear translocation of p65 was blocked by porphyran in a concentration-dependent manner. In addition, immuno blot analysis using specific antibodies indicated that both the phosphorylation and degradation of I κ B- α were also inhibited by a pretreatment with porphyran (Fig. 6B). These results suggest that porphyran blocks the nuclear translocation of NF- κ B through the inhibition of the phosphorylation and subsequent degradation of I κ B- α . Specific NF- κ B inhibitor peptide (SN50) blocked LPS-induced NO production in a concentration-dependent manner, while inactive control peptide SN50M had no effect (Fig. 7). These results indicate that NF- κ B activation is required for LPS-induced NO production, and may support the idea that inhibition of NF- κ B activation by porphyran is its main action mechanism to inhibit NO production in LPS-stimulated RAW264.7 cells.

Superoxide radical scavenging activity of porphyran – It has been reported that sulfated polysaccharides from marine algae including fucoidan and porphyran have antioxidant activities (27-32), and naturally occurring antioxidant compounds such as β -carotene have suppressive effect on NF- κ B activation in macrophages stimulated with LPS (33). Thus, there is a possibility that antioxidant property of porphyran might be partly responsible for the inhibitory effects on the inflammatory response of LPS-stimulated RAW264.7 cells. To study this point, the antioxidant activity of porphyran was assessed by the superoxide radical scavenging activity. As shown in Fig. 8, porphyran significantly scavenged superoxide radical in a concentration-dependent manner.

Porphyran at final concentration of 500 µg/ml scavenged superoxide anion radical by more than 50%.

DISCUSSION

Porphyran is a linear sulfated polysaccharide composed of galactose and 3, 6-anhydrogalactose, which are partially substituted with galactose-6-sulfate and 6-*O*-methyl-galactose. In the present study, we found that porphyran inhibited the production of NO through the inhibition of iNOS expression in LPS-stimulated RAW264.7 cells. RT-PCR analysis revealed that iNOS mRNA level was significantly reduced by porphyran in LPS-treated RAW264.7 cells. Since β -actin mRNA levels remained unchanged, the possibility of the nonspecific actions of porphyran on gene expression may be excluded. Thus, the inhibition of NO production is due to the inhibition of iNOS mRNA transcription. These results differ from those obtained using murine peritoneal macrophages, which demonstrated that porphyran fraction prepared from *P. yezoensis* induced NO and TNF- α production (34, 35). Although the exact reason for these differences is uncertain now, it may be due to the different cell types used in the studies, as well as differences in sources and preparation procedures of porphyrans used. In fact, it has been known that porphyran has a quite complicated and heterogeneous structure depending on the source of the porphyran, and the even cultivation manner or harvest time also influences the structure of porphyran derived from same algal species (36, 37). Although further studies are required for the clarification of the mechanism of

porphyrin to inhibit NO production in LPS-stimulated macrophages as well as for its structure-activity relationship, it seems that our results can exclude the possibility that porphyrin directly interferes with interaction between NF- κ B protein and DNA, since no significant inhibitory effect of porphyrin on NF- κ B DNA binding was observed in *in vitro* EMSA (Fig. 5C).

NO is a gaseous free radical involved in various physiological processes such as vasodilation, smooth muscle regulation, neurotransmission, and apoptosis (38-41). NO also plays a role in anti-infectious immune responses as an important modulator in both innate and adaptive immunity (42). On the other hand, the excess amount of NO produced by iNOS in activated macrophages contributes to numerous severe inflammatory diseases including sepsis and arthritis (43, 44). Since NO particularly induces the impaired vascular reactivity and causes pathological changes (45), the selective inhibition of iNOS expression in inflammatory cells (*e. g.*, macrophages) may provide an important therapeutic strategy for inflammatory diseases.

Similar to our present results on porphyrin, it has recently been reported that fucoidan, a sulfated polysaccharide derived from brown algae, inhibits NO production and the expression of iNOS in LPS-activated macrophages (3). The inhibitory effect of fucoidan on NO production and iNOS expression in other cell types are also reported (4, 46). Fucoidan shows various biological activities. Particularly, its anti-inflammatory and anti-complement actions have drawn significant attention (47, 48). The suppressive effects on iNOS expression in activated macrophages might partly explain its anti-inflammatory actions. In contrast, it was reported that fucoidan induces NO

production from macrophages *via* a p38 MAP kinase and NF- κ B-dependent mechanism (1). Since fucoidan has highly complicated structure similar to porphyran, it can vary depending on the algal sources. Comparative study on the anti-inflammatory, anti-angiogenic, anticoagulant, and anti-adhesive activities of fucoidans prepared from nine species of brown algae showed that the source and composition of fucoidan likely affect the biological activity to a certain extent (49). Further studies are necessary to clarify the structure-activity relationship of sulfated polysaccharides fucoidan and porphyran especially their effects on NO production in activated macrophages.

LPS can induce the secretion of inflammatory cytokines such as TNF- α and IL-6 in addition to iNOS expression in macrophages. In fact, increased level of TNF- α was detected in the medium of LPS-treated RAW264.7 cells. It is of interest to point out that porphyran had only a marginal inhibitory effect on the level of TNF- α . This observation suggests that porphyran may not interrupt the extracellular interaction between LPS and the specific receptors but affect the intracellular signaling pathways such as the activation process of transcription factor. The promoter region of iNOS gene has several homologous consensus sequences for binding of the transcription factors such as NF- κ B, c-jun/c-Fos heterodimers known as AP-1, and CCAAT box enhancer binding protein (C/EBP) (30, 50). In this study, EMSA using oligonucleotides probe corresponding to the consensus binding site for NF- κ B was carried out, and we found that LPS-induced NF- κ B activation was inhibited by porphyran. It is well known that NF- κ B exists in the cytosol as an inactive trimeric complex in which the p50/p65 protein dimer is associated with I κ B known as an inhibitory subunit (51). When the cells are stimulated

with extracellular stimulus, phosphorylation and subsequent degradation of I κ B occurs. Such changes lead to dissociation of I κ B from the complex and to produce activated NF- κ B, which translocates into nucleus and activates the target gene expression. The immunoblot analysis using the specific antibodies demonstrated that the LPS-induced phosphorylation and degradation of I κ B- α , which is prerequisite for p65 activation, was suppressed in the porphyrin-pretreated cells. In addition, the translocation of p65 into nuclear in response to LPS was also inhibited by porphyrin. Thus, these results suggest that the inhibition of NF- κ B activation through the inhibition of the phosphorylation and degradation of I κ B- α may be an action mechanism of porphyrin to inhibit iNOS expression in LPS-stimulated RAW264.7 cells.

The activation status of various transcription factors are influenced by the intracellular redox condition and controlled by the generation of ROS (30). Recently, it has been suggested that porphyrin and fucoidan acts as a potential antioxidant or radical-scavenger (30-32). In fact, our porphyrin sample showed superoxide radical scavenging activity. Regarding naturally occurring antioxidants, which have inhibitory activities on inflammatory responses of macrophages, it has recently been reported that lycopene, a major carotenoid in tomatoes, inhibited LPS-induced production of NO through the suppression of NF- κ B activation in RAW264.7 cells, but had no effect on TNF- α (52). Similarity between porphyrin and lycopene in terms of the effect on LPS-stimulated RAW264.7 cells suggests that the antioxidant property may be one of the important factors responsible for the anti-inflammatory activities towards LPS-stimulated macrophages. In addition to lycopene, diosgenin, a steroid saponin found in several plants,

which has antioxidant activity, has also been reported to inhibit iNOS expression in LPS/interferon γ -activated murine macrophages without affecting the secretion of TNF- α (53). Since diosgenin inhibited LPS/interferon γ -induced NF- κ B and AP-1 activation, the inhibition of iNOS expression by diosgenin was considered at the transcriptional level. As another example, prodigiosin isolated from marine bacteria *Hahella chejuensis* has also been reported to suppress LPS-induced NO production by inhibiting p38 MAPK, JNK and NF- κ B activation in mouse peritoneal macrophages, but had no effect on the production of cytokines including TNF- α (54). Although porphyran, lycopene, diosgenin, and prodigiosin are chemically unrelated compounds, they may have a similar action mechanism on activated macrophages. Namely, these compounds inhibit NO production through the inhibition of NF- κ B activation without affecting the production of TNF- α , suggesting that NF- κ B separately regulates iNOS expression and secretion of TNF- α . Other several natural compounds with antioxidant or free radical scavenging activities have been reported to inhibit NO production by activated macrophages through the inhibition of the NF- κ B activation (3, 55-57). Thus, it seems likely that the specific inhibition of NO production *via* suppression of NF- κ B activation may be a common action mechanism of anti-inflammatory agents with antioxidant activity. Further studies are required to clarify this point as well as exact action mechanism of porphyran on the inhibition of NO production in activated macrophages.

In conclusion, our *in vitro* studies demonstrated that porphyran inhibited NO production in macrophages activated by LPS through the inhibition of NF- κ B activation. This inhibition may explain some of the anti-inflammatory effects of porphyran.

References

1. Nakamura, T., Suzuki, H., Wada, Y., Kodama, T., and Doi, T. (2006) Fucoidan induces nitric oxide production *via* p38 mitogen-activated protein kinase and NF- κ B-dependent signaling pathways through macrophage scavenger receptors. *Biochem. Biophys. Res. Commun.* **343**, 286-294
2. Oomizu, S., Yanase, Y., Suzuki, H., Kameyoshi, Y., and Hide, M. (2006) Fucoidan prevents C epsilon germline transcription and NF- κ B p52 translocation for IgE production in B cells. *Biochem. Biophys. Res. Commun.* **350**, 501-507
3. Yang, J.W., Yoon, S.Y., Oh, S.J., Kim, S.K., and Kang, K.W. (2006) Bifunctional effects of fucoidan on the expression of inducible nitric oxide synthase. *Biochem. Biophys. Res. Commun.* **346**, 345-350
4. Do, H., Pyo, S., and Sohn, E.H. (2010) Suppression of iNOS expression by fucoidan is mediated by regulation of p38 MAPK, JAK/STAT, AP-1 and IRF-1, and depends on up-regulation of scavenger receptor B1 expression in TNF- α - and IFN- γ -stimulated C6 glioma cells. *J. Nutritional Biochem.* **21**, 671-679
5. Morrice, L.M., McLean, M.W., Long, W.F., and Williamson, F.B. (1983) Porphyrin primary structure. *Eur. J. Biochem.* **133**, 673-684
6. Yoshimura, T., Tsuge, K., Sumi, T., Yoshiki, M., Tsuruta, Y., Abe, S., Nishino, S., Sanematsu, S., and Koganemaru, K. (2006) Isolation of porphyrin-degrading marine microorganisms from the surface of red alga, *Porphyra yezoensis*. *Biosci. Biotechnol. Biochem.* **70**, 1026-1028

7. Takahashi, K., Hirano, Y., Araki, S., and Hattori, M. (2000) Emulsifying ability of porphyran prepared from dried nori, *Porphyra yezoensis*, a red alga. *J. Agric. Food. Chem.* **48**, 2721-2725
8. Ishihara, K., Oyamada, C., Matsushima, R., Murata, M., and Muraoka, T. (2005) Inhibitory effect of porphyran, prepared from dried “Nori”, on contact hypersensitivity in mice. *Biosci. Biotechnol. Biochem.* **69**, 1824-1830
9. Kwon, M.J., and Nam, T.J. (2006) Porphyran induces apoptosis related signal pathway in AGS gastric cancer cell lines. *Life Sci.* **79**, 1956-1962
10. Ren, D., Noda, H., Amano, H., Nishino, T., and Nishizawa, K. (1994) Study on antihypertensive antihyperlipidemic effects of marine algae. *Fish Sci.* **60**, 33-40
11. Tsuge, K., Okabe, M., Yoshimura, T., Sumi, T., Tachibana, H., and Yamada, K. (2004) Dietary effect of porphyran from *Porphyra yezoensis* on growth and lipid metabolism of Sprague-Dawley rats. *Food Sci. Technol. Res.* **10**, 147-151
12. Inoue, N., Yamano, N., Sakata, K., Nagao, K., Hama, Y., Yanagita, T. (2009) The sulfated polysaccharide porphyran reduces apolipoprotein B100 secretion and lipid synthesis in HepG2 cells. *Biosci. Biotechnol. Biochem.* **73**, 447-449
13. Hama, Y., Nakagawa, H., Kurosawa, M., Sumi, T., Xia, X., and Yamaguchi, K. (1998) A gas chromatographic method for the sugar analysis of 3, 6-anhydrogalactose-containing algal galactans. *Anal Biochem.* **265**, 42-48
14. Jiang, Z., Okimura, T., Yokose, T., Yamasaki, Y., Yamaguchi, K., and Oda, T. (2010) Effects of sulfated fucan, ascophyllan, from the brown Alga *Ascophyllum nodosum* on various cell lines: A comparative study on ascophyllan and fucoidan. *J. Biosci.*

Bioeng. **110**, 113-117

15. Kuramoto, T., Uzuyama, H., Hatakeyama, T., Tamura, T., Nakashima, T., Yamaguchi, K., and Oda, T. Cytotoxicity of a GalNAc-specific C-type lectin CEL-I toward various cell lines (2005). *J. Biochem.* **137**, 41-50
16. Yamanishi, T., Hatakeyama, T., Yamaguchi, K., and Oda, T. (2009) CEL-I, an *N*-acetylgalactosamine (GalNAc)-specific C-type lectin, induces nitric oxide production in RAW264.7 mouse macrophage cell line. *J. Biochem.* **146**, 209-217
17. Chen, C.Y., Peng, W.H., Tsai, K.D., and Hsu, S.L. (2007) Luteolin suppresses inflammation-associated gene expression by blocking NF- κ B and AP-1 activation pathway in mouse alveolar macrophages. *Life Sci.* **81**, 1602-1614
18. Chang, S.H., Mun, S.H., Ko, N.Y., Lee, J.H., Jun, M.H., Seo, J.Y., Kim, Y.M., Choi, W.S., and Her, E. (2005) The synergistic effect of phytohemagglutinin and interferon- γ on the expression of tumor necrosis factor- α from RAW 264.7 cells. *Immunol Lett.* **98**, 137-143
19. Yamanishi, T., Yamamoto, Y., Hatakeyama, T., Yamaguchi, K., and Oda, T. (2007) CEL-I, an invertebrate *N*-acetylgalactosamine-specific C-type lectin, induces TNF- α and G-CSF production by mouse macrophage cell line RAW264.7 cells. *J. Biochem.* **142**, 587-595
20. Zingarelli, B., Hake, P.W., Yang, Z., O'Conor, M., Denenberg, A., and Wong, H.R. (2002) Absence of inducible nitric oxide synthase modulates early reperfusion-induced NF- κ B and AP-1 activation and enhances myocardial damage. *FASEB J.* **16**, 327-342

21. Kim, J.W., and Kim, C. (2005) Inhibition of LPS-induced NO production by taurine chloramine in macrophages is mediated through Ras-ERK-NF- κ B. *Biochemical Pharmacology* **70**, 1352-1360
22. Pokharel, Y.R., Liu, Q.H., Aryal, D.K., Kim, Y.G., Woo, E.R., and Kang, K.W. (2007) 7,7'-Dihydroxy bursehernin inhibits the expression of inducible nitric oxide synthase through NF- κ B DNA binding suppression. *Nitric Oxide: Biology and Chemistry* **16**, 274-285
23. Nishimiki, M., Rao, N. A., and Yagi, K. (1972) The occurrence of superoxide anion in the reaction of reduced phenazine methosulfate and molecular oxygen. *Biochem. Biophys. Res. Commun.* **46**, 849-864
24. Muller, J.M., Ziegler-Heitbrock, H.W., and Baeuerle, P.A. (1993) Nuclear factor kappa B, a mediator of lipopolysaccharide effects. *Immunobiology* **187**, 233-256.
25. Guha, M., and Mackman, N. (2001) LPS induction of gene expression in human monocytes. *Cell. Signal.* **13**, 85-94
26. Wang, T., Zhang, X., and Li, J.J. (2002) The role of NF- κ B in the regulation of cell stress responses. *Int. Immunopharmacol.* **2**, 1509-1520
27. Leung, M.Y.K., Liu, C., Koon, J.C.M., and Fung, K.P. (2006) Polysaccharide biological response modifiers. *Immunol. Lett.* **105**, 101-114
28. Zhang, Z., Zhang, Q., Wang, J., Zhang H., Niu, X., and Li, P. (2009) Preparation of the different derivatives of the low-molecular-weight porphyrin from *Porphyrin haitanensis* and their antioxidant activities *in vitro*. *Inter. J. Biol. Macromol.* **45**, 22-26

29. Zhang, Q., Li, N., Zhou, G., Lu, X., Xu, Z., and Li, Z. (2003) *In vivo* antioxidant activity of polysaccharide fraction from *Porphyra haitanesis* (Rhodophyta) in aging mice. *Pharmacol. Res.* **48**, 151-155
30. Lee, A.K., Sung, Y.C., Kim, Y.C., and Kim, S.G. (2003) Inhibition of lipopolysaccharide-inducible nitric oxide synthase, TNF- α and COX-2 expression by sauchinone effects on I- κ B α phosphorylation, C/EBP and AP-1 activation. *Br. J. Pharmacol.* **139**, 11-20
31. Zhang, Q., Yu, P., Li, Z., Zhang, H., Xu, Z., and Li, P. (2003) Antioxidant activities of sulfated polysaccharide fractions from *Porphyra haitanesis*. *J. Applied Phycol.* **15**, 305-310
32. Zhang, Z., Zhang, Q., Wang, J., Shi, X., Song, H., and Zhang, J. (2009) *In vitro* antioxidant activities of acetylated, phosphorylated and benzoylated derivatives of porphyran extracted from *Porphyra haitanensis*. *Carbohydrate Poly.* **78**, 449-453
33. Bai, S.K., Lee, S.J., Na, H.J., Ha, K.S., Han, J.A., Lee, H., Kwon, Y.G., Chung, C.K., and Kim, Y.M. (2005) Beta-carotene inhibits inflammatory gene expression in lipopolysaccharide-stimulated macrophages by suppressing redox-based NF-kappaB activation. *Exp. Mol. Med.* **37**, 323-334
34. Yoshizawa, Y., Enomoto, A., Todoh, H., Ametani, A., and Kaminogawa, S. (1993) Activation of murine macrophage by polysaccharide fractions from a marine alga (*Porphyra yezoensis*). *Biosci. Biotechnol. Biochem.* **57**, 1862-1866
35. Yoshizawa, Y., Ametani, A., Tsunehiro, J., Nomura, K., Itoh, M., Fukui, F., and Kaminogawa, S. (1995) Macrophage stimulation activity of the polysaccharide

- fraction from a marine alga (*Porphyra yezoensis*): Structure-function relationships and improved solubility. *Biosci. Biotechnol. Biochem.* **59**, 1933-1937
36. Rees, D.A. (1961) Estimation of the relative amounts of isomeric sulfate esters in some sulfated polysaccharides. *J. Chem. Soc.* 5168-5171
 37. Araki, S., Oohusa, T., Saitoh, M., and Sakurai, T. (1977) The quality of “Nori”, dried laver, in special reference to the contents of 3, 6-anhydro-galactose in porphyran. *Jpn. J. Phycol.* **25**, 19-23
 38. Huang, P.L., Huang, Z., Mashimo, H., Bloch, K.D., Moskowitz, M.A., Bevan, J.A., and Fishman, M.C. (1995) Hypertension in mice lacking the gene for endothelial nitric oxide synthase. *Nature* **377**, 239-242
 39. Nelson, R.J., Demas, G.E., Huang, P.L., Fishman, M.C., Dawson, V.L., Dawson, T.M., and Snyder, S.H. (1995) Behavioral abnormalities in male mice lacking neuronal nitric oxide synthase. *Nature* **378**, 383-386
 40. Dimmeler, S., and Zeiher, A.M. (1997) Nitric oxide and apoptosis: another paradigm for the double-edged role of nitric oxide. *Nitric Oxide: Biology and Chemistry* **1**, 275-281
 41. Lowenstein, C.J., Dinerman, J.L., and Snyder, S.H. (1994) Nitric oxide: a physiologic messenger. *Ann Intern Med.* **120**, 227-237
 42. Bogdan, C., Rölinghoff, M., and Diefenbach, A. (2000) Reactive oxygen and reactive nitrogen intermediates in innate and specific immunity. *Curr. Opin. Immunol.* **12**, 64-76
 43. Szabão, C. (1998) Role of nitric oxide in endotoxic shock. An overview of recent

- advances. *Ann. NY Acad. Sci.* **851**, 422-425
44. Weinberg, J.B. (2000) Nitric oxide synthase 2 and cyclooxygenase 2 interactions in inflammation. *Immunol. Res.* **22**, 319-341
 45. Symeonides, S., and Balk, R.A. (1999) Nitric oxide in the pathogenesis of sepsis. *Infect Dis. Clin. North Am.* **13**, 449-463
 46. Cui, Y., Zhang, L., Zhang, T., Luo, D., Jia, Y., Guo, Z., Zhang, Q., Wang, X., and Wang, X. (2010) Inhibitory effect of fucoidan on nitric oxide production in lipopolysaccharide-activated primary microglia. *Clinical. Exp. Pharm. Physiol.* **37**, 422-428
 47. Berteau, O., and Mulloy, B. (2003) Sulfated fucans, fresh perspectives: structures, functions, and biological properties of sulfated fucans and an overview of enzymes active toward this class of polysaccharide. *Glycobiology* **13**, 29R-40R
 48. Blondin C., Chaubet, F., Nardella, A., Siquin, C., and Jozefonvicz, J. (1996) Relationships between chemical characteristics and anticomplementary activity of fucans. *Biomaterials* **17**, 597-603
 49. Cumashi, A., Ushakova, N.A., Preobrazhenskaya, M.E., D’Incecco, A., Piccoli, A., Totani, L., Tinari, N., Morozovich, G.E., Berman A.E., Bilan M.I., Usov, A.I., Ustyuzhanina, N.E., Grachev, A.A., Sanderson, C.J., Kelly, M., Rabinovich, G.A., Iacobelli, S., Nifantiev, N.E., and on behalf of the Consorzio Interuniversitario Nazionale per la Bio-Oncologia (CINBO), Italy. (2007) An comparative study of the anti-inflammatory, anticoagulant, antiangiogenic, and antiadhesive activities of nine different fucoidans from brown seaweeds. *Glycobiology* **17**, 541-552

50. Cieslik, Y., Zhu, Y., and Wu, K.K. (2002) Salicylate suppresses macrophage nitric-oxide synthase-2 and cyclo-oxygenase-2 expression by inhibiting CCAAT/enhancer-binding protein- β binding *via* a common signaling pathway. *J. Biol. Chem.* **277**, 49304-49310
51. Romics, L., Kodys, K., Dolganiuc, A., Graham, L., Velayudham, A., Mandrekar P, and Szabo G. (2004) Diverse regulation of NF- κ B and peroxisome proliferator-activated receptors in murine nonalcoholic fatty liver. *Hepatology* 40, 376-385
52. Feng, D., Ling, W.-H., and Duan R.-D. (2010) Lycopene suppresses LPS-induced NO and IL-6 production by inhibiting the activation of ERK, p38 MAPK, and NF- κ B in macrophages. *Inflamm. Res.* **59**, 115-121
53. Jung, D. H., Park, H. J., Byun, H. E., Park, Y. M., Kim, T. W., Kim, B. O., Um, S. H., Pyo, S. (2010) Diosgenin inhibits macrophage-derived inflammatory mediators through downregulation of CK2, JNK, NF- κ B and AP-1 activation. *Int. Immunopharmacol.* **10**, 1047-1054
54. Huh, J. E., Yim, J. H., Lee, H. K., Moon, E. Y., Rhee, D. K., Pyo, S. (2007) Prodigiosin isolated from *Hahella chejuensis* suppresses lipopolysaccharide-induced NO production by inhibiting p38 MAPK, JNK and NF- κ B activation in murine peritoneal macrophages. *Int. Immunopharmacol.* **7**, 1825-1833
55. Pergola, C., Rossi, A., Dugo, P., Cuzzocrea, S., and Sautebin, L. (2006) Inhibition of nitric oxide biosynthesis by anthocyanin fraction of blackberry extract. *Nitric Oxide: Biology and Chemistry* **15**, 30-39

56. Han, Y.J., Kwon, Y.G., Chung, H.T., Lee, S.K., Simmons, R.L., Billiar, T.R., and Kim, Y.M. (2001) Antioxidant enzymes suppress nitric oxide production through the inhibition of NF- κ B activation: role of H₂O₂ and nitric oxide in inducible nitric oxide synthase expression in macrophages. *Nitric Oxide: Biology and Chemistry* **5**, 504-513
57. Tang, S., Shen, X.Y., Huang, H.Q., Xu, S.W., Yu, Y., Zhou, C.H., Chen, S.R., Le, K., Wang, Y.H., and Liu, P.Q. (2011) Cryptotanshinone suppressed inflammatory cytokines secretion in RAW264.7 macrophages through inhibition of the NF- κ B and MAPK signaling pathways. *Inflammation* **34**, 111-118

FIGURE LEGENDS

Fig. 1. Cytotoxic effects of porphyran and fucoidan on RAW264.7 cells. (A) Adherent cells (3×10^4 cells/well in 96-well plates) in the growth medium were treated with varying concentrations of porphyran (●) or fucoidan (○) at 37°C. After 24 h, the cell viabilities were estimated by Alamar blue assay as described under "MATERIALS AND METHODS". Data represent means \pm SD of triplicate measurements. Asterisks indicate significant differences between with and without fucoidan ($p < 0.05$). (B) Adherent RAW264.7 cells in 35 mm-diameter culture dishes (2×10^6 cells/dish) were treated with 500 μ g/ml of fucoidan or porphyran in the growth medium at 37°C. After 24 h, DNA fragmentations of the treated cells were examined by diphenylamine assay as described under "MATERIALS AND METHODS". Asterisk indicates significant difference between with and without fucoidan ($p < 0.05$). (C) Adherent RAW264.7 cells in 35 mm-diameter culture dishes equipped with quartz glass were treated with 500 μ g/ml of fucoidan or porphyran in the growth medium at 37°C. After 24 h, the cells were observed under phase contrast microscope (BIOREVO BZ9000, KEYENCE Co., Osaka, Japan). The bar indicates 20 μ m.

Fig. 2. Effects of porphyran on NO production in LPS-treated RAW264.7 cells. Adherent cells (3×10^4 cells/well in 96-well plates) were pre-incubated with various concentrations of porphyran (0-500 μ g/ml) in DMEM supplemented with 10% FBS at 37°C for 1 h, followed by the addition of LPS (final 2 ng/ml). After 18 h incubation, the

NO levels in the supernatants of the treated cells were estimated as described under "MATERIALS AND METHODS". Data represent means \pm SD of triplicate measurements. Asterisks indicate significant differences between with and without porphyrin ($p < 0.05$).

Fig. 3. Effects of porphyrin on LPS-induced iNOS mRNA and iNOS protein levels in RAW264.7 cells. (A) Adherent cells (5×10^5 cells/well in 24-well plates) were pre-incubated with various concentrations of porphyrin (0, 250, or 500 $\mu\text{g/ml}$) in DMEM supplemented with 10% FBS at 37°C for 1 h, followed by the addition of LPS (final 2 ng/ml). After 4 h incubation, the total RNA of each treated cells was subjected to RT-PCR analysis as described under "MATERIALS AND METHODS". (B) Adherent cells in 35 mm-diameter culture dishes (2×10^6 cells/dish) were pre-incubated with various concentrations of porphyrin (0, 250, or 500 $\mu\text{g/ml}$) in DMEM supplemented with 10% FBS at 37°C for 1 h, followed by the addition of LPS (final 2 ng/ml). After 5 h incubation, the whole-cell lysates were analyzed by western blot analysis as described under "MATERIALS AND METHODS"

Fig. 4. Effect of porphyrin on LPS-induced TNF- α secretion from RAW264.7 cells. Adherent cells (5×10^5 cells/well in 24-well plates) were pre-incubated with various concentrations of porphyrin (0, 250, or 500 $\mu\text{g/ml}$) in DMEM supplemented with 10% FBS at 37°C for 1 h, followed by the addition of LPS (final 2 ng/ml). After 4 h incubation, the TNF- α levels in the culture medium of the treated cells were measured by ELISA as

described under "MATERIALS AND METHODS". Data represent means \pm SD of triplicate measurements.

Fig. 5. Effects of porphyran on LPS-induced NF- κ B activation in RAW264.7 cells.

(A) *In vivo* effect of porphyran on DNA binding of NF- κ B. Adherent cells in 35 mm-diameter culture dish (2×10^6 cells/dish) were incubated in serum-free DMEM at 37°C for 2 h, and then the cells were incubated with various concentrations of porphyran (0, 250, or 500 μ g/ml) in serum-free DMEM at 37°C for 1 h, followed by the addition of LPS (final 2 ng/ml). After 30 min incubation, gel shift assay of NF- κ B was performed with nuclear extracts prepared from the treated cells as described under "MATERIALS AND METHODS". (B) EMSA was performed on the nuclear extract prepared from LPS-stimulated RAW264.7 cells in the presence of 100-fold excess unlabeled oligonucleotides with the same NF- κ B consensus sequence. (C) *In vitro* effect of porphyran on DNA binding of NF- κ B. Porphyran (0, 250, or 500 μ g/ml) was added to the assay mixture containing the nuclear extracts prepared from the LPS-stimulated RAW264.7 cells and infrared dye labeled oligonucleotides with NF- κ B consensus sequence, and incubated for 20 min, and then subjected to EMSA.

Fig. 6. Effects of porphyran on the nuclear translocation of NF- κ B p65 and phosphorylation and degradation of I κ B- α in LPS-treated RAW264.7 cells. (A)

Western blot analysis was conducted on NF- κ B p65 in the nuclear extracts prepared from RAW264.7 cells treated with porphyran and subsequent stimulation with LPS as

described under "MATERIALS AND METHODS". (B) Western blot analyzes were conducted on the I κ B- α and phosphorylated I κ B- α in the cytosolic extracts prepared from RAW264.7 cells treated with porphyrin and subsequent stimulation with LPS as described under "MATERIALS AND METHODS".

Fig. 7. Effects of specific NF- κ B inhibitor peptide (SN50) and its inactive control peptide (SN50M) on NO production in LPS-treated RAW264.7 cells. Adherent cells (3×10^4 cells/well in 96-well plates) were pre-incubated with various concentrations of SN50 (●) or SN50M (○) in DMEM supplemented with 10% FBS at 37°C for 2 h, followed by the addition of LPS (final 2 ng/ml). After 18 h incubation, the NO level in the supernatants of the treated cells were estimated as described under "MATERIALS AND METHODS". Data represent means \pm SD of triplicate measurements. Asterisks indicate significant differences between with and without SN50 ($p < 0.05$).

Fig. 8. Scavenging effects of porphyrin on superoxide anion radical. Scavenging activity of porphyrin toward superoxide anion radical was measured as described under "MATERIALS AND METHODS". Experiments were performed in triplicate, and values shown are the means \pm SD. Asterisks indicate significant differences between with and without porphyrin ($p < 0.05$).

Fig.1




Fig.2




Fig.3




Fig.4




Fig. 5




Fig. 6




Fig. 7




Fig. 8

