

Morphological Study of the Adrenal Cortex in Stroke-prone Spontaneously Hypertensive Rats (SHRSP)

—Special Reference to the Zona Glomerulosa by
Electron Microscopy and Morphometry—

Kazuo MAWATARI

*Department of Pathology, Nagasaki University School
of Medicine, Nagasaki, Japan*

Received for publication, March 25, 1982

The adrenal zona glomerulosa of stroke-prone spontaneously hypertensive rats (SHRSP) at 18 week old was examined morphologically and morphometrically. An increase in width of the zona glomerulosa was found including hypertrophy and hyperplasia of the cells. The area of cells in this zone of SHRSP was significantly greater than that of Wistar-Kyoto strain rats (WKY). The mitochondria of this zone were smaller in size, pleomorphic in shape and increased in number, and the percent area of the mitochondrial compartment of SHRSP was significantly lesser than that of WKY. The lipid droplets in zona glomerulosa cells of SHRSP were numerous and the percent area of lipid droplets compartment of SHRSP was significantly greater than that of WKY. Therefore, the zona glomerulosa cells were appeared to be hypertrophic. The smooth-surfaced endoplasmic reticulum and the Golgi apparatus were well developed. The relationship between these pathological changes and functional state in the zona glomerulosa of SHRSP was discussed.

INTRODUCTION

The spontaneously hypertensive rat (SHR)²⁶⁾ has been used as a model of human essential hypertension. Recently, OKAMOTO and YAMORI et al.²⁷⁾ established a sub-strain (stroke-prone SHR, SHRSP) from SHR; this strain is characterized by severe hypertension and a high incidence of cerebrovascular lesion.

The association of hypertension with endocrine disorders in human and in experimental animals has been described in previous studies. AOKI¹⁾ has reported on the relationship between endocrine organs and hypertension in SHR. And he pointed out²⁾

that the adrenal gland was necessary for the development and maintenance of high blood pressure in SHR. However, BAER et al.³⁾ failed to show that the adrenal was involved in pathogenesis of spontaneous hypertension.

Several authors have reported the studies on the plasma renin activity (PRA)¹⁵⁾¹⁸⁾²⁹⁾ and the plasma aldosterone concentration (PAC)¹⁹⁾²⁵⁾ in SHRSP. OGINO et al.²⁵⁾ revealed that PAC of 5 month old SHRSP was nearly equal to PAC of WKY.

Concerning the alterations of the adrenal cortex in SHR, some descriptions¹⁷⁾²²⁾²³⁾³¹⁾ were reported histologically, histochemically and morphometrically. On the other hand, there is no report on the adrenal cortex in SHRSP. This paper describes on morphologic and morphometric studies of the adrenal cortex in male SHRSP, with special emphasis on the zona glomerulosa, which is the major site of aldosterone production.

MATERIALS and METHODS

Nine male SHRSP and nine male WKY which were obtained from the colony of the Department of Pharmacology in Nagasaki University were used. All animals were given a standard food stock diet (FUNABASHI, Chiba) and maintained in a regular day and night rhythm. The blood pressure was recorded by tail plethysmographic method.

All rats were sacrificed by decapitation at 18 week old. Both adrenal glands were removed rapidly and trimmed of adherent fat and the right adrenal gland was weighed. The left adrenal gland was divided into two equal parts with sharp blade. A half was fixed in 10% aqueous formaldehyde solution, embedded in paraffin and cut in serial sections. These sections were stained with hematoxylin and eosin, Van Gieson's method for elastic fiber, Watanabe's method for reticulin fiber, PAS and Mallory-Azan's methods.

The other half was cut into smaller fragments and fixed in 0.1M phosphate-buffered 4% glutaraldehyde followed by 1.3% osmium tetroxide. After dehydration in acetone, the tissue was embedded in Epon 812. Semithin sections were cut and stained with toluidine blue to identify three zones of the adrenal cortex for ultrathin sections. Ultrathin sections were stained with uranyl acetate and lead citrate. Specimens were observed with a JEOL 100B electron microscopy.

As morphometric procedure, the average width of the zona glomerulosa and the area of cells, nucleus, mitochondria and lipid droplets in the zona glomerulosa were recorded. The percentage of the cytoplasm area (percent area) occupied by mitochondria or lipid droplets was also examined.

In order to study the average width of the zona glomerulosa 10 light micrographs of toluidine blue semithin sections from each animal were taken at a magnification of $\times 200$ and enlarged photographically to $\times 1,400$. Five to eight electron micrographs of the zona glomerulosa were taken from each specimen at a magnification of $\times 2,000$ or $\times 5,000$ and enlarged to $\times 8,000$ or $\times 20,000$ for the measurement of the area of cells,

nucleus, mitochondria and lipid droplets. On these randomly sampled micrographs each parameter was determined by a modular system for semiautomatic quantitative evaluation images of A. S. M., LEITZ, West Germany.

Comparison of group means for these parameters was performed by Student's t-test.

RESULTS

The systolic blood pressure of SHRSP was significantly higher than that of WKY. The body weight of SHRSP was lesser than that of WKY. The adrenal weight per 100 g body weight (relative adrenal weight) of SHRSP did not differ significantly from that of WKY. These data are shown in Table 1.

At autopsy, the incidence of cerebral lesions was 6/9 of SHRSP, and none of WKY had the cerebral lesion.

Histological findings. The zona glomerulosa was increased in width and consisted of six to seven cell layers in SHRSP(Fig. 1-a, b), whereas the zone was consisted of three to four cell layers in WKY(Fig. 2-a, b).

The zona glomerulosa cells in WKY were round, polygonal or elongated in shape and had clear cytoplasm with abundant lipid droplets (Fig. 2-b). The cells of this zona in SHRSP were the same shape and had enlarged clear cytoplasm containing numerous lipid droplets (Fig. 1-a, b). Therefore, the zona glomerulosa cells of SHRSP represented hypertrophic character.

In both groups of WKY and SHRSP, the "transitional zone" between zona glomerulosa and zona fasciculata was composed of two or more cell layers (Fig. 1-b, Fig. 2-b). The transitional zone cells were smaller in size than the zona glomerulosa cells, elongated in shape and contained few lipid droplets in the cytoplasm. But this zone in SHRSP (Fig. 1-b) was increased in width compared with WKY (Fig. 2-b).

Intracortical nodules constituted from zona fasciculata cells, which were large and round cells with lipid-rich cytoplasm, were found in SHRSP (Fig. 3) more frequently than in WKY.

Capsular arteries of SHRSP showed moderate medial thickening (Fig. 4) but few

Table 1 Comparison of Blood pressure, Body weight and Relative adrenal weight of SHRSP with WKY

	SHRSP	WKY
Blood Pressure (mmHg)	263 \pm 3.9	125 \pm 2.5
Body Weight (g)	286 \pm 15.4	331 \pm 9.2
Relative Adrenal Weight (mg/100gBW)	7.4 \pm 0.36	7.8 \pm 0.50

Each value represents the group mean \pm SE




Fig. 1-a. Outer adrenal cortex of SHRSP. G= zona glomerulosa, F=zona fasciculata. ($\times 200$).




Fig. 1-b. Zona glomerulosa(G)of SHRSP shows an increase in width with cellular hypertrophy and hyperplasia. Transitional zone (T) is prominent. ($\times 400$).




Fig. 2-a. Outer adrenal cortex of WKY. G= zona glomerulosa, F=zona fasciculata. ($\times 200$).




Fig. 2-b. Zona glomerulosa(G) and transitional zone (T) of WKY. ($\times 400$).



Fig. 3. Cortical nodule (arrow) of zona fasciculata in SHRSP. ($\times 40$).




Fig. 4. Medial thickening of cortical arteries in SHRSP. ($\times 200$).




Fig. 5. Fibrinoid necrosis of cortical artery in SHRSP. ($\times 40$).

intimal fibrous thickening or thromboembolism was appeared. Fibrinoid necrosis of the cortical artery was found in only one rat of SHRSP (Fig. 5).

Ultrastructural findings. In WKY, the zona glomerulosa cells were arranged in cords and here and there arranged in stratified or perivascular rosette-like structure (Fig. 6). The cells were round, polygonal or elongated in shape with round or ovoid shaped nucleus. The cells were separated by narrow space sometimes interrupted by desmosome-like junction structure (Fig. 7). The mitochondria were various in shape and in size. The mitochondrial matrix was slightly electron-dense and two mitochondrial membranes were clearly observed (Fig. 7). The mitochondria generally had short tubular cristae, situated near the periphery of these organelles. Lipid droplets with various size were found in the cytoplasm. Sometimes lipid droplets were surrounded by smooth-surfaced endoplasmic reticulum in tubular appearance but had no remarkable membranous material. The smooth-surfaced endoplasmic reticulum was seen in the cytoplasmic matrix of the zona glomerulosa cells among other organelles. This organelle seemed to be well developed and vesicular in shape (Fig. 7). A few profiles of rough-surfaced endoplasmic reticulum, free or clustered ribosomes and glycogen particles were scattered throughout the cytoplasm. The Golgi apparatus were generally well developed near the nuclei associated by amorphous dense-bodies and mitochondria and were formed by many lamellar rows of cisterna (Fig. 7).




Fig. 6. Zona glomerulosa cells of WKY. Note the arrangement in peri-vascular rosette-like structure. ($\times 6,700$).




Fig. 7. Zona glomerulosa cells of WKY. N=nucleus, M=mitochondria, L=lipid droplets, G=Golgi apparatus. Desmosome-like structure (arrow). ($\times 12,000$).

In SHRSP, the zona glomerulosa cells (Fig. 8-11) had no remarkable difference in the arrangement or structure and in shape from WKY. The nucleus was similar to that of WKY. The mitochondria of the zona glomerulosa cells in SHRSP showed more various in shape (Fig. 9, 10) than in WKY; round, polygonal, elongated, dumb-bell shaped, kidney shaped, etc. These organelles represented smaller in size but increased in number. The mitochondrial matrix was moderately electron-dense. The mitochondrial cristae was tubular as WKY but was prominent compared with WKY (Fig. 10). Some elongated mitochondria exhibited regular arrays of lamellar cristae which usually run parallel to long axis of the mitochondria (Fig. 10, 11-a). Amorphous electron-dense intramitochondrial droplets appeared more frequently in SHRSP (Fig. 11-b) than in WKY. The zona glomerulosa cells in SHRSP had a great deal of lipid droplets in the cytoplasm (Fig. 8-a). Among these droplets very large lipid droplets measuring up to $12\text{--}14\text{ }\mu\text{m}^2$ in area, which were never seen in WKY, appeared sometimes (Fig. 8-b). The smooth-surfaced endoplasmic reticulum in SHRSP was well developed and showed more vesicular or dilated appearance (Fig. 10). The Golgi apparatus, ribosomes and glycogen particles were also observed and prominent in SHRSP (Fig. 9, 10).

Morphometrical findings. Data of various zona glomerulosa cell components are




Fig. 8-a. Zona glomerulosa cells of SHRSP. Numerous lipid droplets (L) are seen. ($\times 7,900$).




Fig. 8-b. Note large lipid droplets (L). ($\times 5,000$).




Fig. 9. Zona glomerulosa cells of SHRSP. N=nucleus, M=mitochondria, L=lipid droplets, G=Golgi apparatus. Note pleomorphism in mitochondria. ($\times 12,000$).




Fig. 10. Zona glomerulosa cell of SHRSP. Mitochondria (M) have electron-dense matrix and tubular cristae. Lipid droplets (L) are surrounded by smooth-surfaced endoplasmic reticulum (arrow). ($\times 34,000$).

given in Table 2. The average width of the zona glomerulosa of SHRSP ($60.7 \pm 1.60 \mu\text{m}$) was significantly ($p < 0.001$) greater than that of WKY ($49.0 \pm 1.03 \mu\text{m}$). The width of the zona glomerulosa in SHRSP with cerebral lesions ($67.0 \pm 8.03 \mu\text{m}$) was greater than that in SHRSP without cerebral lesions ($54.2 \pm 12.58 \mu\text{m}$). However, this difference was not statistically significant because the number of each group was too small to compare.

The area of the zona glomerulosa cells of SHRSP ($75.4 \pm 2.15 \mu\text{m}^2$) was significantly ($p < 0.001$) larger than that of WKY ($62.7 \pm 2.28 \mu\text{m}^2$), whereas the area of nucleus did not differ significantly from each other (SHRSP: $23.4 \pm 0.56 \mu\text{m}^2$ vs WKY: $22.1 \pm 0.70 \mu\text{m}^2$).

The area of mitochondria of SHRSP ($0.28 \pm 0.006 \mu\text{m}^2$) was significantly ($p < 0.001$) lesser than that of WKY ($0.39 \pm 0.010 \mu\text{m}^2$). And the percent area of the mitochondrial compartment of SHRSP ($15.6 \pm 0.38 \%$) was significantly ($p < 0.001$) lesser than that of WKY ($21.6 \pm 0.60 \%$).

Lipid droplets were observed on 90% of the zona glomerulosa cells in SHRSP and on 69% of that in WKY. Although the area of lipid droplets did not differ significantly from each other (SHRSP: $0.80 \pm 0.031 \mu\text{m}^2$ vs WKY: $0.88 \pm 0.052 \mu\text{m}^2$), the percent area



Fig. 11-a. Zona glomerulosa cell of SHRSP. Note regular arrays of lamellar cristae in elongated mitochondria. ($\times 48,000$).



Fig. 11-b. Zona glomerulosa cell of SHRSP. Note electron-dense intramitochondrial droplets (arrow). ($\times 48,000$)

Table 2 Synopsis of morphometric parameters of the zona glomerulosa

	SHRSP	WKY	p
Width of the zona glomerulosa (μm)	60.7 ± 1.60	49.0 ± 1.03	<0.001
Area of cells (μm^2)	75.4 ± 2.15	62.7 ± 2.28	<0.001
Area of nucleus (μm^2)	23.4 ± 0.56	22.1 ± 0.70	NS
Area of mitochondria (μm^2)	0.28 ± 0.006	0.39 ± 0.010	<0.001
Percent area of mitochondria (%)	15.6 ± 0.38	21.6 ± 0.60	<0.001
Area of lipid droplets (μm^2)	0.80 ± 0.031	0.88 ± 0.052	NS
Percent area of lipid droplets (%)	21.6 ± 1.09	16.9 ± 1.40	$0.001 < <0.01$

Each value represents the group mean \pm SE
p, level of significance of the difference between SHRSP and WKY
NS, not significant

of lipid droplets compartment of SHRSP (21.6 ± 1.09 %) was significantly ($0.001 < p < 0.01$) greater than that of WKY (16.9 ± 1.40 %).

DISCUSSION

It is postulated that the adrenal gland plays an important role to regulate the development or maintenance of hypertension in SHR. TSUCHIYAMA et al.³¹⁾ described the morphological alterations of the adrenal cortex in SHR histochemically and ultra-structurally, and pointed out that these changes were seen even in prehypertensive stage. NICKERSON showed by quantitative morphological techniques that the zona glomerulosa and the zona fasciculata of 21 week old male SHR²²⁾ and the zona reticularis of aging female SHR²³⁾ differed from those of WKY. Another morphological studies on the zona glomerulosa with relation to its functional alterations have been reported in SHR. The present study is the first description of the zona glomerulosa in SHRSP.

In this paper, the zona glomerulosa of SHRSP was characteristically increased in width. Morphometrically the average width of the zona glomerulosa of SHRSP was significantly greater than that of WKY. This widened zone of SHRSP consisted of hypertrophy and hyperplasia of the zona glomerulosa cells. The width of the zona glomerulosa in SHR was also greater than that of controls²²⁾. The increased zona glomerulosa was observed in a response to ACTH administration²⁴⁾, renovascular hypertension²⁸⁾, renal encapsulation-induced hypertension¹³⁾ and sodium-depletion¹¹⁾. Biochemically, the renin-angiotensin system of SHRSP was activated¹⁵⁾¹⁸⁾¹⁹⁾²⁹⁾, but PAC was not elevated²⁵⁾, and participated in maintaining high blood pressure. MATSUNAGA et al.¹⁸⁾ and KAWASHIMA et al.¹⁵⁾ pointed out the close relationship between PRA and cerebral lesions in SHRSP, that is, when the higher PRA was found in SHRSP the higher incidence of cerebral lesions was observed. It is of interest that the width of the zona glomerulosa in SHRSP with cerebral lesions was greater than that in rats without cerebral lesions, though it was not statistically significant. Considering these facts, the activated renin-angiotensin system may play an important role²⁸⁾ to induce hypertrophy and hyperplasia of the zona glomerulosa in SHRSP.

The area of the zona glomerulosa cells of SHRSP was significantly greater than that of WKY. The significant increase of the area of cells in SHRSP may probably due to increased lipid droplets compartment because the area of nucleus and the percent area of mitochondria were not increased in these cells. The smooth-surfaced endoplasmic reticulum and the Golgi apparatus, though they were not measured, seemed to be not effective enough to increase the area of cells.

Under the condition of sodium or magnesium depletion and ACTH administration the mitochondria are increased in number and in size, and the smooth-surfaced endoplasmic reticulum and the Golgi apparatus appear to be well developed. On the other hand, lipid droplets tend to be decreased in number and in size⁶⁾⁸⁾¹¹⁾¹⁴⁾. Opposite alter-

ations of these organelles have been observed under a hypofunctional state by hypophysectomy²⁴⁾, sodium supplement¹²⁾ or aldosterone administration⁸⁾.

There are at least two steps in the biochemical process of aldosterone formation which involve mitochondrial enzymes⁷⁾. One is the transformation of cholesterol to pregnenolone and the other, from 18-hydroxycorticosterone to aldosterone⁵⁾.

In this study on the zona glomerulosa cells of SHRSP, the mitochondria were small in area and pleomorphic in shape but increased in number. NICKERSON²²⁾ observed a decreased volume of mitochondria also in SHR. YAMORI et al.³³⁾ and MOLL et al.²⁰⁾ reported impairments of mitochondrial enzymes in SHR. Therefore, above-mentioned changes of mitochondria may be induced by the impairment of mitochondrial enzymes or the impairment of maturation of mitochondria.

The presence of straight tubular cristae in some mitochondria has been reported in the case of functional solicitation; sodium depleted animals¹¹⁾, aldosterone administration⁸⁾, hypophysectomy in animals¹⁰⁾ and primary aldosteronism in human³⁰⁾. This type of cristae was also described in the mitochondria of a few cells of the zona glomerulosa in normal rats¹⁰⁾. At the present study this alteration was found not only in SHRSP but also in WKY.

The percent area of lipid droplets compartment of the zona glomerulosa cells in SHRSP was significantly increased, but the area of lipid droplets in SHRSP did not differ from in WKY. AOKI¹⁾, TSUCHIYAMA et al.³¹⁾ and FRINK et al.⁹⁾ reported that lipid droplets of the zona glomerulosa cells in SHR were increased. On the other hand, a decreased volume of lipid droplets in SHR was shown quantitatively by NICKERSON²²⁾. The cause of this difference is not known but it will be related with diet, strain or developmental environment.

The smooth-surfaced endoplasmic reticulum tended to be well developed in SHRSP. β -hydroxysteroid dehydrogenase system and 21-hydroxylase system are located within the smooth-surfaced endoplasmic reticulum³²⁾. Therefore, these organelles may reflect the activity of the zona glomerulosa cells.

The Golgi apparatus also appeared to be hypertrophic in SHRSP. This organelle takes a part in conjugation of steroid hormone¹⁶⁾. The changes of the smooth-surfaced endoplasmic reticulum and the Golgi apparatus in SHRSP may be the response to an activated state of the zona glomerulosa cells.

These fine structural alterations may indicate the following possibilities. The zona glomerulosa cells in SHRSP are stimulated in the steroidogenesis by elevated demands for aldosterone. It may be supported by the fact that PRA in SHRSP was elevated¹⁵⁾¹⁸⁾. And so a lot of cholesterol are taken into the zona glomerulosa cells as a precursor of aldosterone. Then, if the mitochondria of SHRSP have some kinds of dysfunction, it may lead to the increase in number of the mitochondria to compensate for unfavorable function. But it may be limited to spend cholesterol because the impairment of mitochondrial function causes an insufficient conversion to steroid hormone. Therefore, lipid droplets where cholesterol is stored²¹⁾ are increased in number in cytoplasm. The significant

increase of cell area may be induced by increased lipid droplets. And the zona glomerulosa cells may have undergone hypertrophy to hyperplasia.

Biochemically, though PRA of SHRSP was elevated¹⁵⁾¹⁸⁾ OGINO et al.²⁵⁾ reported that PAC of SHRSP did not differ from that of WKY at 5 month old. These results will support the speculation in the present study.

However, further studies on the renin-angiotensin aldosterone system in SHRSP are necessary to understand the pathogenesis of hypertension.

ACKNOWLEDGEMENT

The author wish to thank greatly Dr. H. TSUCHIYAMA, Professor of Pathology, and Drs. K. KAWAI, Associate Professor of Pathology, Nagasaki University for their suggestive and valuable criticism and comments. And the author also express my gratitude to Dr. M. OZAKI, Professor of Pharmacology, and Dr. M. NIWA, Associate Professor of Pharmacology, Nagasaki University for their kind provision for rats. The co-operative research students and skillful technical assistants in the Department of Pathology, Nagasaki University are acknowledged.

REFERENCES

- 1) AOKI, K., TANKAWA, H., FUJINAMI, T., MIYAZAKI, A. and HASHIMOTO, Y.: *Jpn. Heart J.* 4: 426-442, 1963.
- 2) AOKI, K.: *Jpn. Heart J.* 4: 443-461, 1963.
- 3) BAER, L., KNOWLTON, A. and LARAGH, J. H.: Spontaneous hypertension: its pathogenesis and complications (OKAMOTO, K. ed.). p. 203, Igaku Shoin Ltd., Tokyo, 1972.
- 4) BARTSCH, G., BAUMGARTNER, U. and ROHR, H. P.: *Path. Res. Pract.* 162: 291-300, 1978.
- 5) CRISTENSEN, A. K. and GILIM, S. W.: The gonads (McKERNNS, K. W. ed.). Appelton Century Crafts, New York, 1969.
- 6) DOMOTO, D. T., BOYD, J. E., MULROW, P. J. and KASHGARIAN, M.: *Am. J. Pathol.* 72: 433-446, 1973.
- 7) FAWCETT, D. W., LONG, J. A. and JONES, A. L.: *Progr. Hormone Res.* 25: 315-380, 1969.
- 8) FISHER, E. R. and HORVAT, B.: *Arch. Pathol.* 92: 172-179, 1971.
- 9) FRINK, R., KRUPP, P. P., ROOT, D. T. and WARSHAW, D. M.: *Expl. Cell Biol.* 48: 245-254, 1980.
- 10) FUJITA, H.: *Z. Zellforsch.* 125: 480-496, 1972.
- 11) GIACOMELLI, F., WIENER, J. and SPIRO, D.: *J. Cell Biol.* 26: 499-521, 1965.
- 12) HIRANO, T.: *Acta Pathol. Jpn.* 26: 589-601, 1976.
- 13) KASEMSRI, S. and NICKERSON, P. A.: *Am. J. Pathol.* 82: 143-156, 1976.
- 14) KAWAI, K., SUGIHARA, H. and TSUCHIYAMA, H.: *Acta Pathol. Jpn.* 28: 265-278, 1978.

- 15) KAWASHIMA, K., SHIONO, K. and SOKABE, H.: *Clin. Exp. Hyp.* 2:229-245, 1980.
- 16) LONG, J. A. and JONES, A. L.: *Lab. Invest.* 17: 355-370, 1967.
- 17) MARUYAMA, T.: *Jpn. Circul. J.* 33: 1271-1284, 1969.
- 18) MATSUNAGA, M., YAMAMOTO, J., HARA, A., YAMORI, Y., OGINO, K. and OKAMOTO, K.: *Jpn. Circul. J.* 39: 1305-1311, 1975.
- 19) MATSUNAGA, M., KOMURO, T., YAMAMOTO, J., HARA, A., MORIMOTO, K. and YAMORI, Y.: *Jpn. Heart J.* 21: 738-751, 1980.
- 20) MOLL, D., DALE, S. L. and MELBY, J. C.: *Endocrinology* 96: 416-420, 1975.
- 21) MOSES, H. L.: *Science* 163: 1203-1205, 1969.
- 22) NICKERSON, P. A.: *Am. J. Pathol.* 84: 545-560, 1976.
- 23) NICKERSON, P. A., FELD, L. G. and VANLIEW, J. B.: *Am. J. Pathol.* 97: 433-448, 1979.
- 24) NUSSDORFER, G. G., MAZZOCCHI, G. and REBUFFAT, P.: *Endocrinology* 92: 141-151, 1973.
- 25) OGINO, K., HARA, A., YAMAMOTO, J. and MATSUNAGA, M.: Spontaneous hypertension: its pathogenesis and complications, Proc. 2nd Int. Symp. on SHR. pp. 352-353, DHEW Publication No. (NIH) 77-1179, 1976.
- 26) OKAMOTO, K. and AOKI, K.: *Jpn. Circul. J.* 27: 283-293, 1963.
- 27) OKAMOTO, K., YAMORI, Y. and NAGAOKA, A.: *Circ. Res.* 34-35, Suppl. 1: 143-153, 1974.
- 28) REBUFFAT, P., BELLONI, A. S., MAZZOCCHI, G., VASSANELLI, P. and NUSSDORFER, G. G.: *J. Anat.* 129: 561-570, 1979.
- 29) SHIBOTA, M., NAGAOKA, A., SHINO, A. and FUJITA, T.: *Am. J. Physiol.* 236: H409-H416, 1979.
- 30) SYMINGTON, T.: Functional pathology of the human adrenal gland. p. 132, E&S Livingstone Ltd., London, 1969.
- 31) TSUCHIYAMA, H., SUGIHARA, H. and KAWAI, K.: Spontaneous hypertension: its pathogenesis and complications (OKAMOTO, K. ed.). pp. 177-184, Igaku Shoin Ltd., Tokyo, 1972.
- 32) VOLK, T. L.: *Lab. Invest.* 25: 349-355, 1971.
- 33) YAMORI, Y., OOSHIMA, A. and OKAMATO, K.: *Jpn. Heart J.* 14: 162-164, 1973.