

# The Effect of Epidermal Growth Factor, Basic Fibroblast Growth Factor, Transforming Growth Factor- $\beta$ , and Insulin on the DNA Synthesis of Renal Cell Carcinoma Cell Lines.

Naoki NISHIMURA

*Department of Urology, Nagasaki University School of Medicine,  
Nagasaki, 852, Japan.*

*Received for publication, January 28, 1991*

**SUMMARY:** In this experiment, the effect of various growth factors (epidermal growth factor, basic fibroblast growth factor, transforming growth factor- $\beta$  and insulin) on the DNA synthesis of three renal cell carcinoma cell lines (ACHN, VMRC-RCW, NT) has been investigated in a serum free condition. These growth factors stimulated the DNA synthesis of all renal cell carcinoma cell lines dose-dependently. Transforming growth factor- $\beta$ , a known growth inhibitor for renal tubular cells, stimulated the DNA synthesis of renal cell carcinoma cells. The conditioned medium (which did not include any serum) contained very little autocrine growth factor for renal cell carcinoma cell itself. These results suggest that paracrine growth factors are mostly related to the growth of renal cell carcinoma cells than autocrine growth factor. The renal cell carcinoma cells, which are the transformed form of renal tubular cells and due to this transformed character, TGF- $\beta$  which is basically a growth inhibitor for tubular cell but stimulates the renal cell carcinoma cell.

## INTRODUCTION

It has been reported that different growth factors are present in the extracts of normal and malignant tissues<sup>1, 2)</sup>, blood<sup>3)</sup> and urine<sup>4)</sup>. These growth factors can stimulate or inhibit cellular differentiation as well as they play important role in cellular proliferation. Growth factors have also been identified from conditioned media of hepatoma<sup>5)</sup>, prostatic cancer<sup>6)</sup>, colon carcinoma<sup>7)</sup> and renal cell carcinoma cell lines<sup>8)</sup>. Growth factors stimulate cancer cell proliferation by the autocrine or paracrine manner. It is reported that renal cell carcinoma was originated from renal tubular cells, especially from proximal tubular cells<sup>9)</sup>. If we can make a comparative study on the response of different growth factors on renal cell carcinoma cells and

normal renal tubular cells, than it will be possible to understand the mechanism of growth of cancer cells. In this experiment, we have investigated the production of autocrine growth factors by renal cell carcinoma cell lines using the serum free culture system and studied the effects of different exogenous growth factors on these cell lines.

## MATERIALS AND METHODS

Materials.

EGF was purified from submaxillary glands of male mice using the method described by Savage and Cohen<sup>1)</sup>. Bovine basic FGF was obtained from TOYOBO, Osaka, Japan. Insulin was purchased from Sigma, St. Louis, MO. Human TGF- $\beta$  was from Wako Pure Chemicals, Osaka, Japan. Dulbecco's modified Eagle's






Fig. 2. Effect of the conditioned medium on DNA synthesis of renal cell carcinoma cell lines.

**Table 1.** Effects of various growth factors on DNA synthesis of renal cell carcinoma cell lines.

addition	DNA synthesis (cpm)		
	ACH-N	VMRC-RCW	NT
none	40493 ± 1124	2617 ± 309	16642 ± 958
EGF 1 ng/ml	79458 ± 1448	3199 ± 988	29372 ± 410
bFGF 2 ng/ml	64090 ± 3657	3556 ± 90	26739 ± 114
TGF- $\beta$ 1 ng/ml	53767 ± 3859	3186 ± 526	23278 ± 260
insulin $10^{-7}$ M	49436 ± 127	3421 ± 255	19511 ± 12
10%FCS	50375 ± 2371	11313 ± 657	22607 ± 1039

Experimental conditions were as described in "Materials and Methods". Values are expressed as means  $\pm$  SD for triplicated experiences.

## DISCUSSION

In order to see the effect of different growth factors on RCC cell line, we adopted the DNA synthesis assay method rather than the cell count technique. In the cell count technique, cells may die or detach from the dish that might give false result. In this experiment, we used the serum free culture system, because there are different growth factors present in the serum that might act as a cofactor which may have a direct stimulatory effect to exogenous growth factors.

According to these results all these RCC cell lines were stimulated by exogenous growth factors better than autocrine growth factors under the serum free condition. EGF, IGF-I and FGF have been reported to stimulate the growth and TGF- $\beta$  has been reported to inhibit the

growth of cultured renal tubular cells<sup>12, 13, 14</sup>. But our results showed that TGF- $\beta$  have weakly stimulated the growth of RCC cells. TGF- $\beta$  inhibits the DNA synthesis of normal tubular cells, but when these tubular cells are transformed to RCC cells, then this inhibitory effect is altered and TGF- $\beta$  acts as a stimulatory factor to RCC cells. The same results were obtained when hepatic cells<sup>15</sup> and bronchial epithelial cells<sup>16</sup> were investigated. It is probable that the abolition of inhibitory effect of growth factors to normal cells might allow them to transform to malignant cells. Our result showed EGF has strongly stimulated the growth of RCC cell. Transforming growth factor- $\alpha$ , which is considered as a transformed type of EGF and acts through the receptor<sup>17</sup> of EGF, stimulates the growth of RCC cells. Renal cell carcinoma is usually a hypervascular tumor. It is probable that FGF stimulates the growth of endothelial




Fig. 3. Dose-response effect of EGF, basic FGF, TGF- $\beta$ , and insulin on DNA synthesis of renal cell carcinoma cell lines.

cells and participates in the neovascularization of RCC. FGF also stimulates the growth of RCC cells. Although it was reported that RCC cells produced autocrine growth factors<sup>8, 18)</sup>. We could not find the production of autocrine growth factor by RCC cells in our experiment. In the presence of serum, RCC cells produce IL-6 in culture medium<sup>18)</sup>. It seems that for the production of autocrine growth factor by RCC

cells in culture, same triggers are needed. Further study should be directed to find our the trigger for the production of autocrine growth factors by RCC cells.

#### ACKNOWLEDGMENTS

I wish to thank Prof. Y. Saito, Dr. S. Kanda and Dr. P. K. Saha for helpful discussion

through out the study. I also thank Mr. T. Shimogama and Miss M. Yoshimoto for their outstanding technical assistance.

### REFERENCES

- 1) Cohen, S.: *J. Biol. Chem.* **237**:1555 (1962).
- 2) Gospodarowicz, D.: *Nature* **249**:123 (1974).
- 3) Kohler, N. and Lipton, A.: *Exp. Cell Res.* **87**:297 (1974).
- 4) Cohen, S. and Carpenter, S.: *Proc. Natl. Acad. Sci. USA.* **72**:1317 (1975).
- 5) Peter, O., Mona, L. C., Philip, S., Antonio, F. and Thomas, E. S.: *J. Cell physiol.* **131**:165 (1987).
- 6) T. Ikeda, Mario, N. L. and Hans, M.: *Biochemistry* **26**:2406 (1987)
- 7) Jean-Michel, C., Françoise, G., Maryse, R., Dominique, B., Jacques, M. and Gilbert, P.: *Int. J. Cancer.* **42**:895 (1988).
- 8) T. Nakamoto, A. Usui, K. Oshima, H. Ikemoto, S. Mitani and T. Usui.: *Biochem. Biophys. Res. Commun.* **153**:818 (1988).
- 9) H. Holthofer, A. Miettinen, R. Paasivuo, V. P. Lehto, E. Linder, O. Alfthan and I. Virtanen.: *Lab. Invest.* **49**:317 (1983).
- 10) Borden, E. C., Hogan, T. F. and Voelkel, J. G.: *Cancer Res.* **42**:4948 (1982).
- 11) N. Tsuda, Y. Saito.: *Saibou* **15**:161 (1983). (Japanese)
- 12) S. Kanda, K. Nomata, Saha, P. K., N. Nishimura, J. Yamada, H. Kanetake and Y. Saito.: *Cell Biol. Int. Rep.* **13**:687 (1989).
- 13) Fine, J. A., Holley, R. W., Nasri, H. and Badie-Dezfooly, B.: *Proc. Natl. Acad. Sci.* **82**:6163 (1982).
- 14) S. Kanda, T. Igawa, M. Taide, Saha, P. K., N. Nishimura, K. Nomata, H. Kanetake and Y. Saito.: *Cell Biol. Int. Rep.* **14**:877 (1990).
- 15) T. Nakamura, Y. Tomita, R. Hirai, K. Yamaoka, K. Kaji and A. Ichihara.: *Biochem. Biophys. Res. Commun.* **133**:1042 (1985).
- 16) T. Masui, Wakefield, L. M., Lechner, J. F., LaVeck, M. A., Sporn, M. B. and Harris, C. C.: *Proc. Natl. Acad. Sci. USA.* **83**:2438 (1986).
- 17) Sporn, M. B. and Roberts, A. B.: *Nature* **313**:745 (1985).
- 18) Miki, M. Iwano, Y. Miki, M. Yamamoto, Bo tang, K. Yokokawa, T. Sonoda, T. Hirano and T. Kishimoto.: *FEBS Letters* **250**:607 (1989).