

Predicting West Nile Virus Infection Risk from the Synergistic Effects of Rainfall and Temperature

Journal:	<i>Journal of Medical Entomology</i>
Manuscript ID	JME-2015-0246.R3
Manuscript Type:	Research Article
Date Submitted by the Author:	16-Mar-2016
Complete List of Authors:	Shand, Lindsay; University of Illinois, Statistics Brown, William; University of Illinois at Urbana-Champaign, Pathobiology Chaves, Luis; Nagasaki University, Institute of Tropical Medicine (NEKKEN), Department of Vector Ecology and Environment; Tomonori, Goldberg, Tony; University of Wisconsin, Pathobiological Sciences Hamer, Gabriel; Texas A&M, Entomology Haramis, Linn; Illinois Dept of Public Health, Division of Environmental Health Kitron, Uriel; Emory University, ENVS Walker, Edward; Michigan State University, Microbiology and Molecular Genetics Dept Ruiz, Marilyn; University of Illinois at Urbana-Champaign, College of Veterinary Medicine
Please choose a section from the list:	Vector-Borne Diseases, Surveillance, Prevention
Field Keywords:	West Nile Virus, Modeling, Public Health Entomology
Organism Keywords:	Culex

1 Shand et al.: WNV risk prediction
2 from rainfall and temperature
3
4 Journal of Medical Entomology
5 Full-length research article, Vector-Borne Diseases,
6 Surveillance, Prevention
7
8
9
10
11

Marilyn O'Hara Ruiz
Department of Pathobiology
2001 South Lincoln Avenue
Urbana, Illinois 61802
217-265-5115 (office phone)
217-244-7421 (office FAX)
moruiz@illinois.edu

12 Predicting West Nile Virus Infection Risk from the Synergistic Effects of Rainfall and
13 Temperature
14

15 L. Shand,¹ W.M. Brown,² L. F. Chaves,³ T.L. Goldberg,⁴ G. L. Hamer,⁵ L. Haramis,⁶ U. Kitron⁷ E. D.
16 Walker⁸ and M.O. Ruiz²

17
18 ¹Department of Statistics, University of Illinois, Urbana, IL 61801

19 ²Department of Pathobiology, University of Illinois, Urbana, IL 61801

20 ³Institute of Tropical Medicine (NEKKEN), Nagasaki University, Japan

21 ⁴Department of Pathobiological Sciences, University of Wisconsin, Madison, WI 53706

22 ⁵Department of Entomology, Texas A&M University, College Station, TX 77843

23 ⁶Division of Environmental Health, Illinois Department of Public Health, Springfield, IL 62761

24 ⁷Department of Environmental Sciences, Emory University, Atlanta, GA 30322

25 ⁸Department of Microbiology and Molecular Genetics, Michigan State University, East Lansing,
26 MI 48824

27

28

29

Abstract

30 Mosquito-based surveillance is a practical way to estimate the risk of transmission of West Nile
31 virus (WNV) to people. Variations in temperature and precipitation play a role in driving
32 mosquito infection rates and transmission of WNV, motivating efforts to predict infection rates
33 based on prior weather conditions. Weather conditions and sequential patterns of
34 meteorological events can have particularly important, but regionally distinctive, consequences
35 for WNV transmission, with high temperatures and low precipitation often increasing WNV
36 mosquito infection. Predictive models that incorporate weather can thus be used to provide
37 early indications of the risk of WNV infection. The purpose of this study was: first, to assess the
38 ability of a previously published model of WNV mosquito infection to predict infection for an
39 area within the region for which it was developed: and second, to improve the predictive ability
40 of this model by incorporating new weather factors that may affect mosquito development.
41 The legacy model captured the primary trends in mosquito infection, but it was improved
42 considerably when calibrated with local mosquito infection rates. The use of interaction terms
43 between precipitation and temperature improved model performance. Specifically,
44 temperature had a stronger influence than rainfall, so that lower than average temperature
45 greatly reduced the effect of low rainfall on increased infection rates. When rainfall was lower,
46 high temperature had an even stronger positive impact on infection rates. The final model is
47 practical, stable and operationally valid for predicting West Nile virus infection rates in future
48 weeks when calibrated with local data.

49 West Nile virus, climate and weather, risk model, Illinois

50

51

52 Predicting West Nile Virus Infection Risk from the Synergistic Effects of Rainfall and

53 Temperature

54

55 Since the introduction of West Nile virus (WNV) into the Western Hemisphere via New

56 York City in 1999, WNV has spread throughout the Americas and poses an ongoing and serious

57 threat to human and animal health. Over 40,000 cases of illness from WNV were reported

58 through public health surveillance systems in the United States between 1999 and 2014 (CDC

59 2014). The number and location of cases has varied each year, and the ability to predict

60 outbreaks has proven to be challenging. After a period of relatively low activity between the

61 years 2008 to 2011, a large outbreak in 2012, with 5674 human cases reported in the United

62 States, renewed concern about the need for public health preparedness, and spurred efforts to

63 determine better ways to anticipate and reduce the risk of exposure to WNV (Nasci 2013).

64 Mosquito-based surveillance is a recommended, standard, and practical way to estimate

65 the risk of transmission of WNV and other mosquito-borne pathogens to people (Macdonald

66 1956, Moore et al. 1993, Hokit et al. 2013). Select species of mosquitoes in the genus *Culex*

67 comprise over 95 percent of the positive tests for WNV in the United States and are the primary

68 focus of mosquito surveillance efforts (Andreadis 2012). After trapping of blood-fed vector

69 mosquitoes and virus diagnostic testing, the minimum infection rate and the maximum

70 likelihood estimator for mosquito infection rates based on pooled samples are common

71 measures used to estimate the true infection rate (Walter et al. 1980, Hepworth 2005, Gu et al.

72 2003, Biggerstaff 2009, Ebert et al. 2010).

73 Variations in temperature and precipitation play a role in driving the WNV infection rate
74 and transmission, motivating efforts to predict WNV mosquito infection rates from prior
75 weather conditions. Warmer weather increases potential for transmission because it reduces
76 the number of days between virus ingestion to effective transmission (extrinsic incubation
77 period), shortens the length of time between blood meals (gonotrophic cycle), and leads to an
78 earlier start to seasonal mosquito activity (Turell et al. 2001, Dohm 2002, Turell et al. 2005,
79 Reisen et al. 2010, Hartley et al. 2012). Mosquito abundance also generally increases with
80 warmer temperatures, but very hot conditions can have the opposite effect; and shorter
81 lifespans in *Culex* mosquitoes may reduce transmission as fewer individuals live long enough to
82 become infectious (Chaves et al. 2013, Ciota et al. 2014).

83 Hydrologic conditions also affect WNV transmission. *Culex* mosquitoes reproduce in
84 standing water, but heavy rainfall can reduce *Culex* survival both at the adult stage and during
85 larval development (Gardner et al. 2012, Jones et al. 2012). Rainfall influences near-surface
86 humidity, and studies have found that higher humidity induced oviposition in gravid *Culex*
87 *nigripalpus* (Day and Curtis 1999) and *Culex quinquefasciatus* (Chaves and Kitron 2011). Thus,
88 rainfall may increase the potential for pathogen transmission as females seek blood meals prior
89 to oviposition. The frequency, strength, and timing of rainfall events can also affect water
90 chemistry and the degree to which standing water is suitable for mosquito pre-adult
91 development (Shaman and Day 2007, Chaves and Kitron 2011, Gardner et al. 2013).

92 The net result of these effects is that high temperatures combined with low
93 precipitation have often led to higher than average mosquito infection, but these effects vary
94 by region, and the effect of rainfall is especially variable. Weekly patterns of lower than average

95 rainfall and higher temperature, for example, explained about 70 percent of the variability in
96 WNV mosquito infection rates in a study focused on the Chicago, Illinois area (Ruiz et al. 2010).
97 Similarly, drought followed by wet conditions preceded the reporting of WNV human illness in
98 Florida (Shaman et al. 2005;). Drought, during which mosquitoes and birds are in closer
99 proximity due to reduced water availability, could cause local sylvatic amplification of WNV,
100 and subsequent rainfall could then allow dispersal of infected vectors and hosts (Shaman et al.
101 2005). Especially during very hot and dry periods, human-introduced water can create
102 mosquito habitats that might not be otherwise available (Reisen et al. 2008, Barker et al. 2009,
103 Becker, Leisnham, and LaDeau 2014). The relationship between prior rainfall and WNV
104 outbreaks has varied in prior analyses. Outbreaks of WNV in Europe in 2010, for example, were
105 preceded two to four weeks earlier by warmer than average conditions, but the outbreaks were
106 less clearly associated with relative humidity and rainfall (Paz et al. 2013). Similarly, warmer
107 than average winter temperatures and higher than average rainfall preceded the 2012 outbreak
108 in Dallas, Texas, but variables that measured rainfall were not significant in a multivariate
109 analysis (Chung et al. 2013).

110 The purpose of our study was two-fold. First, we assessed the ability of a previously
111 published model of WNV mosquito infection developed for the Chicago region (Ruiz et al. 2010)
112 to predict infection for a subset of that region – specifically for DuPage County, Illinois. For this
113 objective, we compared the measured WNV mosquito infection rate (MIR) for the period from
114 2004 to 2013 with the MIR estimated by a linear model that resulted from the prior work (See
115 Supplementary Materials), referred to henceforth as the “legacy model”. Then, we worked with
116 public health and mosquito abatement personnel in DuPage County in 2014 to learn about the

117 local characteristics of mosquito testing and delivery of public health warnings, so that a
118 predictive model for WNV could be developed and implemented effectively in this setting.
119 Second, we refined the legacy model both to develop a model that takes into account the local
120 conditions and to exploit weather data more fully by considering interaction effects between
121 rainfall and temperature. The broader context of this work is to provide a practical,
122 generalizable, and operationally valid approach to predicting WNV mosquito infection that can
123 be incorporated into public health assessments using data from prior weather conditions.

124 **Materials and Methods**

125 **Study region.**

126 DuPage County, Illinois, is located west of the city of Chicago (Figure 1). It comprises an
127 area of 848 km² and is the second most populous county in the state of Illinois, with a
128 population of 932,126 in 2013 (US Census Bureau). Mosquito control in the county is organized
129 through a combination of mosquito abatement districts, townships, municipalities, and several
130 large landholders. The study period of interest was from 2005 to 2014 and model development
131 included data on weather conditions and mosquito infection rates during this period. All data
132 were organized by week, with weeks starting on Saturday.

133 **Weather data**

134 Daily temperature and precipitation measures were based on two local weather
135 stations: Midway (MDW) and O'Hare (ORD) (Figure 1). Weekly precipitation (rainfall in cm) was
136 calculated from the daily average for each week from the two stations. Weekly temperature
137 was measured as the mean of the temperature (°C) from the daily temperature readings from
138 the two stations. Temperature data were further used to calculate a variable called a "Degree

139 Week" (*DW*) constructed similarly to the more common Degree Day, but with differences
140 accumulated over weeks, rather than days (Ruiz et al. 2010). The *DW* is the cumulative sum of
141 the difference of all prior weekly temperatures from a threshold value of 22°C. The
142 temperature threshold of 22°C was used because compared to other values, it led to the
143 highest correlation between *DW* and the weekly local MIR based on cross-correlations across a
144 range of threshold values from 10° to 24°C and time lags from one to ten weeks (Baker et al.
145 1984, Curriero et al. 2005, Kunkle et al. 2006). For a given week: $\Delta DW = T_{\text{mean}} - T_{\text{base}}$ if the weekly
146 Temperature (T_{mean}) is greater than the threshold ($T_{\text{base}} = 22^\circ\text{C}$) and 0 otherwise. To remove the
147 seasonal trend from the model, weather variables were measured as the weekly differences
148 from the 30-year Climate Normals for 1981-2010, provided by the U.S. National Weather
149 Service (Figure 2). These differences captured the patterns outside the seasonal trends and
150 focused the analysis on characterizing how weeks differed from the expected values. We also
151 considered variables that measured the prior year's average precipitation as was done in Ruiz
152 et al. (2010). To improve our understanding of this relationship with *MIR*, we considered the
153 effect of the average precipitation for four equal parts of the prior year starting with week 1,
154 rather than the year as a whole.

155 **Mosquito data**

156 The results of mosquito pools tested for WNV during the years from 2005 to 2013 from
157 specimens and collected from gravid traps located in DuPage County were provided by the
158 Illinois Department of Public Health (IDPH). These data were submitted to the IDPH Web Portal,
159 where Illinois agencies upload WNV mosquito test results. For 2014 data, mosquito test results
160 were received directly from the DuPage County Department of Public Health. Test results were

161 selected to include only the most common female vector species mosquitoes, which in this
162 region are *Culex pipiens* and *Culex restuans* (Hamer et al. 2008, Andreadis 2012). PCR and
163 VecTests were reported from 2005 to 2007 and PCR and RAMP tests, from 2009 to 2014. PCR
164 tests comprised from 49 percent to 65 percent of all samples, depending on the year. The IDPH
165 protocol stipulates pool sizes no larger than 50 individuals, and 19,115 (99 percent) of the
166 19,345 pools tested were within this guideline. The number of gravid trap locations in the study
167 region during the years of interest varied from 136 trap locations in 2007 to 72 in 2014 (Figure
168 1). Test result data were grouped by week and the *MIR* was calculated for a given week where:
169 $\text{minimum infection rate} = 1000 * (\text{number of positive pools}) / (\text{total number of mosquitoes in}$
170 $\text{pools tested})$, using the CDC Excel Add-in for pooled infection rates (Biggerstaff 2009). As with
171 the weather data, the *MIR* variable was calculated as the difference from the countywide
172 average *MIR* from 2005-2013 (Figure 2).

173 **Model development**

174 To determine how well the legacy model published in Ruiz et al. (2010) performed for
175 DuPage County alone, we first used the coefficients from the weather-only (*MIR* independent)
176 version of this model and local weather station data and compared visually the actual *MIR* for
177 DuPage County with the predicted weekly *MIR* values. For the new model, initially, we
178 considered all weather variables - including 1-8 week lags of temperature and rainfall and the
179 prior year's precipitation measured in quarters, halves, and the full year. We used Pearson's
180 correlation *r* values to assess the strength of associations between weather variables and *MIR*
181 at different time lags to determine how far back in time to include weekly lagged weather

182 variables and to determine the relative strength of the associations with prior seasons'
183 precipitation.

184 Using the same general approach as the legacy model, we developed new linear
185 regression models to predict the weekly DuPage County *MIR*. All models were fitted using the
186 least squares method with the R package *stats* (R Core Team 2013). We selected the model
187 variables using adjusted R^2 (R^2_{adj}) and Akaike Information Criteria (AIC) with both backward and
188 forward stepwise regression with a significance level threshold of $\alpha=0.1$. Calendar weeks 18-38
189 (from the end of April to mid-September) from each year were used to develop the model. Data
190 were treated as a weekly time series, with weekly weather data starting four weeks prior to the
191 *MIR* data, to account for the temporal lags prior to the first *MIR* measurement in week 18.

192 We investigated the effect of the temporal autocorrelation of *MIR* by developing *MIR*
193 lag dependent models that included prior levels of *MIR* to predict future levels. We then added
194 all interaction terms between the temperature and precipitation weekly lagged weather
195 variables in interactions models. One important practical goal was to determine if it was
196 possible to use the *MIR* measured from mosquitoes collected and tested during the current
197 season for real-time predictions. Thus, we compared four model types in the model
198 development phase: *MIR* dependent models without and with interaction terms, and *MIR*
199 independent models without and with interaction terms.

200 The new models for DuPage County were fitted initially using data from the years 2005
201 through 2012, while data from 2013 and 2014 were used to test the models' predictive ability.
202 Since the difference from the weekly average *MIR* was used to fit the model, the *MIR* weekly
203 averages were added to the model estimates to produce the predicted *MIR* values. The

204 predicted residual sum of squares (PRESS), calculated as the sum of squared errors of out-of-
205 sample prediction values for 2005 to 2013, was used as a measure to compare the model
206 predictions (Chaves and Pascual 2007). Out-of-sample predictions were made by randomly
207 dropping one weekly observation at a time to predict, while using the remainder of the data to
208 fit the model. Once we selected the best model for DuPage County and were reaching the end
209 of the 2014 mosquito season, we refit the model including the year 2013 data to recalculate
210 and improve the models' coefficients. Finally, we compared the best new local model with the
211 legacy model predictions, using the mean square prediction error (MSPE) and standard error
212 (SE) of MSPE for model prediction for the year 2014, a year that was not used to fit the
213 coefficients of either of the two models.

214

215

Results

Data exploration

217 During the study period, the three years with the highest rates of human illness in
218 DuPage County were 2005, 2006, and 2012, with at least 40 or more cases of WNV illness
219 (Table 1). These years also had high average mosquito infection rates of 5.57, 6.88, and 8.74
220 respectively. In the two years 2010 and 2013, average *MIR* was similar to the years with more
221 human illness, but the peak *MIR* week was later. Weekly precipitation was often lower in the
222 three weeks prior to the peak *MIR* and the DW temperature higher at the peak *MIR* week
223 during higher *MIR* years.

224 The initial comparison between the actual *MIR* and predicted *MIR* using the legacy
225 model for DuPage County indicated that the model captured the main trend of infection rates

226 but did not always correctly estimate the amplitude or timing of mosquito infection, especially
227 in years with low infection rates (Figure 3A).

228 Weekly average precipitation showed moderate correlation with *MIR*. The assessment
229 of correlations between weather variables and *MIR* at different weekly time lags determined
230 that the average weekly precipitation and *DW* were most strongly correlated with *MIR* at lags 1-
231 4, with correlation dropping after a 4-week lag (Figure 4). The correlation between *MIR* and *DW*
232 was particularly strong at short lags and showed a clear pattern of decreasing correlation with
233 increased time lag. The Pearson's correlations between *MIR* and lagged *MIR* were 0.88 (n=209,
234 $p < 0.0001$) at one week and 0.73 (n=208, $p < 0.0001$) at two weeks. We found that the average
235 precipitation of weeks 27-39 of the previous year showed the highest negative correlation with
236 *MIR* (Table 2). Therefore, we considered this variable in the new DuPage County *MIR* model.

237 **Model selection**

238 After observing the timing of data availability following mosquito collection and testing
239 in the county, we determined that the data for a 2-week lagged autoregressive *MIR* term may
240 be available for use in a real-time prediction model, but the data would not be available in time
241 for including the 1-week lagged *MIR*. Initial model diagnostics revealed additional temporal
242 correlation among the residuals, even after the seasonal de-trending of *MIR*. Thus, we also
243 included the temporal variable *week* as a predictor. *Week* was a more significant and influential
244 variable in the *MIR* independent models. The four best models, after AIC variable selection,
245 based on the R^2_{adj} and the smallest AIC included variables significant with $p < 0.1$. As a last step,
246 the least important interaction terms were also excluded in cases where the model fits were

247 not significantly changed as a result. From six to nineteen variables were selected for the four
248 models (Table 3).

249 Comparing model structures, two things became clear. First, in the models that included
250 prior *MIR* (models 1 and 3), *MIR* had an exceptionally strong effect on the model prediction in
251 all cases; and second, the interaction terms significantly improved the model fit. The *MIR*
252 dependent model with interaction effects (model 3) explained the most variation ($R^2_{\text{adj}} = 0.721$;
253 $\text{AIC} = 315.2$), but the strong contribution of the interaction terms was seen especially when *MIR*
254 from previous weeks was not included in the model (comparison of model 2 and model 4).
255 Though the autoregressive *MIR* term was an important factor statistically, we determined that
256 its inclusion in the model could overwhelm the effect of weather on *MIR* prediction. In other
257 words, the *MIR* autoregressive terms tended to mimic the prior weeks' *MIR* pattern, making
258 predictions less sensitive to actual changes in weather.

259 Based on these observations, we decided that the *MIR* independent models were
260 preferred. They modeled more clearly the relationship of next week's *MIR* with the weather
261 variables and obviated the need to wait for field-based collections and testing. We then
262 considered whether implementation of the more complex interaction term model was
263 warranted over a simpler model. For DuPage County, the best independent main effects model
264 (model 2) explained only about half of the variation in *MIR* ($R^2_{\text{adj}}=0.451$; $\text{AIC}=409.0$), whereas
265 the best *MIR* independent 2nd order interaction model (model 4) explained 66 percent
266 ($\text{AIC}=353.5$). For these reasons, we selected the latter model for implementation. The larger
267 PRESS statistic of 0.537 of model 4 also showed the strength of this model over model 2
268 ($\text{PRESS}=0.481$). Quadratic terms were also tested in model construction due to a possible

269 quadratic relationship between *MIR* and *DW* lags seen in the exploratory analysis, but did not
270 improve the model. Finally, comparing the best local model with the legacy model, the 2014
271 MSPE and SE of the new local DuPage model was 4.54 and 1.70 respectively, which was lower
272 (less error) when compared to the legacy model MSPE of 5.52 and SE 2.08.

273 **Model inference**

274 The best model's final variable selection included weekly precipitation (*prec*) at 1 to 3
275 week lags, weekly *DW* at 1 to 4 week lags, the average precipitation in the third quarter of the
276 previous year (*previous year prec. weeks 27-39*), and 9 interaction terms, for a total of 17
277 factors (Table 3). All of the included terms had significant effects at $\alpha=0.1$ on *MIR* predictions.
278 As with the legacy model, *DW* had a larger overall effect on infection rate than precipitation.

279 Considering the overall effect of the weather variables, an increase in average *DW* in the
280 4 prior weeks led to higher than average infection rate estimates. Precipitation effects varied,
281 however, with a positive effect of rainfall in the week immediately prior, but a negative effect in
282 the second and third prior weeks. Unlike *DW*, rainfall four weeks prior did not have an effect on
283 the model estimates. The strongest main effects variable was *DW* with a 1-week lag with an
284 effect of 1.10. For precipitation, the strongest variable was lower than average precipitation
285 during weeks 27-39 of the previous year, which led to higher *MIR* estimates with an effect of -
286 0.21.

287 **Discussion**

288 Since temperature and precipitation are largely interdependent events, interaction
289 terms more realistically represented the relationship between temperature and precipitation
290 and their combined effect on infection rate. This was an important improvement over the

291 legacy model and provided insight into how weather affected the mosquito infection rate. In
292 particular, the interaction terms revealed that though higher *DW* generally increased *MIR*,
293 higher *DW* two weeks prior in combination with higher than average precipitation in weeks 1,
294 2, and 3 prior each resulted in lower *MIR*. In other words, higher precipitation slightly reduced
295 the magnitude of temperature's effect on *MIR*, as seen by the effect of *DW* on *MIR* decreasing
296 from 2.41 to a magnitude of 2.16 when the average precipitation in preceding weeks is below
297 average by 1.85 cm (Figure 5A). In addition, with lower than average precipitation, temperature
298 became an even stronger predictor of *MIR* (Figure 5A) and with lower than average
299 temperatures, precipitation had minimal to no effect on *MIR* (Figure 5B). Refer to the
300 Supplementary Materials for figures with all interaction plots.

301 The legacy model (Ruiz et al 2010) captured the overall shape of the mosquito infection
302 curve when applied to a sub-region of the area for which it was developed, but significant
303 improvements were possible by developing a new model to account for local weather
304 conditions, by using the local *MIR*, by introducing additional terms, and by using more years of
305 data in the model. We found that the general linear regression approach used by the legacy
306 model, with *MIR* based on prior weather conditions, provided a reproducible methodology to
307 estimate *MIR* in a location and time period that was not part of the original model. The
308 assessment of the use of prior *MIR* in the new DuPage County model led us to conclude that a
309 model that is not dependent on *MIR* measured in previous weeks is both statistically sound and
310 operationally preferred. In situations where the *MIR* can be reliably measured across the entire
311 study region, the *MIR* dependent model may give good predictions most weeks, but with the
312 caution that a prediction immediately after a rapid change in weather may not capture the true

313 effect of weather and thus over-emphasize the effect of past *MIR*. The inclusion of significant
314 interaction terms between rainfall and temperature greatly improved the model's fit and
315 provided more detailed insight into the relationship between weather and mosquito infection
316 rate.

317 Both higher temperature and below average precipitation led to an increase in *MIR*,
318 which conforms to prior expectations (Shaman et al. 2005, Paz and Albersheim 2008, Paz et al.
319 2013). Additionally, temperature had a greater influence than precipitation on mosquito
320 infection as demonstrated in the results of all four models, where the effect of *DW* had a much
321 stronger effect than the precipitation variables (see Table 3). Significant interactions revealed
322 that when temperature was much lower than average, low precipitation had little to no effect
323 on the prediction and when precipitation was much lower than average, temperature had an
324 even greater influence. It is this second situation that is most likely to lead to illness from WNV,
325 and we recommend that public health personnel should develop the information they provide
326 to the public on the risk of WNV in the following week by incorporating both the predicted *MIR*
327 and the prior weather patterns.

328 Because lower than average precipitation during weeks 27-39 of the previous year
329 resulted in higher *MIR* during the current mosquito season, the next summer's *MIR* can be
330 approximated prior to the onset of the WNV season, a point also made by Hahn et al. (2015).
331 Reasons behind the significance of the previous fall and winter's precipitation remain unclear.
332 It is possible, for example, that less rainfall during the fall and winter are correlated with the
333 amount of rainfall during later periods, and the effect is indirect rather than direct. It is also
334 possible that this variable improves the model mostly during the early part of the season, and it

335 may not be as important for the critical period of virus amplification. Less moisture in the soil
336 at the start of the season might lead to a more patchy distribution of mosquito larval sites, thus
337 influencing spatial patterns of interactions between birds and mosquitoes. Vegetation
338 characteristics, related to the avian hosts and their interactions with WNV vectors, may also be
339 affected by weather (Gibbs et al. 2006, LaDeau et al. 2008). Mosquito abundance may be
340 higher following a dry fall due to a reduction of predator species (Walsh et al. 2008).
341 Abundance may also be affected by a mild winter with higher survival rates of overwintering
342 *Culex pipiens* and *restuans*, while cooler weather earlier in the fall may lead to earlier, more
343 successful hibernation, and earlier warmer conditions in the spring could provide conditions for
344 early emergence (Walsh et al. 2008). The simple linear models used in the current study would
345 not be suitable to determine these complex biological interactions. However, both
346 precipitation and temperature during the prior year and the winter and spring weeks leading up
347 to mosquito season of the same year should be evaluated in future work.

348 Several factors may influence the calculation of MIR estimates used to build the models.
349 For WNV surveillance, the best policy management decisions are often tempered by funding
350 and public perceptions related to pesticide use and to the risk of human illness (Shaw et al.
351 2010, Tedesco et al. 2010, Dickenson and Paskewitz 2012,). Thus, temporal and spatial
352 variability in testing effort and in mosquito abatement is likely, but it is difficult to measure.
353 Pooled samples for testing mosquitoes are another issue. The testing of mosquitoes is usually
354 done with pools of variable size, rather than testing individuals. This characteristic, in
355 combination with the relative inability to discriminate between latent and active infection
356 levels, and the differences in results from different testing methods can lead to errors in the

357 measurement of mosquito infection rates (Bustamante and Lord 2010, Speybroeck et al. 2012).
358 Although of interest from a research perspective, these measures are not easily managed
359 across administrative areas, and different approaches in other places may need to be
360 considered if this *MIR* model is applied in other locations.

361 An important area of research is to explore more fully the effects of weather on avian
362 hosts, mosquito abundance and human behavior relative to the risk of WNV illness. The
363 relationship between mosquito infection and the abundance of *Culex* vectors could not be
364 assessed in our analysis, so the model does not use a vector index measure, which is often used
365 to determine the risk of human exposure (e.g. Chung et al. 2013). Abundance measures were
366 not available in this study because the number of tested mosquitoes, not the full count from
367 each collection was recorded in the IDPH database. DuPage County did have some light trap
368 and larval sampling to monitor vector mosquito abundance, but these were not collected
369 systematically across all entities and could not be incorporated into the model.

370 One future analysis would be to determine how weather influences the abundance of
371 vector mosquito populations both temporally and spatially (see Yoo 2014 for example), and
372 develop an approach to incorporate this into predictions of MIR. For example, Lebl et al. (2013)
373 analyzed light trap counts of *Culex* mosquitoes relative to weather in northeast Illinois and
374 found abundance was positively correlated with temperature during the prior two weeks and
375 negatively associated with increased wind speed. Chaves et al. (2013) found that *Culex pipiens*
376 abundance in the island of Jeju-do Korea was positively associated with temperature, but with
377 heterogeneities at local scales, as mosquito abundance decreased with rainfall in the north,
378 while it increased with minimum temperature in the south. Morin and Comrie (2013)

379 developed a climate-based approach to link temperature and rainfall conditions in the southern
380 U.S. to the population dynamics of the WNV vector *Culex quinquefasciatus* and extended their
381 approach to consider future conditions under climate change, finding that dry and hot
382 conditions may reduce populations. Kunkel *et al.* (2006) used a long-term database on vector
383 mosquito abundance in Central Illinois to link weather to the so-called “crossover” of the early-
384 season dominance of *Culex restuans* that gives way to the later-season *Culex pipiens*. The timing
385 of their crossover was related to weather and often coincided with WNV amplification
386 (Westcott *et al.* 2011). Studies that incorporate both biotic and abiotic factors to model
387 mosquito abundance are relatively rare, and future work should be directed in this area to
388 create a more nuanced WNV risk estimate.

389 The main intent of our work was to build a stable local model that would provide a
390 reliable way to predict MIR quickly and effectively. With our model, we were able to provide
391 regionally calibrated model-based estimates of MIR two to three weeks sooner than MIR
392 estimation that needed test results from mosquitoes collected by a variety of agencies to be
393 completed by all groups and compiled into a common MIR value. Of immediate interest would
394 be to apply our methods to other locations to develop a similar weather-only model for further
395 comparison where vegetation and landscape factors are different from those in northern
396 Illinois. We do not expect that our model will apply to all other locations, but we expect that its
397 general structure can form a template for similar MIR prediction models elsewhere and
398 ultimately may be a way to estimate MIR, even in the absence of lab testing for WNV.

399

400 Acknowledgments

401 This research was made possible through funding from the Wheaton Mosquito Abatement

402 District, the Forest Preserve District of DuPage County, and the DuPage County Health

403 Department. A. Bolds encouraged the initiation of the operational model and facilitated

404 communication with the mosquito abatement and public health agencies in DuPage County.

405 This material is also based upon work supported by the National Science Foundation Ecology of

406 Infectious Diseases program under Awards 0429124 and 0840403.

407

408

409

References Cited

- 410 **Andreadis, T. G. 2012.** The contribution of *Culex pipiens* complex mosquitoes to transmission and
411 persistence of West Nile virus in North America. J Am Mosquito Contr 28: 137-151.
412
- 413 **Baker, D., B. Sharratt, H. Chiang, J. Zandlo, and D. Ruschy. 1984.** Base temperature selection for the
414 prediction of European corn borer instars by the growing degree day method. Agr Forest
415 Meteorol 32: 55-60.
416
- 417 **Barker, C. M., B. G. Bolling, W. C. Black, C. G. Moore, and L. Eisen. 2009.** Mosquitoes and West Nile
418 virus along a river corridor from prairie to montane habitats in eastern Colorado. J Vector Ecol
419 34: 276-293.
420
- 421 **Becker, B., P. T. Leisnham, and S. L. LaDeau. 2014.** A tale of two city blocks: differences in immature
422 and adult mosquito abundances between socioeconomically different urban blocks in Baltimore
423 (Maryland, USA). Int J Environ Res Publ Health 11: 3256-3270.
424
- 425 **Bertolotti, L., U. D. Kitron, E. D. Walker, M. O. Ruiz, J. D. Brawn, S. R. Loss, G. L. Hamer, and T. L.
426 Goldberg. 2008.** Fine-scale genetic variation and evolution of West Nile Virus in a transmission
427 "hot spot" in suburban Chicago, USA. Virology 374: 381-389.
428
- 429 **Biggerstaff, B. J. 2009.** PooledInfRate, Version 4.0: A Microsoft Office Excel Addin to Compute
430 Prevalence Estimates from Pooled Samples computer program, version By Biggerstaff, B. J., Fort
431 Collins, CO.
432
- 433 **Bustamante, D. M., and C. C. Lord. 2010.** Sources of error in the estimation of mosquito infection rates
434 used to assess risk of arbovirus transmission. Am J Trop Med Hyg 82: 1172-1184.
435
- 436 **CDC. 2014.** West Nile Virus Statistics and Maps, Division of Vector-Borne Diseases,
437 <http://www.cdc.gov/westnile/statsMaps/>.
438
- 439 **Chaves, L. F., Y. Higa, S. H. Lee, J. Y. Jeong, S. T. Heo, M. Kim, N. Minakawa, and K. H. Lee. 2013.**
440 Environmental forcing shapes regional house mosquito synchrony in a warming temperate
441 island. Environ Entomol 42: 605-613.
442
- 443 **Chaves, L. F., and U. D. Kitron. 2011.** Weather variability impacts on oviposition dynamics of the
444 southern house mosquito at intermediate time scales. B Entomol Res 101: 633-641.
445
- 446 **Chaves, L. F., and M. Pascual. 2007.** Comparing models for early warning systems of neglected tropical
447 diseases. PLoS Neglect Trop D 1: e33.
448
- 449 **Chung, W. M., C. M. Buseman, S. N. Joyner, S. M. Hughes, T. B. Fomby, J. P. Luby, and R. W. Haley.
450 2013.** The 2012 West Nile encephalitis epidemic in Dallas, Texas. JAMA – J Am Med Assoc 310:
451 297-307.

- 452
453 **Ciota, A. T., and L. D. Kramer. 2013.** Vector-virus interactions and transmission dynamics of West Nile
454 virus. *Viruses* 5: 3021-3047.
455
- 456 **Curriero, F. C., S. M. Shone, and G. E. Glass. 2005.** Cross correlation maps: a tool for visualizing and
457 modeling time lagged associations. *Vector-Borne Zoonot* 5: 267-275.
458
- 459 **Day, J. F., and G. A. Curtis. 1999.** Blood feeding and oviposition by *Culex nigripalpus* (Diptera:
460 Culicidae) before, during, and after a widespread St. Louis encephalitis virus epidemic in Florida.
461 *J Med Entomol* 36: 176-181.
462
- 463 **Dickinson, K., and S. Paskewitz. 2012.** Willingness to pay for mosquito control: how important is West
464 Nile virus risk compared to the nuisance of mosquitoes? *Vector-Borne Zoonot* 12: 886-892.
465
- 466 **Dohm, D. J., M. L. O'Guinn, and M. J. Turell. 2002.** Effect of environmental temperature on the ability
467 of *Culex pipiens* (Diptera: Culicidae) to transmit West Nile virus. *J Med Entomol* 39: 221-225.
468
- 469 **Ebert, T. A., R. Brilansky, and M. Rogers. 2010.** Reexamining the pooled sampling approach for
470 estimating prevalence of infected insect vectors. *Ann Entomol Soc Am* 103: 827-837.
471
- 472 **Gardner, A. M., T. K. Anderson, G. L. Hamer, D. E. Johnson, K. E. Varela, E. D. Walker, and M. O. Ruiz.**
473 **2013.** Terrestrial vegetation and aquatic chemistry influence larval mosquito abundance in
474 catch basins, Chicago, USA. *Parasite Vector* 6:9.
475
- 476 **Gardner, A. M., G. L. Hamer, A. M. Hines, C. M. Newman, E. D. Walker, and M. O. Ruiz. 2012.**
477 Weather variability affects abundance of larval *Culex* (Diptera: Culicidae) in storm water catch
478 basins in suburban Chicago. *J Med Entomol* 49: 270-276.
479
- 480 **Gibbs, S. E., M. C. Wimberly, M. Madden, J. Masour, M. J. Yabsley, and D. E. Stallknecht. 2006.**
481 Factors affecting the geographic distribution of West Nile virus in Georgia, USA: 2002-2004.
482 *Vector-Borne Zoonot* 6: 73-82.
483
- 484 **Gu, W., R. Lampman, and R. J. Novak. 2003.** Problems in estimating mosquito infection rates using
485 minimum infection rate. *J Med Entomol* 40: 595-596.
486
- 487 **Hahn, M. B., A. J. Monaghan, M. H. Hayden, R. J. Eisen, M. J. Delorey, N. P. Lindsey, R. S. Nasci, and**
488 **M. Fischer. 2015.** Meteorological conditions associated with increased incidence of West Nile
489 virus disease in the United States, 2004-2012. *Am J Trop Med Hyg* 92: 1013-1022.
490
- 491 **Hamer, G. L., U. D. Kitron, J. D. Brawn, S. R. Loss, M. O. Ruiz, T. L. Goldberg, and E. D. Walker. 2008.**
492 *Culex pipiens* (Diptera: Culicidae): a bridge vector of West Nile virus to humans. *J Med Entomol*
493 45: 125-128.
494
- 495 **Hartley, D. M., C. M. Barker, A. Le Menach, T. Niu, H. D. Gaff, and W. K. Reisen. 2012.** Effects of

- 496 temperature on emergence and seasonality of West Nile virus in California. Am J Trop Med Hyg
497 86: 884-894.
- 498
- 499 **Hepworth, G. 2005.** Confidence intervals for proportions estimated by group testing with groups of
500 unequal size. J Agr Biol Envir St 10: 478-497.
- 501
- 502 **Hokit, G., S. Alvey, J. M. Geiger, G. D. Johnson, M. G. Rolston, D. T. Kinsey, and N. Tall Bear. 2013.**
503 Using undergraduate researchers to build vector and West Nile virus surveillance capacity. Int J
504 Environ Res Publ Health 10: 3192-3202.
- 505
- 506 **Jones, C. E., L. P. Lounibos, P. P. Marra, and A. M. Kilpatrick. 2012.** Rainfall influences survival of *Culex*
507 *pipiens* (Diptera: Culicidae) in a residential neighborhood in the mid-Atlantic United States. J
508 Med Entomol 49: 467-473.
- 509
- 510 **LaDeau, S. L., P. P. Marra, A. M. Kilpatrick, and C. A. Calder. 2008.** West Nile virus revisited:
511 Consequences for North American ecology. Bioscience 58: 937-946.
- 512
- 513 **Lebl, K., K. Brugger, and F. Rubel. 2013.** Predicting *Culex pipiens/restuans* population dynamics by
514 interval lagged weather data. Parasite Vector 6: 129.
- 515
- 516 **Macdonald, G. 1956.** Theory of the eradication of malaria. B World Health Organ 15: 369-387.
- 517
- 518 **Moore, C. G. M., R.G.; Mitchell, C.J.; Nasci, R.S.; Tsai, T.F.; Calisher, C.H.; Marfin, A.A; Moore, P.S.;**
519 **Gubler, D.J. 1993.** Guidelines for Arbovirus Surveillance Programs in the United States. Division
520 of Vector-Borne Infectious Diseases, Centers for Disease Control and Prevention, Fort Collins,
521 Colorado.
- 522
- 523 **Morin, C. W., and A. C. Comrie. 2013.** Regional and seasonal response of a West Nile virus vector to
524 climate change. B Natl Acad Sci USA 110: 15620-15625.
- 525
- 526 **Nasci, R. S. 2013.** Monitoring and controlling West Nile Virus: are your prevention practices in place? J
527 Environ Health 75: 42-44.
- 528
- 529 **Paz, S., and I. Albersheim. 2008.** Influence of warming tendency on *Culex pipiens* population
530 abundance and on the probability of West Nile fever outbreaks (Israeli Case Study: 2001-2005).
531 Ecohealth. 5: 40-48.
- 532
- 533 **Paz, S., D. Malkinson, M. S. Green, G. Tsioni, A. Papa, K. Danis, A. Sirbu, C. Ceianu, K. Katalin, E.**
534 **Ferenczi, H. Zeller, and J. C. Semenza. 2013.** Permissive summer temperatures of the 2010
535 European West Nile fever upsurge. PloS One 8: e56398.
- 536
- 537 **R Development Core Team 2013.** R: A language and environment for statistical computing. computer
538 program, version By R Development Core Team, Vienna, Austria <http://www.R-project.org/>.
- 539

- 540 **Reisen, W. K., Y. Fang, and A. C. Brault. 2008.** Limited interdecadal variation in mosquito (Diptera:
541 Culicidae) and avian host competence for Western equine encephalomyelitis virus (Togaviridae:
542 Alphavirus). *Am J Trop Med Hyg* 78: 681-686.
543
- 544 **Reisen, W. K., T. Thiemann, C. M. Barker, H. Lu, B. Carroll, Y. Fang, and H. D. Lothrop. 2010.** Effects of
545 warm winter temperature on the abundance and gonotrophic activity of *Culex* (Diptera:
546 Culicidae) in California. *J Med Entomol* 47: 230-237.
547
- 548 **Ruiz, M. O., L. F. Chaves, G. L. Hamer, T. Sun, W. M. Brown, E. D. Walker, L. Haramis, T. L. Goldberg,
549 and U. D. Kitron. 2010.** Local impact of temperature and precipitation on West Nile virus
550 infection in *Culex* species mosquitoes in northeast Illinois, USA. *Parasite Vector* 3: 19.
551
- 552 **Shaman, J., and J. F. Day. 2007.** Reproductive phase locking of mosquito populations in response to
553 rainfall frequency. *PloS One* 2: e331.
554
- 555 **Shaman, J., J. F. Day, and M. Stieglitz. 2005.** Drought-induced amplification and epidemic transmission
556 of West Nile virus in southern Florida. *J Med Entomol* 42: 134-141.
557
- 558 **Shaw, I. G. R., P. F. Robbins, and J. P. Jones. 2010.** A Bug's life and the spatial ontologies of mosquito
559 management. *Ann Assoc Am Geogr* 100: 373-392.
560
- 561 **Speybroeck, N., C. J. Williams, K. B. Lafia, B. Devleeschauwer, and D. Berkvens. 2012.** Estimating the
562 prevalence of infections in vector populations using pools of samples. *Med Vet Entomol* 26:
563 361-371.
564
- 565 **Tedesco, C., M. Ruiz, and S. McLafferty. 2010.** Mosquito politics: Local vector control policies and the
566 spread of West Nile virus in the Chicago region. *Health Place* 16: 1188-1195.
567
- 568 **Turell, M. J., D. J. Dohm, M. R. Sardelis, M. L. Oguinn, T. G. Andreadis, and J. A. Blow. 2005.** An
569 update on the potential of north American mosquitoes (Diptera: Culicidae) to transmit West
570 Nile Virus. *J Med Entomol* 42: 57-62.
571
- 572 **Turell, M. J., M. L. O'Guinn, D. J. Dohm, and J. W. Jones. 2001.** Vector competence of North American
573 mosquitoes (Diptera: Culicidae) for West Nile virus. *J Med Entomol* 38: 130-134.
574
- 575 **Walsh, A. S., G. E. Glass, C. R. Lesser, and F. C. Curriero. 2008.** Predicting seasonal abundance of
576 mosquitoes based on off-season meteorological conditions. *Environ Ecol Stat* 15: 279-291.
577
- 578 **Walter, S. D., S. W. Hildreth, and B. J. Beaty. 1980.** Estimation of infection rates in population of
579 organisms using pools of variable size. *Am J Epidemiol* 112: 124-128.
580
- 581 **Westcott, N. E., S. D. Hilberg, R. L. Lampman, B. W. Alto, A. Bedel, E. J. Muturi, H. Glahn, M. Baker, K.
582 E. Kunkel, and R. J. Novak. 2011.** Predicting the seasonal shift in mosquito populations
583 preceding the onset of the West Nile virus in Central Illinois. *B Am Meteorol Soc* 92: 1173-1180.

584

585 **Yoo, E. H. 2014.** Site-specific prediction of West Nile virus mosquito abundance in Greater
586 Toronto area using generalized linear mixed models. *Int J Geogr Inf Sci* 28: 296-313.

587

588

589 **Tables**590 **Table 1.** DuPage County West Nile virus-related annual conditions data summary.

Year	MIR Mean (SD)	Week of	Avg Prec. (cm) of 3 weeks before peak week*	DW at peak week*	WNV Human Cases
		Max MIR (Peak)			
2005	5.57 (6.14)	32	-0.74	11.95	47
2006	6.88 (8.50)	34	-0.68	6.67	43
2007	2.76 (3.18)	33	1.23	-1.52	10
2008	1.13 (1.77)	37	0.59	-2.31	1
2009	0.78 (0.77)	37	0.36	-5.22	0
2010	5.66 (6.90)	35	-1.61	16.98	17
2011	2.63 (3.73)	36	-0.77	12.46	2
2012	8.74 (7.91)	32	-0.08	29.02	56
2013	4.52 (5.56)	36	-1.40	5.42	6
2014	3.27 (5.07)	35	3.42	-0.57	6

591 *Differences from weekly averages using the 30-year Normal of both temperature and
592 precipitation.

593

594

595

596

597

598 **Table 2.** Pearson's correlation (r) between *MIR* and the previous year's average precipitation
 599 over blocks of 52, 26, and 13 weeks.

600

601

	Weeks	Weeks	Weeks	Weeks	Weeks	Weeks	Weeks
	1-52	1-26	27-52	1-13	14-26	27-39	40-52
r	-0.262	0.014	-0.439	-0.063	0.048	-0.460	-0.091
p-value	< 0.001	0.853	< 0.001	0.391	0.514	< 0.001	0.213

602

603

604

605

606 **Table 3.** Variable effects (standard errors) and fit of 4 model types: (1) Main effect model
 607 dependent on MIR, (2) Main effect model independent of MIR, (3) Interaction model
 608 dependent on MIR, (4) Interaction model independent of MIR.

	Model 1	Model 2	Model 3	Model 4
R² adjusted	0.661	0.511	0.721	0.658
PRESS	0.635	0.481	0.635	0.537
AIC	340.6	409.0	315.2	353.5
Main Effects				
(at week lag)				
<i>MIR (2nd order)</i>	0.14(0.02)***		0.11(0.02)***	
<i>Week</i>	-0.08(0.05)	-0.25(0.06)***	-0.10(0.05)*	-0.23 (0.05)***
	Model 1	Model 2	Model 3	Model 4
<i>Prec. (1)</i>			0.06(0.04)	0.09(0.05)*
<i>Prec. (2)</i>	-0.06(0.04)	-0.09(0.05)	-0.05(0.04)	-0.03(0.05)
<i>Prec. (3)</i>			-0.07(0.04)	-0.09(0.05)
<i>DW (1)</i>	0.72(0.12)***	0.99(0.14)***	1.12(0.23)***	1.21(0.25)***
<i>DW (2)</i>			0.01(0.32)	0.13(0.36)
<i>DW (3)</i>			-0.59(0.32)	-0.83(0.37)*
<i>DW (4)</i>	-0.53(0.11)***	-0.41(0.13)**	0.01(0.21)	0.42(0.26)
<i>Previous Year Prec.</i> <i>(wks 27-39)</i>	-0.14 (0.05)***	-0.30(0.05)***	-0.08(0.04)	-0.18(0.05)***

Interaction Terms

<i>Prec. (1)*DW (2)</i>	-0.83(0.20)***	-1.15(0.22)***
<i>Prec. (1)*DW (3)</i>	0.86(0.20)***	1.13(0.22)***
<i>Prec. (2)*DW (2)</i>	0.89(0.37)*	1.29(0.41)**
<i>Prec. (2)*DW (3)</i>	-1.71(0.51)**	-2.53(0.57)***
<i>Prec. (2)*DW (4)</i>	0.78(0.22)***	1.18(0.24)***
<i>Prec. (3)*DW(1)</i>	0.72(0.26)**	0.83(0.29)**
<i>Prec. (3)*DW(2)</i>	-0.68(0.25)**	-0.85(0.28)**
<i>DW(2)*DW(4)</i>	-0.12(0.03)***	-0.71(0.17)***
<i>DW(2)*DW(3)</i>		0.55(0.18)**

609

610 R^2 adjusted and predicted residual sum of squared errors (PRESS) are reported for each model.
 611 * indicated the variable is significant at 5% level, ** significant at 1% level and *** significant at
 612 0.1% level.

613

614 **Figure Legends**

615 **Figure 1.** Map of the study region with the two weather station locations and the average
616 number of trap locations at which mosquitoes were tested. The legacy model was developed
617 from data combined from Cook and DuPage counties. The current objectives focus on DuPage
618 County, only. The average number of traps is for the years from 2005 to 2014 summarized for
619 hexagons of 200 hectares.

620 **Figure 2** (A) Average weekly Mosquito Infection Rate (*MIR*) with normal precipitation and (B)
621 Average *MIR* with normal temperature. The average *MIR* is a weekly average from the DuPage
622 County study area from 2005 to 2014.

623 **Figure 3.** (A) Measured *MIR* and legacy model estimates (Predicted $MIR = a + 0.35$ (3wk *Prec.*
624 moving average at 3 week lag) + 0.42 (*DW* at 1week lag) – 1.57 (*previous year prec.*) (MSPE*
625 2.640). (B) Measured *MIR* and new model estimates with interactions (MSPE 1.826). *Mean
626 squared prediction error. Supplementary Material includes a graph of the full range of years
627 shown as a subset of four years in Figure 3A.

628 **Figure 4.** Correlations between weather variables and DuPage *MIR* at lags of from 1 to 8 weeks.

629 **Figure 5.** Interaction Plots between the variables *DW* and *precipitation* of preceding weeks and
630 the variable *MIR*. All variables are measured as the difference from the weekly average.



631 Covariates were scaled before plotting. A: Effect of *DW* when precipitation is low/high.



632 Equation of solid line: $MIR = -0.003 + 2.41 * DW$. B: Effect of precipitation when *DW* is low/high.


633 Equation of solid line: $MIR = -0.003 - 0.39 * prec.$


634











DuPage Local Model Details.


Included in this file is: details of coefficients and p-values of model terms, diagnostic plots for this model, scatterplots and interaction plots and interpretations for most significant terms.

Supplementary Table 1. Local DuPage Model Effects, Coefficients, and p-values for each term. *Significant at 0.05


Significant at 0.01 *Significant at 0.001

Variable	Effect	Coefficient	Significance (P-Value)
Intercept	0.164	-0.79	
Prec.1wk lag1	0.10	2.19	0.053
Prec.1wk lag2	-0.02	0.04	0.744
Prec.1wk lag3	-0.08	-1.44	0.081
DW lag1	1.12***	0.59	6×10^{-5} ***
DW lag2	0.13	0.11	0.62
DW lag3	-0.83*	-0.62	0.02*
DW lag4	0.40	0.44	0.05*
Prec.1yr Q4	-0.19***	-8.80	0.0004***
Prec.1wk lag1*DW lag 2	-1.26***	-2.86	1.34×10^{-7} ***
Prec.1wk lag1*DW lag 3	1.26***	3.06	1.47×10^{-7} ***
Prec.1wk lag2*DW lag 2	1.28**	2.79	0.004**
Prec.1wk lag2*DW lag 3	-2.65***	-6.24	2×10^{-5} ***
Prec.1wk lag2*DW lag 4	1.33***	3.40	3.71×10^{-7} ***
Prec.1wk lag3*DW lag 1	0.71*	1.47	0.021*
Prec.1wk lag3*DW lag 2	-0.74*	-1.63	0.013*
DW lag2*DW lag4	-0.74***	-0.07	1.25×10^{-4} ***
DW lag2*DW lag3	-0.73**	0.05	0.003**

Supplementary Figure 1. Legacy Model from Ruiz 2010, DuPage Local model, and actual DuPage MIR from 2005 through 2014




Supplementary Figure 2. Diagnostic Plots for Local Model above.




- Residuals vs Fitted Plot (Plot 1) shows residuals randomly distributed about 0
- Q-Q plot (Plot 3) validates the assumption that our residuals follow a normal distribution
- Leverage Plot (Plot 4) shows no large leverage points based on Cook's distance

Supplementary Figure 3. Scatter and Interaction Plots for most significant variables.










- Higher than average DW in past weeks ago leads to higher MIR this week (effect of DW decreases with lag as seen previously by correlation r)
- The lower from average last week's precipitation is, the less positive effect DW has on MIR



- In general, when there is lower than average precipitation in the past few weeks, DW in the past few weeks has a greater positive effect on MIR
- This observation seems to be most clear when looking at interaction plots of precipitation weeks and DW weeks that are closer together temporally.
- If DW is lower than average 3-4 weeks ago, DW 2 weeks ago has a more positive effect on MIR

<https://mc.manuscriptcentral.com/medent>



- In general, lower than average DW values for the past 4 weeks lessens the negative effect of precipitation in the past 3 weeks
- Again, this relationship is more clear between precipitation and DW of consecutive weeks

- Exception: When higher than average DW 2 weeks ago, higher than average precipitation last week has slightly positive effect. This reflects DW has stronger influence on MIR prediction than precipitation
- Precipitation in the past 3 weeks tends to have a negative effect on MIR