

FRACTAL DIMENSIONS FOR RADIOISOTOPE POLLUTION PATTERNS BY NUCLEAR POWER PLANT ACCIDENTS

Keisuke Saito ^{a*}, and Susumu Ogawa ^b

^a Nagasaki University, Graduate school of engineering, Bunkyo-machi, Nagasaki, 852-8521 Japan
jj20140051@cc.nagasaki-u.ac.jp

^b Nagasaki University, Graduate school of engineering, Bunkyo-machi, Nagasaki, 852-8521 Japan
ogawasusumu@nagasaki-u.ac.jp

KEY WORDS: Chernobyl, deposit, fractal, Fukushima, NOAA/AVRR

ABSTRACT:

The radioisotope pollution shows two types of patterns: dry and wet deposits for nuclear power plant accidents. Two surface pollution patterns were analysed by fractal. In Fukushima nuclear power plant accident, surface pollution by wet deposits was estimated to occur. However, actually it was no rain and white crystals were observed on the surface. Then, fractal analysis was carried out for the spatial distribution patterns of radio isotopes on the surface to judge the types of deposits. As a reference, Chernobyl nuclear power plant accident was checked for the spatial distribution patterns of radioisotopes on the surface. The objective patterns by fractal analysis were the surface pollution maps in Fukushima and Chernobyl, Abukuma river watershed map, and NOAA/AVHRR. The calculation of fractal dimensions was carried out with the box counting for binarized images. Fractal analysis results suggested the next conclusions. The radioisotope pollution in Fukushima might occur in both dry and wet deposits. The dry deposit might make the pollution pattern similar to the watershed, while the wet deposit might make the pollution pattern similar to cloud images. Moreover, most radioisotope contaminants might flow on the road in the forest valley and deposit on forest with and without rainfall in Fukushima.

1. INTRODUCTION

Each process was calculated from meteorological data for radioisotope pollutions (Japan Meteorological Agency, 2014) with hydrogen explosions at Fukushima Daiichi nuclear power plant derived from the earthquake and Tsunami on March 11, 2011. The pollution at Iidate village was estimated mainly from wet deposits by hydrogen explosion at the second reactor on March 15 (Imanaka, 2014). However, it was little rainfall at the period and dry deposits might be the main pollution by the leak including air vent. Then, the radioisotope pollution was examined on March 12 to 31. Moreover, fractal analysis was applied to the contribution rate of dry and wet deposits in Fukushima. Fractal dimension is the value corresponding to complexity for the forms.

1.1 Outline of the Fukushima nuclear power plant accident

At 14:46 on March 11, 2011, the great earthquake with magnitude of 9.0 occurred from a focus of Sanriku-oki. At 15:50, the great tsunami attacked Fukushima Daiichi and Daini Nuclear Power Plants. The emergent core cooling system did not work and lost cooling ability. Fukushima Daiichi had meltdown at Reactors 1 to 3, and a hydrogen explosion occurred, which made building structures flying to pieces and a big amount of isotopes leaking into the atmosphere. At 13:10 on March 20, Reactor 4 was fired. On March 21, it rained in the Kanto plain, and high concentration of iodine 131 was detected from drinking water in waterworks in the Tone river watershed on March 22 and 23.

1.2 Location

Fukushima nuclear power plant is located in the place 280km away from Tokyo. On the east side it faces the Pacific Ocean, from which tsunami flowed in. On the other hand, on the west side it faces Abukuma highlands, where the flat land near the plant continues to steep slopes on the surroundings. Location of Fukushima nuclear power plant is shown in Figure 1.



Figure 1. Location

*Corresponding author. Tel. +81 958192611.

E-mail address: jj20140051@cc.nagasaki-u.ac.jp

2. METHODS

In radioisotope pollution analysis, a particle model, the atmospheric diffusion model, and fractal analysis were selected.

2.1 Particle model

In a particle model, the particle was assumed to be sand with specific weight 2.65 for concrete debris generated by hydrogen explosions in the nuclear power plants and to diffuse by wind with radioisotopes and to fall down with Stokes equation.

$$v_s = \frac{D^2(\rho_p - \rho_a)}{18\mu} \quad (1)$$

where v_s = falling velocity
 D = particle size
 ρ_p = particle density, 2650kg/m³
 ρ_a = air density, 1.225kg/ m³
 μ = viscosity, 1.8×10⁻⁵Pas.

The pollution particle trajectory was obtained each height from the meteorological archives (Meteorological data archive) for the explosions polluting to East Fukushima on March 12 and 15.

2.2 Atmospheric diffusion model

All pollutions were calculated with the plume model and pollution Mar 15 was wet deposit.

Plume model: wind speed >1m/s

$$C_0 = \frac{q}{\sqrt{2\pi}\sigma_y U h} \exp\left(-\frac{y^2}{2\sigma_y^2}\right) \quad (2)$$

where C_0 = uniform concentration
 q = emittance
 U = wind speed
 h = vertical mixing layer height
 y = lateral distance
 σ_y = lateral diffusion distance width in the Axis y direction as next.

$$\sigma_y = \tan \sigma_{wd} \times x + \sigma_{yp} + \sigma_{y0} \quad (3)$$

where σ_{wd} = standard deviation of wind fluctuation (4°)
 σ_{yp} = diffusion width from Pasquill diagram
 σ_{y0} = initial diffusion width and half of the grid size (0.5km).

Horizontal diffusion width in Pasquill is next.

$$\sigma_y = 0.1467x^{0.889} \text{ (Stability D)} \quad (4)$$

$$\sigma_y = 0.396x^{0.865} \text{ (Stability B)} \quad (5)$$

Dry deposit speed:

$$k_{dry} = \frac{V_d}{h} \quad (6)$$

where k_{dry} = dry deposit constant (s⁻¹)
 V_d = dry deposit rate (m/s)
 h = mixing layer height (m).

Here, stability D, $h = 200$ m, $V_d = 0.002$ m/s, thus, $k_{dry} = 0.00001$ s⁻¹.

Wet deposit speed:

$$k_{wet} = \frac{LR}{h} \quad (7)$$

where k_{wet} = wet deposit constant (s⁻¹)
 L = precipitation runoff coefficient
 R = precipitation (m/s).

Here, $L = 1.5$, $R = 0.0015$ m/s, $k_{wet} = 0.00001$ s⁻¹.

$V_d = 0.002$ m/s corresponds to particle size 0.005mm, 5mm sand particle with natural falling. When precipitation is less than 1.5mm/h, both dry and wet deposits showed in the same way.

2.3 Fractal analysis

Fractal is an irregular geometry with self-similar. Fractal is determined by fractal dimensions, which distribute 1 to 2 in two dimensional images. In fractal analysis, the objects were a soil pollution map in Fukushima accident (Ministry of Education, Culture, Sports, Science and Technology, 2011), a soil pollution map in Chernobyl accident (International Advisory Committee, 1991), Abukuma river watershed map, and NOAA/AVHRR data. In fractal dimension calculation, the box counting method was selected for binarized images.

$$D = \lim_{r \rightarrow 0} \frac{\log N_r}{\log(1/r)} = - \lim_{r \rightarrow 0} \frac{\log N_r}{\log r} \quad (8)$$

where D = fractal dimension
 r = the side length of a box
 N_r = number of boxes to cover the shape by box with side length r .

3. RESULTS

3.1 Particle model

The main hydrogen explosions occurred four times as shown in Table 1. But, only the fourth reactor showed the increase of spatial dose and the others did not show any increase of spatial dose. Moreover, the wind direction at the fourth reactor explosion was north.

Reactor	1	2	3	4
Date/Time	Mar12 15:36	Mar15 18:10	Mar14 11:01	Mar15 6:14
Wind Direction	SE	SE	W	N

Table 1. Hydrogen explosions and the wind directions

From these evidences, the possibility of pollution for Iitate village may be contributed by the first reactor explosion on March 12 and the second reactor explosion on March 15. With a time series of wind profiles (Meteorological data archive), a particle model showed the estimate of pollution on March 12 in Figure 2. Wind directions each height were quite different.




Figure 2. Particle flying traces estimated from wind directions and speeds on March 12.

From this result, the possibility of pollution should be contributed by radioisotopes flying at altitude 100m to 300m.

On the other hand, it rained at Iidate village at 16:00 on March 15 shown in Figure 3. At 18:10, the second reactor exploded. Raincloud passed over Fukushima from west and caught flying radioisotopes over East Fukushima at 21:00 as shown in Figure 4. By 24:00 most polluted particles fell down in wet deposit as shown in Figure 5. Polluted particles less than 800m in altitude were estimated to fall down in wet deposit. Polluted particles more than 1000m in altitude passed away to the Pacific Ocean.




Figure 3. Hyetograph on March 15 to 16 at Iidate village.




Figure 4. Flying polluted particles and rainclouds at 21:00 on March 15.




Figure 5. Flying polluted particles and rainclouds at 0:00 on March 16.

3.2 A time-series of spatial dose and precipitation

A time-series of spatial dose during March 12 and 31 and the contribution to pollution for Iidate village with wind directions east to south and precipitation are shown in Figures 4 and 5.

Figure 6 shows a time-series of spatial dose and the ratio of pollution for Iidate village with 19.6%. Figure 7 shows a time-series of spatial dose and the ratios of dry to wet deposits, respectively 6.5% for wet and 93.5% for dry deposits. Precipitation at Iidate village during the period is shown in Figure 8.

During this period, it rained a little with rain intensity 0.5 to 1.5 mm/h and pollution was dry deposits except March 15. It means that pollution at Iidate village was contributed by mainly dry deposits from the leak of radioisotopes including air vent.

3.3 Atmospheric diffusion model

As shown in Figure 9, most of pollution for Iidate village was contributed by the next three events.

1) Pollution on March 15

where the atmospheric stability was D, the mean wind direction was south southeast with 157.5°, the mean wind speed was 1.3 m/s during 12:00 to 22:30. It rained at 22:30 to 23:30. The mean spatial dose at the gate of Fukushima Daiichi nuclear power plant was 657 $\mu\text{Sv/h}$. It was only wet deposit during this period.

2) Pollution on March 18

where the atmospheric stability was B, the mean wind direction was southeast with 147°, the mean wind speed was 2.0 m/s during 12:00 to 17:30. The mean spatial dose at the gate of Fukushima Daiichi nuclear power plant was 3017 $\mu\text{Sv/h}$.

3) Pollution on March 20

where the atmospheric stability was B, the mean wind direction was southeast with 146°, the mean wind speed was 1.7 m/s during 12:00 to 17:30. The mean spatial dose at the gate of Fukushima Daiichi nuclear power plant was 2832 $\mu\text{Sv/h}$.




Figure 6. A time series of spatial dose (unit: $\mu\text{Sv/h}$) and the ratio of pollution for Iidate village.




Figure 7. A time series of spatial dose (unit: $\mu\text{Sv/h}$) and the ratio of dry and wet deposits.




Figure 8. Precipitation at Iidate village on March 11 to 31.




Figure 9. Main pollution routes for Iidate.

The overlapped atmospheric pollution map shows almost a normal distribution. From the altitudes in DEM (Figure 10), the deposit ratios were assumed to be 0.3 on the adverse slopes and 0.1 on the downward slopes, and Figures 11 and 12 were obtained.




Figure 10. DEM




Figure 11. Overlapped radioisotope pollutions on March 15, 18 and 20.





Figure 12. Actual pollution map by aircraft monitoring (upper) and synthesized radioisotope pollutions with altitude correction (lower). (Unit: $\mu\text{Sv/h}$)

3.4 Fractal analysis

Fractal dimensions were quite different between a soil pollution map in Fukushima nuclear power plant accident and Chernobyl accident, respectively 1.83 and 1.53. The fractal dimension for Abukuma river watershed was 1.83, while the one of NOAA/AVHRR with a cloud image was 1.56. It means that the fractal dimension of soil pollution in Fukushima nuclear power plant accident corresponded to the one of Abukuma river watershed.

On the other hand, the fractal dimension of soil pollution in Chernobyl nuclear power plant accident corresponded to the one of NOAA/AVHRR with a cloud image, as shown in Figure 13.




Figure 13. NOAA/AVHRR image

Objects	Fractal dimension
Soil pollution map in Fukushima	1.83
Soil pollution map in Chernobyl	1.53
Abukuma river watershed	1.83
NOAA/AVHRR with cloud image	1.56

Table 2 Fractal dimensions of radioisotope pollutions

4. DISCUSSION

4.1 Radioisotope pollution at Iitate village

The pollution at Iitate village stated from hydrogen explosion on March 12, but most of contribution was three pollutions on March 15, 18, and 20. Pollution on March 12 was brought by radioisotopes about 100m in altitude. On March 15, 18, and 20, radioisotopes at less than 200m, 500m, and 500m in altitude respectively flowed in the village, but on March 15, it rained directly after sedimentation and others without rainfall. The pollution distributions were determined by mainly the landforms. Especially, radioisotopes collided against the slopes of mountains with altitude 500m to 1192m and adsorbed the surface with pollution.

4.2 Dry deposit and wet deposit

On March 12 to 31, precipitation was 24.5mm in total and ranged 0.5 to 1.5mm/h. On the other hand, in Chernobyl, it rained with more than 5mm at most on April 29 in radioisotope emission during 10 day period, as shown in Figure 14.




Figure 14. Precipitation distribution in Chernobyl on April 29, 1986. (International Advisory Committee,1991)




Figure 15. Radioisotope pollution patterns in Chernobyl. (International Advisory Committee,1991)

As a result, the wet deposit brought a pattern of radioisotope pollution as shown in Figure 15. The change of the altitude is small there and the deposit developed without the effect of the landforms. Therefore, in Fukushima dry deposit was dominant with the effect of the landforms, while in Chernobyl wet deposit was dominant with the effect of precipitation.

5. CONCLUSIONS

From the above results, the next items were concluded.

- (1) The radioisotope pollution for Iitate village from Fukushima Daiichi power plant was emitted by the leak mainly on March 15, 18 and 20.
- (2) The radioisotope pollution from Fukushima Daiichi nuclear power plant was limited because the diffusion of radioisotopes was controlled by mountain in altitude 500m to 1192m.
- (3) The deterministic difference between Fukushima Daiichi nuclear power plant and Chernobyl nuclear power plant in radioisotope pollution was that the former was affected by landform more than precipitation while the latter was affected by precipitation more than the landform.

In this paper, the deposit ratio was obtained from the altitude in DEM, but the next elements are required to be estimated considering the movement of radioisotopes after the deposit.

- (1) Runoff from rivers and groundwater. (Figure 16)
- (2) Absorption and runoff by land covers. (Figure 17)
- (3) Transportation by public networks. (Figure 18)





Figure 16. Runoff from rivers and groundwater



- カテゴリー
- 1 水域 Water
 - 2 都市 Urban
 - 3 水田 Paddy
 - 4 畑地 Crop
 - 5 草地 Grass
 - 6 落葉樹 Deciduous forest
 - 8 常緑樹 Evergreen forest
 - 10 裸地、砂地 Bare surface
 - 11 雪氷 Snow and ice
 - その他 other

Figure 17. Land cover




Figure 18. Transportation by public network

From these elements, emission by explosions and leaks, advective diffusion in the atmosphere, and movement on the ground surface should be analysed as the comprehensive pollution mechanism near future.

REFERENCES

- Imanaka, T., 2014. Estimate of initial external exposure in Iitate village, *Science*, 84, 3, pp. 322-332.
- IAEA, 1991. International Advisory Committee, The international Chernobyl Project, Technical report, Vienna,
- Japan Meteorological Agency, 2014. Various data and materials, 2014.
- National Institute of Informatics, 2011. Archives of 2011 Great East Japan Earthquake, <http://agora.ex.nii.ac.jp/earthquake/-201103-eastjapan/weather/data/wind-20110311/surf/>
- Mandelbrot, B.B., 1982. *The Fractal Geometry of Nature*, W.H.Freeman and Company, New York.
- Ministry of Education, Culture, Sports, Science and Technology, 2011. Aircraft monitoring.
- Takayasu, H., 1986. *Fractal*, Asakura Publishing Co., Ltd.