

1 Algicidal hydroxylated C18 unsaturated fatty acids from the red alga *Tricleocarpa jejuensis*:  
2 Identification, synthesis and biological activity

3

4 Shijiao Zha<sup>1</sup>, Kazuyoshi Kuwano<sup>1</sup>, Tomohiro Shibahara<sup>2</sup>, Fumito Ishibashi<sup>1\*</sup>

5

6 <sup>1</sup>Graduate School of Fisheries and Environmental Sciences and <sup>2</sup>Faculty of Fisheries, Nagasaki  
7 University, 1-14 Bunkyo-Machi, Nagasaki, 852-8521, Japan.

8

9 \*Corresponding author

10 fumito@nagasaki-u.ac.jp

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37 Abstract

38 Bioassay-guided separation of a methanol extract of *Tricleocarpa jejuensis* by monitoring  
39 algicidal activity against the red tide phytoplankton *Chattonella antiqua* led to the isolation of an  
40 active fraction consisting of a mixture of four isomeric compounds. The active compounds were  
41 identified as (*E*)-9-hydroxyoctadec-10-enoic acid (**1**), (*E*)-10-hydroxyoctadec-8-enoic acid (**2**), (*E*)-  
42 11-hydroxyoctadec-12-enoic acid (**3**) and (*E*)-12-hydroxyoctadec-10-enoic acid (**4**) by NMR, IR and  
43 mass spectral data. The structures were confirmed by comparison of the NMR and MS data with  
44 those of authentic samples of **1**~**4** obtained by unambiguous syntheses. Synthesized hydroxy acids  
45 **1**~**4** and related compounds were assessed for algicidal activity against *C. antiqua* and it was found  
46 that all of **1**~**4** had high activity (>80% mortality at 24 h) at a concentration of 20 µg/mL. A  
47 structure–activity relationship study using 11 related compounds revealed that the presence of the  
48 hydroxyl group is important for the activity and the double bond may be replaced with a triple bond.

49  
50 Keywords

51 *Tricleocarpa jejuensis*; hydroxylated *trans*-unsaturated fatty acid; oxylipin; anti-microalgal activity;  
52 *Chattonella antiqua*.

53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72

73 1. Introduction

74

75 Harmful algal blooms (HABs), commonly known as red tides, due to eutrophication of coastal  
76 waters occur world-wide and cause serious damage to aquatic ecosystems and public health. The  
77 recent dominant species of HABs in Japan are *Chattonella antiqua* (Raphidophyceae), *Karenia*  
78 *mikimotoi* (Dinophyceae) and *Heterocapsa circularisquama* (Dinophyceae), which have caused  
79 mass mortality of cultivated fish and shellfish. Various physical, chemical, physico-chemical, and  
80 biological methods to control HABs have been developed [1]; however, many of them are  
81 unacceptable for practical use in marine environments due to the second pollution, high cost, or  
82 difficulty of handling.

83 Macroalgae have been shown to produce and release allelopathic substances toxic to HAB  
84 species [2,3]. Consequently, considerable studies on the isolation and identification of the  
85 allelochemicals of macroalgae have been conducted [4] with the goal of developing an  
86 environmentally benign, natural product-based, anti-red tide agent. The algicidal (antialgal)  
87 compounds isolated so far include polyunsaturated fatty acids (PUFAs) from *Cladosiphon*  
88 *okamuranus* [5], *Botryococcus braunii* [6], *Ulva fasciata* [7] *Lithophyllum yessoense* [8], and  
89 *Sargassum thunbergii* [9]; glycerolipids from *Ishige sinicola* [10] and *Ulva prolifera* [11,12];  
90 terpenoids from *Dictyota dichotoma* [13], *Gracilaria lemaneiformis* [14,15], *Dictyopteris undulata*  
91 [16], and *Ulva pertusa* [17]; and phenolics [15,17]. Many of these compounds are reported to have  
92 potent algicidal activity at concentrations of low  $\mu\text{g/mL}$  range against some of the raphidophytes and  
93 dinoflagellates responsible for red tides. We screened 17 species of macroalgae including 9  
94 Rhodophyta, 6 Phaeophyta, and 2 Chlorophyta collected from the coastal region of Nagasaki  
95 Prefecture, Japan, for their algicidal activity against the red tide phytoplankton *Chattonella antiqua*  
96 and found that a methanol extract of the red alga *Tricleocarpa jejuensis* had cell lysis activity at a  
97 concentration of 0.1 mg/mL (Supplementary data). Herein, we describe the separation, structure  
98 elucidation, synthesis and structure–activity study of the algicidal principles of *T. jejuensis*.

99

100 2. Materials and methods

101

102 2.1. General experimental procedure

103

104 NMR spectra were recorded on a Varian System 500PS SN spectrometer (500 MHz for  $^1\text{H}$  and  
105 125 MHz for  $^{13}\text{C}$ ), a JOEL JNM AL400 spectrometer (400 MHz for  $^1\text{H}$  and 100 MHz for  $^{13}\text{C}$ ) or a  
106 Varian Gemini 300 spectrometer (300 MHz for  $^1\text{H}$  and 75 MHz for  $^{13}\text{C}$ ) in  $\text{CDCl}_3$  using  
107 tetramethylsilane and  $\text{CDCl}_3$  as the internal standards for  $^1\text{H}$  and  $^{13}\text{C}$  nuclei, respectively. High  
108 resolution (HR) electron impact mass spectroscopy (EIMS) was carried out on a JEOL JMS-700N

109 spectrometer. Electron spray ionization (ESI) and direct analysis in real time (DART) mass spectra  
110 (MS) were obtained on a JEOL JMS-T100TD spectrometer. IR spectra were recorded on a  
111 ThermoFisher Scientific Nicolet Nexus 670NT spectrophotometer. Optical rotation was measured on  
112 a JASCO P-2200 polarimeter using a 10-cm microcell. GC-EIMS analysis was performed using an  
113 Agilent Technologies GC7890A-MS7000A system equipped with an HP-1MS capillary column  
114 (length 30 m, inside diameter 0.250 mm, film thickness 0.25  $\mu\text{m}$ ) in EI mode at 70 eV. GLC  
115 conditions: carrier gas, He; flow rate, 1.8 mL/min; oven, 120  $^{\circ}\text{C}$ , 5 min isothermal, 120  $^{\circ}\text{C}$ ~300  $^{\circ}\text{C}$   
116 with 10  $^{\circ}\text{C}/\text{min}$ .

117 Silica gel gravity and medium pressure column chromatography separations were performed  
118 using Kanto Chem. Co. Ltd. Silica Gel N (spherical neutral) 100-210  $\mu\text{m}$  and 40-60  $\mu\text{m}$ ,  
119 respectively. Preparative TLC was performed using Merck Silica Gel 60 F<sub>254</sub> (20  $\times$  20 cm, layer  
120 thickness 1.0 mm).

121

## 122 2.2. Plant material

123

124 A specimen of *T. jejuensis* was collected from Ishigaki Island of Okinawa Prefecture, Japan, in  
125 June 2016. All samples were saved in a freezer and brought to the laboratory in plastic bags. After  
126 thawing at rt (ca. 25  $^{\circ}\text{C}$ ), the samples were briefly washed with tap water to remove possible  
127 contaminants, and dried in air.

128

## 129 2.3. Cultivation of phytoplankton

130

131 *Chattonella antiqua*, isolated from Shimabara Bay, Japan in 2010 by Dr. Tatsuya Oda, Nagasaki  
132 University, was cultured aseptically in PES medium at 20  $^{\circ}\text{C}$  under 40  $\mu\text{mol}/\text{m}^2/\text{s}$  using 40 W  
133 fluorescent lamps with a 12 h day cycle and 12 h night cycle and sub-cultured after approximately  
134 14 days.

135

## 136 2.4. Algicidal assay

137

138 The algicidal assay was performed according to Kakisawa's procedure [5], with a slight  
139 modification. In brief, a methanol solution of the extract or sample at varying concentrations was  
140 added to the cell suspension (cell density ca.  $2 \times 10^4$  cells/mL) of *C. antiqua* in a 48-well microplate  
141 to make the final concentrations of 5, 20, or 80  $\mu\text{g}/\text{mL}$  (methanol concentration  $\leq 1\%$ ). After  
142 incubation at 20  $^{\circ}\text{C}$  for 24 h, the cell mortality was calculated under microscope observation ( $\times 400$ ).  
143 The assay was performed in triplicate. Algicidal activity (AA) was calculated using a formula: AA  
144 (%) =  $(1 - T/C) \times 100$ , where T and C represent number of the living cells in the presence and absence

145 of the compound tested, respectively. Swollen and burst cells were considered dead cells.

146

## 147 2.5. Extraction and isolation of algicidal compounds

148

149 *T. jejuensis* (240 g dry wt) was powdered using a blender, extracted twice with MeOH (2 L × 2)  
150 for 3 days, whereupon the MeOH was evaporated under reduced pressure. The crude extract was  
151 partitioned between hexane and 80% aqueous MeOH. After almost of the MeOH had been removed  
152 *in vacuo*, the aqueous layer was partitioned between water and EtOAc. The EtOAc layer (1.5 g) was  
153 separated through HP20 resin by successive elutions with 20%, 40%, 60%, 80%, and 100% MeOH,  
154 and finally with acetone. The active fraction eluted with 100% MeOH (734 mg) was separated by  
155 silica gel column chromatography followed by TLC using hexane-EtOAc (1:1) as the solvent, to  
156 give two active fractions (Fr. TC5-1 and TC5-2). Fr. TC5-2 (13.1 mg) was separated by reversed-  
157 phase HPLC (Capcell Pak C18, 10 mm × 250 mm, 90% MeOH) to give active fraction **f5** (4.4 mg).

158 Fraction **f5**: a colorless oil,  $[\alpha]_D^{20} -0.67^\circ$  (c 0.1, MeOH). ESIMS  $m/z$  321  $[M+Na]^+$ , EIMS (bis-  
159 TMS derivative)  $m/z$  442 ( $M^+$ ), 427, 357, 329, 227, 199. HR-EIMS (bis-TMS derivative) calcd for  
160  $C_{24}H_{50}O_3Si_2$ : 442.3298, found 442.3299.  $^1H$  NMR (500 MHz)  $\delta$  0.879 and 0.881 (3H, t x 2,  $J=7.0$   
161 Hz), 1.22-1.41 (14H, m), 1.42-1.45 (1H, m), 1.45-1.59 (1H, m), 1.64 (2H, m), 2.57 (2H, m), 2.35  
162 (2H, t,  $J=7.5$  Hz), 3.45-3.65 (1H, br), 4.03 (1H, m), 5.443 and 5.445 (1H, dd ×2,  $J=7.1, 1.0$  Hz), 5.62  
163 (1H, m).  $^{13}C$  NMR (125 MHz)  $\delta$  14.05, 14.10, 14.10, 14.11, 22.60, 22.66, 24.58, 24.66, 25.39, 25.43,  
164 25.49, 28.52, 28.75, 28.79, 28.96, 29.11, 29.14, 29.18, 29.26, 29.3, 29.55, 31.36, 31.82, 31.84, 31.87,  
165 32.00, 32.18, 33.54, 37.25, 33.59, 33.59, 37.32, 73.20, 73.25, 73.26, 73.30, 131.97, 132.20, 132.28,  
166 132.31, 132.93, 132.94, 132.97, 133.16, 177.09, 177.12, 177.16, 177.18. IR (KBr)  $\nu_{max}$  980, 1260,  
167 1445, 1710, 2870, 2920  $cm^{-1}$ .

168

## 169 2.6. *p*-Bromophenacyl esterification of **f5**

170

171 A mixture of **f5** (1 mg) and  $K_2CO_3$  (spray dried, 8 mg) in dry acetone (1 mL) was stirred at rt for  
172 15 min. A 0.1 M acetone solution of *p*-bromophenacyl bromide (0.090 mL, 9.0 mmol) was added  
173 and the whole was stirred for 5 h. The mixture was diluted with  $CH_2Cl_2$  (1 mL) and filtered. The  
174 filtrate was concentrated and the residue was purified by silica gel TLC (0.25 mm thickness; 10 x 20  
175 cm; solvent, hexane-EtOAc (2:1)) to afford fraction **f5a** (Rf 0.53, 0.2 mg) and **f5b** (Rf 0.48, 0.2 mg).

176 Fraction **f5a**.  $^1H$  NMR (500 MHz)  $\delta$  0.880 and 0.884 (3H, t × 2,  $J=6.8$  Hz), 1.20-1.65 (21H,  
177 m), 1.69 (1H, m), 2.00-2.05 (2H, m), 2.48 (2H, deformed-t,  $J=7.5$  Hz), 3.67 (1H, s), 4.03  
178 (1H, m), 5.28 (2H, s), 5.41-5.48 (1H, m), 5.59-5.67 (1H, m), 7.64 (2H, d,  $J=8.6$  Hz), 7.78  
179 (2H, d,  $J=8.6$  Hz).

180 Fraction **f5b**.  $^1H$  NMR (500 MHz)  $\delta$  0.877 and 0.881 (3H, t × 2,  $J=7.1$  Hz), 1.20-1.65 (21H,

181 m), 1.66-1.75 (2H, m), 1.99-2.07 (2H, m), 2.479 and 2.483 (2H, t × 2, J=7.5 Hz), 4.03 (1H,  
182 m), 5.29 (2H, s), 5.41-5.48 (1H, m), 5.59-5.67 (1H, m), 7.64 (2H, d, J=8.6 Hz), 7.78 (2H,  
183 d, J=8.6 Hz).

184

## 185 2.7. Chemicals

186

187 (E)-Octadec-9-enoic acid (elaidic acid) was prepared by nitrous acid mediated isomerization of  
188 oleic acid [18]. (R)-(+)-Ricinoleic acid was purchased from Tokyo Kasei, Tokyo.

189

## 190 2.8. Synthesis

191

### 192 2.8.1. Octadec-10-ynoic acid (**5**)

193

194 To a cooled (−78 °C) solution of 10-undecynoic acid (1.00 g, 5.49 mmol) in anhydrous THF  
195 (40 mL) and HMPA (10 mL), was added dropwise via a syringe a 2.5 M cyclohexane solution of  
196 BuLi (5.27 mL, 13.2 mmol) over a period of 30 min. The mixture was warmed up to 0 °C and kept  
197 at this temperature for 2 h. The mixture was cooled again to −78 °C and 1-bromoheptane (0.95 mL,  
198 6.04 mmol) was injected. The whole was stirred at rt for 18 h before being quenched with 10%  
199 NH<sub>4</sub>Cl and 1M HCl solutions. The THF was removed *in vacuo*, and the residue was acidified to pH  
200 1 with 1 M HCl and extracted twice with EtOAc. The organic layer washed with brine, dried over  
201 Na<sub>2</sub>SO<sub>4</sub>, and concentrated *in vacuo*. The crude product was purified with flash chromatography on  
202 silica gel eluted with hexane-EtOAc (4:1) to give **5** (0.421 g, 1.50 mmol, 27 %) as white crystals, mp  
203 43 °C, with 50% recovery of 10-undecynoic acid. <sup>1</sup>H NMR (300 MHz) δ 0.88 (3H, t, J=6.7 Hz),  
204 1.25-1.40 (16H, m), 1.40-1.54 (4H, m), 1.57-1.70 (2H, m), 2.14 (4H, t, J=7.0 Hz), 2.35 (2H, t,  
205 J=7.62 Hz), 9.96-10.42 (1H, br.). <sup>13</sup>C NMR (100 MHz) δ 14.00, 18.65, 22.55, 24.56, 28.69, 28.74,  
206 28.76, 28.87, 28.93, 29.02, 29.07, 29.08, 29.70, 31.70, 33.97, 80.12, 80.29, 180.28. DART-MS *m/z*  
207 (rel intensity) 282 (26), 281 (100), 215 (17), 180 (16). HR-DART-MS [M+H]<sup>+</sup> *m/z* 281.24840 (calcd  
208 for C<sub>18</sub>H<sub>33</sub>O<sub>2</sub>: 281.24806).

209

### 210 2.8.2. Methyl octadec-10-ynoate (**6**)

211

212 To a solution of **5** (123 mg, 0.440 mmol) in a mixture of CH<sub>2</sub>Cl<sub>2</sub> (6 mL) and MeOH (6 mL),  
213 was added 2M ethereal solution of TMSCH<sub>2</sub>N<sub>2</sub> (0.9 mL, 1.8 mmol) and the mixture was stirred at rt  
214 until TLC revealed the disappearance of the acid. The reaction was then quenched with one drop  
215 AcOH and the solvent was removed *in vacuo* to afford **6** (129 mg, 0.439 mmol, 100%) as a colorless  
216 oil. This was used for the next step without further purifications. <sup>1</sup>H NMR (300 MHz) δ 0.88 (3H, t,

217  $J=7.0$  Hz), 1.23-1.41 (16H, m), 1.41-1.54 (4H, m), 1.58-1.70 (2H, m), 2.14 (4H, t,  $J=7.0$ ), 2.30 (2H,  
218 t,  $J=7.6$  Hz), 3.67 (3H, s).

219

### 220 2.8.3. (*Z*)-Methyl octadec-10-enoate (**7**)

221

222 The acetylenic fatty acid ester **6** (107 mg, 0.363 mmol) was hydrogenated over 5% Pd/CaCO<sub>3</sub>  
223 poisoned with Pb (67.8 mg) in EtOAc (10 mL) under H<sub>2</sub> (balloon pressure) for 35 min at rt. The  
224 mixture was filtered through a short column on silica gel and concentrated *in vacuo* to give olefin **7**  
225 (125 mg, 0.420 mmol, 96%) as a pale yellow oil. This was used for the next step without further  
226 purifications. <sup>1</sup>H NMR (500 MHz)  $\delta$  0.88 (3H, t,  $J=7.0$  Hz), 1.22-1.40 (20H, m), 1.55-1.68 (2H, m),  
227 1.96-2.07 (4H, m), 2.30 (2H, t,  $J=7.6$  Hz), 3.66 (3H, s), 5.32-5.37 (2H, m). <sup>13</sup>C NMR (125 MHz)  $\delta$   
228 14.10, 22.67, 24.95, 27.18, 27.20, 29.13, 29.22 ( $\times 2$ ), 29.27, 29.33, 29.72, 29.76, 31.86, 34.10, 51.42,  
229 129.80, 129.94, 174.33. DART-MS  $m/z$  (rel intensity) 298 (20), 297 (100). HR-DART-MS [M+H]<sup>+</sup>  
230  $m/z$  297.28021 (calcd for C<sub>19</sub>H<sub>37</sub>O<sub>2</sub>: 297.27936).

231

### 232 2.8.4. (*E*)-Methyl 9-hydroxyoctadec-10-enoate (**8**) and (*E*)-Methyl 12-hydroxyoctadec-10E-enoate 233 (**9**)

234

235 A mixture of **7** (87.3 mg, 0.294 mmol), SeO<sub>2</sub> (28.0 mg, 0.252 mmol), and *t*-BuOOH (5 M in  
236 decane, 0.213 mL, 1.18 mmol) in dry CH<sub>2</sub>Cl<sub>2</sub> (2 mL) was stirred at rt for 50 h. The reaction was then  
237 quenched by addition of 10% Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> solution (5 mL) and extracted three times with CH<sub>2</sub>Cl<sub>2</sub>. The  
238 combined organic extracts were dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated *in vacuo*. The residue was  
239 purified by a column chromatography on silica gel eluted with hexane-EtOAc (4:1~3:1) to give a  
240 mixture of **8** and **9** (67.6 mg, 0.216 mmol, 74%). Further elution of the column with EtOAc gave  
241 methyl (*E*)-9,12-dihydroxyoctadec-10-enoate (13.9 mg, 0.0424 mmol, 14%) as a mixture of  
242 diastereomers. The mixture of **8** and **9** was separated by MPLC on silica gel eluted with hexane-  
243 EtOAc (9:1).

244 Compound **8**; a colorless oil. <sup>1</sup>H NMR (500 MHz)  $\delta$  0.88 (3H, t,  $J=7.0$  Hz), 1.15-1.38 (18H, m),  
245 1.45-1.75 (5H, m), 1.97-2.07 (2H, m), 2.30 (2H, t,  $J=7.6$  Hz), 3.67 (3H, s), 4.02 (1H, q,  $J=6.5$  Hz),  
246 5.38-5.48 (1H, m), 5.63 (1H, dt,  $J=15.2, 6.5$  Hz). <sup>13</sup>C NMR (75 MHz)  $\delta$  14.11, 22.64, 24.89, 25.39,  
247 29.04, 29.09, 29.13, 29.16, 29.16, 29.31, 31.82, 32.16, 34.06, 37.23, 51.44, 73.18, 132.26, 132.92,  
248 174.30. DART-MS  $m/z$  (rel intensity) 311 (20), 296 (32), 295 (100), 177 (11). HR-DART-MS [M+H-  
249 H<sub>2</sub>O]<sup>+</sup>  $m/z$  295.26377 (calcd for C<sub>19</sub>H<sub>35</sub>O<sub>2</sub>: 295.26371).

250 Compound **9**; a colorless oil. <sup>1</sup>H NMR (500 MHz)  $\delta$  0.88 (t, 3H,  $J=7.0$  Hz), 1.23-1.40 (18H, m),  
251 1.42-1.67 (5H, m), 2.01 (2H, q,  $J=6.7$  Hz), 2.30 (2H, t,  $J=7.0$  Hz), 3.67 (3H, s), 4.03 (1H, q,  $J=6.7$   
252 Hz), 5.44 (1H, ddt,  $J=15.3, 7.0, 1.2$  Hz), 5.62 (1H, dt,  $J=15.3, 7.0$  Hz). <sup>13</sup>C NMR (75 MHz)  $\delta$  14.11,

253 22.58, 24.89, 25.43, 29.00, 29.07, 29.11, 29.15, 29.20, 29.20, 31.80, 31.13, 34.07, 37.32, 51.43,  
254 73.20, 132.07, 133.05, 174.32. DART-MS  $m/z$  (rel intensity) 312 (12), 311 (26), 295 (100), 284 (20),  
255 282 (23), 256 (20). HR-DART-MS  $[M+H-H_2O]^+$   $m/z$  295.26214 (calcd for  $C_{19}H_{35}O_2$ : 295.26371).

256

257 2.8.5. *(E)*-12-Hydroxyoctadec-10-enoic acid (**4**)

258

259 A solution of **9** (21.7 mg, 0.0694 mmol) in a mixture of 10% NaOH (1 mL) and MeOH (4  
260 mL) was heated at reflux for 7.5 h. After cooling, the MeOH was removed *in vacuo*, the aqueous  
261 residue was diluted with water, acidified with 3 M HCl, extracted twice with ether, washed with  
262 brine and concentrated. The crude product was purified by silica gel TLC developed with hexane-  
263 EtOAc (1:1) to give **4** (18.4 mg, 0.0616 mmol, 89%) as white crystals, mp 47.5~49.5 °C.  $^1H$  NMR  
264 (500 MHz)  $\delta$  0.88 (3H, t,  $J=7.0$  Hz), 1.22-1.41 (18H, m), 1.42-1.57 (2H, m), 1.57-1.66 (2H, m), 2.02  
265 (2H, q,  $J=7.0$  Hz), 2.34 (2H, t,  $J=7.3$  Hz), 4.04 (1H, q,  $J=6.7$  Hz), 4.50-6.50 (2H, br), 5.42-5.47 (1H,  
266 m), 5.62 (1H, dt,  $J=15.2, 6.6$  Hz).  $^{13}C$  NMR (125 MHz)  $\delta$  14.07, 22.58, 24.58, 25.42, 28.83 ( $\times 2$ ),  
267 29.01, 29.04, 29.06, 29.20, 31.80, 32.10, 33.95, 37.26, 73.30, 132.19, 132.94, 179.22. DART-MS  $m/z$   
268 (rel intensity) 298 (38), 297 (35), 282 (22), 281 (100), 187 (22). HR-DART-MS  $m/z$   $[M+H-H_2O]^+$   
269 281.24701 (calcd for  $C_{18}H_{33}O_2$ : 281.24806).

270

271 2.8.6. *(E)*-9-Hydroxyoctadec-10-enoic acid (**1**)

272

273 The title compound was obtained from **8** in 85% yield in a similar procedure used for the  
274 synthesis of **4**. Mp 49~50.5 °C.  $^1H$  NMR (500 MHz)  $\delta$  0.88 (3H, t,  $J=7.1$  Hz), 1.22-1.41 (18H, m),  
275 1.12-1.50 (1H, m), 1.51-1.57 (1H, m), 1.58-1.66 (2H, m), 2.02 (2H, q,  $J=7.1$  Hz), 2.34 (2H, t,  $J=7.4$   
276 Hz), 4.03 (1H, q,  $J=6.7$  Hz), 4.67-5.60 (2H, br), 5.41-5.48 (1H, m), 5.62 (1H, dt,  $J=15.4, 6.7$  Hz).  
277  $^{13}C$  NMR (125 MHz)  $\delta$  14.08, 22.64, 24.65, 25.37, 28.96, 29.10, 29.12, 29.15, 29.17, 29.29, 31.82,  
278 32.16, 33.95, 37.21, 73.23, 132.34, 132.85, 179.22. DART-MS  $m/z$  (rel intensity) 298 (31), 297 (22),  
279 282 (42), 281 (100). DART-MS  $m/z$  298, 287, 282, 281, 263. HR-DART-MS  $m/z$   $[M+H-H_2O]^+$   
280 281.24717 (calcd for  $C_{18}H_{33}O_2$ : 281.24806).

281

282 2.8.7. 12-Hydroxyoctadec-10-ynoic acid (**10**)

283

284 To a cooled ( $-78$  °C) and stirred solution of 10-undecynoic acid (424 mg, 2.33 mmol) in dry  
285 THF (24 mL), was added dropwise a 2.5 M solution of BuLi in hexane (2.05 mL, 5.12 mmol). After  
286 10 min at that temperature, the cooling bath was removed and the whole was stirred at rt for 45 min.  
287 The mixture was cooled again to  $-78$  °C and heptanal (293 mg, 2.56 mmol) dissolved in THF (2  
288 mL) was injected. The cooling bath was removed and the mixture was stirred at rt for 1.5 h. The



289 reaction was then quenched with 2 M HCl solution and extracted twice with ether. The ethereal  
290 extracts were combined, washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated. The crude product  
291 was purified by silica gel column chromatography eluted with hexane-EtOAc (2:1) to give **10** (332  
292 mg, 1.12 mmol, 48%) as white crystals, mp 35~36.5 °C. <sup>1</sup>H NMR (300 MHz) δ 0.88 (3H, t, *J*=6.5  
293 Hz), 1.22-1.55 (18H, m), 1.58-1.72 (4H, m), 2.20 (2H, dt, *J*=6.8, 1.8 Hz), 2.34 (2H, t, *J*=7.6 Hz),  
294 4.36 (1H, dt, *J*=6.5, 1.8 Hz), 5.50-6.52 (2H, br). <sup>13</sup>C NMR (125 MHz) δ 14.02, 18.57, 22.53, 24.53,  
295 25.11, 28.50, 28.53, 28.69, 28.77, 28.89, 28.92, 31.72, 33.97, 38.05, 62.68, 81.21, 85.36, 179.47.

296

#### 297 2.8.8. (*E*)-12-Hydroxyoctadec-10-enoic acid (**4**) from compound **10**

298

299 Clean cut Li (42 mg, 6.0 mmol) was added in small portions to liquid NH<sub>3</sub> (ca. 3 mL) at –  
300 78 °C. After 10 min, a solution of **10** (35.4 mg, 0.120 mmol) in dry THF/*t*-BuOH (3:1, 1.5 mL) was  
301 added as drops to the deep blue solution of Li metal in liquid NH<sub>3</sub> and the mixture was stirred at this  
302 temperature for 2 h. The reaction was quenched by addition of solid NH<sub>4</sub>Cl (0.5 g) and the cooling  
303 bath was removed. After the NH<sub>3</sub> was evaporated, the residue was acidified with 3 M HCl solution,  
304 extracted twice with ether, washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and evaporated. The crude product  
305 was purified by reversed-phase HPLC (Capcell Pak C18, 10 mm x 250 mm) eluted with 85%  
306 CH<sub>3</sub>CN to give **4** (13.7 mg, 0.0459 mmol, 38%) as white crystals, mp 49~51 °C.

307

#### 308 2.8.9. (*E*)-Methyl 11-hydroxyoctadec-12-enoate (**13**)

309

310 A 1.0 M toluene solution of DIBAL (1.36 mL, 1.36 mmol) was injected via a syringe to a  
311 stirred solution of 1-heptyne (153 mg, 1.59 mmol) in dry hexane (4 mL) at rt under Ar atmosphere.  
312 After the mixture had been stirred at 60 °C for 5 h, it was cooled to –78 °C (dry ice-acetone bath)  
313 and a solution of **11** (195 mg, 0.909 mmol) in toluene (2 mL) was added as drops. After 20 min, the  
314 cooling bath was replaced with an ice-salt bath and the mixture was stirred for 1 h. The reaction was  
315 then quenched by addition of a saturated solution of Rochelle's salt (0.2 mL), stirred overnight, dried  
316 over MgSO<sub>4</sub>, and filtered through a pad of Celite, washed well with EtOAc, and concentrated. The  
317 crude product was purified by a column chromatography on silica gel eluted with hexane-EtOAc  
318 (5:1) to give **13** (79.3 mg, 0.254 mmol, 28%) as a pale yellow oil with 77.0 mg (40%) recovery of  
319 aldehyde **11**. <sup>1</sup>H NMR (300 MHz) δ 0.89 (3H, deformed t, *J*=7.0 Hz), 1.20-1.55 (19H, m), 1.55-1.68  
320 (4H, m), 2.02 (2H, q, *J*=7.0 Hz), 2.30 (2H, t, *J*=7.6 Hz), 3.67 (3H, s), 4.02 (1H, q, *J*=6.5 Hz), 5.44  
321 (1H, dd, *J*=15.4, 7.0 Hz), 5.63 (1H, dt, *J*=15.4, 6.8 Hz). <sup>13</sup>C NMR (75 MHz) δ 14.02, 22.47, 24.91,  
322 25.44, 28.83, 29.09, 29.18, 29.32, 29.47 (×2), 31.32, 32.11, 34.07, 37.27, 51.42, 73.18, 132.17,  
323 132.97, 174.33. DART-MS *m/z* (rel intensity) 312 (15), 296 (22), 295 (100), 293 (38), 282 (21). HR-  
324 DART-MS *m/z* [M+H-H<sub>2</sub>O]<sup>+</sup> 295.26347 (calcd for C<sub>19</sub>H<sub>35</sub>O<sub>2</sub>: 295.26371).

325

326 2.8.10. (*E*)-11-hydroxyoctadec-12-enoic acid (**3**)

327

328 The title compound was obtained by alkaline hydrolysis of **13** at rt in 78% yield in a similar  
329 procedure used for the synthesis of **4**. White crystals, mp 49 °C. <sup>1</sup>H NMR (500 MHz) δ 0.89 (3H, t,  
330 *J*=7.1 Hz), 1.23-1.41 (19H, m), 1.41-1.57 (2H, m), 1.57-1.668 (2H, m), 2.03 (2H, q, *J*=7.1 Hz), 2.34  
331 (2H, t, *J*=7.3 Hz), 4.04 (1H, q, *J*=6.8 Hz), 5.44 (1H, dd, *J*=15.4, 7.1 Hz), 5.63 (1H, dt, *J*=15.4, 6.7  
332 Hz), 9.75-9.77 (1H, br.). <sup>13</sup>C NMR (125 MHz) δ 14.03, 22.49, 24.65, 25.43, 28.85, 29.00, 29.16,  
333 29.28, 29.46 (×2), 31.34, 32.13, 33.93, 37.26, 73.27, 132.29, 132.89, 179.23. DART-MS *m/z* (rel  
334 intensity) 298 (36), 297 (46), 282 (76), 281 (100). HR-DART-MS *m/z* [M+H-H<sub>2</sub>O]<sup>+</sup> 281.24863  
335 (calcd for C<sub>18</sub>H<sub>33</sub>O<sub>2</sub>: 281.24806).

336

337 2.8.11. 9-(Tetrahydropyran-2-yl)oxy-1-nonyne (**15**)

338

339 A solution of **14** (1.10 g, 7.83 mmol), dihydro-2*H*-pyran (1.24 g, 14.7 mmol), and *p*-  
340 TsOH·H<sub>2</sub>O (0.05 g) in dry CH<sub>2</sub>Cl<sub>2</sub> (60 mL) was stirred at rt for 18 h. The mixture was then washed  
341 with 5% NaHCO<sub>3</sub> solution (30 mL) and the aqueous layer was extracted with CH<sub>2</sub>Cl<sub>2</sub>. The combined  
342 organic extracts were washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated. The oily residue was  
343 chromatographed on silica gel eluted with hexane-Et<sub>2</sub>O (19:1) to give **15** (1.52 g, 6.76 mmol, 69 %)   
344 as a colorless oil. <sup>1</sup>H NMR (300 MHz) δ 1.21-1.46 (6H, m), 1.46-1.67 (8H, m), 1.67-1.90 (2H, m),  
345 1.94 (1H, t, *J*=2.6 Hz), 2.18 (2H, dt, *J*=6.9, 2.6 Hz), 3.38 (1H, dt, *J*=9.5, 6.7 Hz), 3.45-3.56 (1H, m),  
346 3.73 (1H, dt, *J*=9.5, 7.0 Hz), 3.81 3.94 (1H, m), 4.58 (1H, dd, *J*=4.1, 2.8 Hz). <sup>13</sup>C NMR (75 MHz) δ  
347 18.33, 19.66, 25.44, 26.07, 28.36, 28.63, 28.89, 29.64, 30.72, 62.30, 67.55, 68.06, 84.66, 98.80.  
348 DART-MS *m/z* (rel intensity) 225 (9), 169 (12), 102 (19), 85 (100). HR-DART-MS *m/z* [M+H]<sup>+</sup>  
349 225.18602 (calcd for C<sub>14</sub>H<sub>25</sub>O<sub>2</sub>: 225.18546).

350

351 2.8.12. (*E*)-18-(Tetrahydropyran-2-yl)oxyoctadec-10-en-9-ol (**16**)

352

353 To a solution of **15** (0.758 g, 3.38 mmol) in dry hexane (5 mL), a 1 M toluene solution of  
354 DIBAL (3.71 mL, 3.71 mmol) was added dropwise at rt under Ar atmosphere, and the mixture was  
355 stirred at 60 °C for 2h. The mixture was then cooled to -78 °C and nonanal (0.577 mg, 4.06 mmol)  
356 dissolved in toluene (4 mL) was added dropwise. After 2 h at -60 °C, the reaction mixture was  
357 warmed up to rt, quenched with water, and acidified with 1 M HCl. The whole was extracted twice  
358 with EtOAc, washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated. The oily residue was  
359 chromatographed on silica gel eluted with hexane-EtOAc (7:1 to 2:1) gave crude **16** (0.548 g, 1.49  
360 mmol, 44%) as a mixture of diastereomers. DART-MS *m/z* (rel intensity) 367 (5), 352 (33), 351

361 (100), 333 (37), 283 (37), 281 (33), 85 (94).

362

363 2.8.13. (*E*)-10-(Acetoxy)octadec-8-en-1-ol (**18**)

364

365 A solution of the crude **16** (0.548 mg, 1.49 mmol) in Ac<sub>2</sub>O (0.5 mL) and pyridine (1 mL) was  
366 stirred at rt for 40 h. The reaction was quenched with water, acidified with 2 M HCl, and extracted  
367 twice with ether. The ethereal extracts were combined, washed with 5% NaHCO<sub>3</sub>, dried over  
368 Na<sub>2</sub>SO<sub>4</sub>, and concentrated. The crude product **17** was dissolved EtOH (12 mL) and a catalytic  
369 amount of *p*-TsOH·H<sub>2</sub>O (0.05 g) was added. After the mixture had been stirred at rt for 25 min, it  
370 was concentrated and purified by a column chromatography on silica gel eluted with hexane-EtOAc  
371 (3:1) to give **18** (160 mg, 0.491 mmol, 33%) as a pale yellow oil. <sup>1</sup>H NMR (300 MHz) δ 0.88 (3H, t,  
372 *J*=6.7 Hz), 1.19-1.44 (20H, m), 1.46-1.68 (4H, m), 1.61 (1H, br. s), 1.96-2.09 (2H, m), 2.04 (3H, s),  
373 3.64 (2H, t, *J*=6.7 Hz), 5.17 (1H, q, *J*=7.0 Hz), 5.36 (1H, m), 5.68 (1H, dt, *J*=15.5, 6.8 Hz). <sup>13</sup>C  
374 NMR (75 MHz) δ 14.07, 21.38, 22.62, 25.15, 25.60, 28.78, 28.95, 29.14, 29.18, 29.32, 29.44, 31.81,  
375 32.11, 32.66, 34.46, 62.93, 75.12, 128.33, 134.31, 170.50. DART-MS *m/z* (rel intensity) 326 (22),  
376 267 (100), 429 (61), 177 (56). HR-DART-MS [M+H-AcOH]<sup>+</sup> *m/z* 267.26766 (calcd for C<sub>18</sub>H<sub>35</sub>O:  
377 267.26879).

378

379 2.8.14. (*E*)-10-(Acetoxy)octadec-8-enoic acid (**19**)

380

381 A mixture of **18** (66.5 mg, 0.204 mmol) and PDC (268 mg, 0.713 mmol) dry DMF (2 mL)  
382 was stirred at rt for 17 h. The mixture was poured into water, extracted twice with ether, washed with  
383 brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated. The crude product was purified by a column  
384 chromatography on silica gel eluted with hexane-EtOAc (3:1) to give **19** (31.6 mg, 0.0928 mmol,  
385 45%) as a pale oil. <sup>1</sup>H NMR (300 MHz) δ 0.88 (3H, t, *J*=7.0 Hz), 1.16-1.41 (18H, m), 1.48-1.69 (4H,  
386 m), 1.95-2.11 (2H, m), 2.04 (3H, s), 2.34 (2H, t, *J*=7.3 Hz), 5.17 (1H, q, *J*=7.0 Hz), 5.36 (1H, m),  
387 5.67 (1H, dt, *J*=15.7, 7.0 Hz), 9.75-9.77 (1H, br.). <sup>13</sup>C NMR (75 MHz) δ 14.09, 21.42, 22.63, 24.56,  
388 25.18, 28.65, 28.80, 29.07, 29.21, 29.33, 29.46, 31.82, 32.07, 33.99, 34.48, 75.14, 128.45, 134.16,  
389 170.58, 179.93. DART-MS *m/z* (rel intensity) 281 [M+H-AcOH]<sup>+</sup> (42), 89 (100), 61 (38). HR-  
390 DART-MS [M+H-AcOH]<sup>+</sup> *m/z* 281.24687 (calcd for C<sub>18</sub>H<sub>33</sub>O<sub>2</sub>: 281.24805).

391

392 2.8.15. (*E*)-10-Hydroxyoctadec-8-enoic acid (**2**)

393

394 The title compound was obtained by alkaline hydrolysis of **19** at rt in 36% yield in a similar  
395 procedure used for the synthesis of **1**. Pale yellow crystals, mp 49 °C. <sup>1</sup>H NMR (500 MHz)  
396 δ 0.88 (3H, t, *J*=6.9 Hz), 1.18-1.58 (19H, m), 1.59-1.72 (3H, m), 2.04 (2H, t, *J*=7.3 Hz), 2.34 (2H, t,

397  $J=7.3$  Hz), 3.45-3.73 (2H, br.), 4.04 (1H, q,  $J=6.7$  Hz), 5.41-5.48 (1H, m), 5.61 (1H, dt,  $J=15.4$ , 6.9  
398 Hz).  $^{13}\text{C}$  NMR (125 MHz)  $\delta$  14.09, 22.65, 24.57, 25.47, 28.26, 28.57, 28.76, 28.82, 29.25, 29.54,  
399 31.86, 32.01, 33.88, 37.28, 73.27, 131.97, 133.12, 179.08. DART-MS  $m/z$  (rel intensity) 299 (31),  
400 298 (34), 282 (38), 281 (100), 279 (58). HR-DART-MS  $[\text{M}+\text{H}-\text{H}_2\text{O}]^+$   $m/z$  281.24670 (calcd for  
401  $\text{C}_{18}\text{H}_{34}\text{O}_3$ : 281.24805).

402

#### 403 2.8.16. *1,10-Dihydroxyoctadec-8-ene (20)*

404

405 A solution of **16** (167.7 mg, 0.455 mmol) in MeOH (3 mL) containing a catalytic amount of  
406 PPTS was allowed to stand at rt for 20 h. After the MeOH had been removed *in vacuo*, the residue  
407 was chromatographed on silica gel eluted with hexane-EtOAc (2:1) to give **20** (59.5 mg, 0.209  
408 mmol, 46%) as a colorless oil.  $^1\text{H}$  NMR (300 MHz)  $\delta$  0.88 (3H, t,  $J=7.0$  Hz), 1.22-1.62 (25H, m),  
409 1.64-1.71 (1H, m), 2.02 (1H, q,  $J=7.0$  Hz), 2.21 (1H, dt,  $J=7.0$ , 1.8 Hz), 3.64 (2H, t,  $J=6.7$  Hz), 4.03  
410 (1H, q,  $J=6.8$  Hz), 5.44 (1H, m), 5.62 (1H, dt,  $J=15.2$ , 6.7 Hz). DART-MS  $m/z$  (rel intensity) 284  
411 (18), 283 (33), 281 (25), 267 (71), 265 (100), 249 (38), 247 (31). HR-DART-MS  $[\text{M}+\text{H}-\text{H}_2\text{O}]^+$   $m/z$   
412 267.26888 (calcd for  $\text{C}_{18}\text{H}_{35}\text{O}$ : 267.26879).

413

#### 414 2.8.17. *p-Bromophenacyl ester of oleic acid (25)*

415

416 A mixture of oleic acid (2.82 g, 10.0 mmol) and  $\text{K}_2\text{CO}_3$  (2.28 g, 16.0 mmol) in dry acetone  
417 (25 mL) was stirred at rt for 30 min. *p*-Bromophenacyl bromide (3.06 g, 11.0 mmol) was then added  
418 and the whole was stirred overnight. The reaction mixture was filtrated and the filtrate was  
419 evaporated. The residue was then extracted with diethyl ether, washed with 5%  $\text{NaHCO}_3$  solution,  
420 dried over anhydrous  $\text{Na}_2\text{SO}_4$ , and concentrated. The crystalline product was recrystallized from  
421 methanol, washed with hexane to remove unreacted oleic acid, giving **25** as a pale yellow powder.  
422  $^1\text{H}$  NMR (400 MHz)  $\delta$  0.88 (3H, t,  $J=7.3$  Hz), 1.17-1.45 (20H, m), 1.70 (2 H, m), 1.96-2.08 (4H, m),  
423 2.48 (2H, t,  $J=7.3$  Hz), 5.28 (2H, s), 5.32-5.38 (2H, m), 7.64 (2H, d,  $J=8.8$  Hz), 7.78 (2H, d,  $J=8.8$   
424 Hz).  $^{13}\text{C}$  NMR (100 MHz)  $\delta$  14.09, 22.67, 24.85, 27.14, 27.18, 29.03, 29.06, 29.14, 29.29 ( $\times 2$ ),  
425 29.49, 29.67, 29.74, 31.87, 33.84, 65.61, 129.05, 129.21 ( $\times 2$ ), 129.73, 129.95, 132.18 ( $\times 2$ ), 132.94,  
426 173.15, 191.44.

427

#### 428 2.8.18. *Selenium dioxide oxidation of oleate 25 (26, 27, and 28)*

429

430 Compounds **26**, **27**, and **28** were synthesized in a similar procedure to that of **8** and **9** in 22%,  
431 20%, and 13% yields, respectively.

432 Compound **26**:  $^1\text{H}$  NMR (300 MHz)  $\delta$  0.88 (3H, t,  $J=6.9$  Hz), 1.15-1.18 (23H, m), 1.97-2.13

433 (2H, m), 2.49 (2H, t,  $J=7.7$  Hz), 4.04 (1H, q,  $J=6.6$  Hz), 5.29 (2H, s), 5.45 (1H, dt,  $J=15.4, 7.2$  Hz),  
434 5.58-5.69 (1H, m), 7.64 (2H,  $J=8.8$  Hz), 7.78 (2H, d,  $J=8.8$  Hz).

435 Compound **27**:  $^1\text{H}$  NMR (300 MHz)  $\delta$  0.88 (3H, t,  $J=6.6$  Hz), 1.18-1.79 (23H, m), 2.02 (2H, q,  
436  $J=6.8$  Hz), 2.48 (2H, t,  $J=7.7$  Hz), 4.04 (1H, q,  $J=6.7$  Hz), 5.29 (2H, s), 5.45 (1H,  $J=15.4, 7.3$  Hz),  
437 5.57-5.68 (1H, m), 7.64 (2H, dt,  $J=8.8, 1.9$  Hz), 7.78 (2H, dt,  $J=8.8, 1.9$  Hz).

438 Compound **28**:  $^1\text{H}$  NMR (300 MHz)  $\delta$  0.88 (3H, t,  $J=6.6$  Hz), 1.15-1.85 (24H, m), 2.49 (2H, t,  
439  $J=7.4$  Hz), 4.09-4.16 (2H, m), 5.29 (2H, s), 5.66-5.74 (2H, m), 7.64 (2H, d,  $J=8.8$  Hz), 7.78 (2H, d,  
440  $J=8.8$  Hz).

441

442 2.8.19. (*E*)-8-Hydroxyoctadec-9-enoic acid (**21**)

443

444 The title compound was obtained by an alkaline hydrolysis of **26** at 50 °C in 38% yield in a  
445 similar procedure used for the synthesis of **4**. White crystals, mp 52-54 °C (lit. mp 54-55 °C) [19].  $^1\text{H}$   
446 NMR (500 MHz)  $\delta$  0.88 (3 H, m,  $J=7.0$  Hz) 1.18-1.40 (18H, m), 1.43-1.68 (4H, m), 2.02 (2 H, q,  
447  $J=6.9$  Hz), 2.33 (2H, t,  $J=7.5$  Hz), 4.04 (1H, q,  $J=6.7$  Hz), 5.44 (1H, dd,  $J=15.28, 7.21$  Hz, 1 H), 5.63  
448 (1 H, dt,  $J=15.2, 6.6$  Hz), 6.20 (2H, br. s).  $^{13}\text{C}$  NMR (125 MHz)  $\delta$  14.07, 22.63, 24.62, 25.45, 28.81,  
449 28.93, 28.97, 29.02, 29.24, 29.48, 31.79, 32.07, 34.05, 37.23, 73.28, 132.11, 132.93, 179.56. EIMS  
450 (bis TMS derivative)  $m/z$  (rel intensity) 442 ( $\text{M}^+$ , 6), 427 (9), 274 (13), 242 (21), 241 (100).

451

452 2.8.20. (*E*)-11-Hydroxyoctadec-9-enoic acid (**22**)

453

454 The title compound was obtained from **27** in 34% yield in the same procedure used for the  
455 synthesis of **4**. Mp 43-46 °C (lit. mp 43-44 °C) [19].  $^1\text{H}$  NMR (500 MHz)  $\delta$  0.88 (3H, t,  $J=7.0$  Hz),  
456 1.18-1.42 (18H, m), 1.42-1.68 (4H, m), 2.02 (2H, q,  $J=7.1$  Hz), 2.32 (2H, t,  $J=7.6$  Hz), 4.04 (1H, q,  
457  $J=6.9$  Hz), 5.43 (1H, m), 5.46 (1H, dt,  $J=15.2, 6.9$  Hz), 5.80-6.97 (2H, br. s).  $^{13}\text{C}$  NMR (125 MHz)  $\delta$   
458 14.09, 22.65, 24.64, 25.48, 28.83, 28.95, 28.99, 29.04, 29.26, 29.50, 31.81, 32.09, 34.07, 37.25,  
459 73.30, 132.13, 132.95, 179.58. HR-EI MS [ $\text{M}-\text{H}_2\text{O}+\text{H}$ ] $^+$   $m/z$  281.24672 (calcd for  $\text{C}_{18}\text{H}_{33}\text{O}_2$ :  
460 281.24806). EIMS (bis TMS derivative)  $m/z$  (rel intensity) 442 ( $\text{M}^+$ , 5), 427 (9), 345 (14), 344 (40),  
461 343 (100), 227 (7).

462

463 2.8.21. 8,11-Dihydroxyoctadec-9-enoic acid (**23**)

464

465 The title compound was obtained from **28** in 19% yield in the same procedure used for the  
466 synthesis of **4**. A colorless oil.  $^1\text{H}$  NMR (300 MHz)  $\delta$  0.88 (3H, t,  $J=7.2$  Hz), 1.20-1.43 (17H, m),  
467 1.43-1.70 (6H, m), 2.35 (2H, t,  $J=7.5$  Hz), 3.38-3.57 (1H, m), 3.65-3.83 (1H, m), 4.05-4.18 (2H, m),  
468 5.63-5.73(2H, m).

469

470 2.8.22. (*E*)-11-Oxo-octadec-9-enoic acid (**24**)

471

472 To a stirred solution of **22** (57.3 mg, 0.192 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (1 mL), Dess-Martin periodinane  
473 (169.7 mg, 0.400 mmol) was slowly added at rt. After stirring overnight, the reaction was quenched  
474 by adding 10% Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> solution. The CH<sub>2</sub>Cl<sub>2</sub> layer was separated and the aqueous layer was  
475 extracted with CH<sub>2</sub>Cl<sub>2</sub>. Organic layers were combined and dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>.

476 Purification by a column chromatography on silica gel eluted with hexane-EtOAc (2:1) and then  
477 reversed phase HPLC (COSMOSIL 5C18-MS-II, 90% methanol) gave **24** (36.0 mg, 0.121 mmol,  
478 63%) as white crystals, mp 52~54 °C. <sup>1</sup>H NMR (300 MHz) δ 0.88 (3H, t, *J*=7.0 Hz), 1.18-1.72 (21H,  
479 m), 2.20 (2H, q, *J*=7.1 Hz), 2.36 (2H, t, *J*=7.0 Hz), 2.53 (2H, t, *J*=7.3 Hz), 6.09 (1H, d, *J*=15.8 Hz),  
480 6.82 (1H, dt, *J*=15.8, 7.0 Hz). <sup>13</sup>C NMR (75 MHz) δ 14.04, 22.57, 24.31, 24.57, 27.98, 28.86, 28.91,  
481 28.94, 29.06, 29.24, 31.65, 32.36, 33.97, 40.06, 130.30, 147.27, 172.16, 201.21. HR-DART-MS  
482 [M+H]<sup>+</sup> *m/z* 297.24295 (calcd for C<sub>18</sub>H<sub>33</sub>O<sub>3</sub>: 297.24297).

483

484 3. Results and discussion

485

486 3.1. Structure elucidation of the algicidal compounds of *T. jejuensis*

487

488 Separation of the methanol extract of *T. jejuensis* by monitoring the algicidal activity against  
489 *C. antiqua* afforded an inseparable mixture of compounds with 100% mortality to the phytoplankton  
490 at 20 µg/mL. The mixture fraction, named **f5**, showed a single molecular ion peak at *m/z* 321  
491 [M+Na]<sup>+</sup> by ESI-MS, indicating that the active compounds were isomeric to each other. A molecular  
492 formula of C<sub>18</sub>H<sub>34</sub>O<sub>3</sub> was established by HR-EI-MS of the bistrimethylsilyl derivative of the mixture.  
493 <sup>13</sup>C NMR spectrum showed signals for carboxylic carbons at δ 177.09, 177.12, 177.16, and 177.18,  
494 olefinic carbons at δ 131.97, 132.20, 132.28, 132.31, 132.93, 132.94, 132.97, and 133.16,  
495 hydroxymethine carbons at δ 73.20, 73.25, 73.26, and 73.30, many methylene carbons, and  
496 overlapping methyl carbons in the sp<sup>3</sup> carbon region. The <sup>1</sup>H NMR spectrum showed overlapping  
497 signals of multiplets at δ 5.59-5.66 (1H), two doublet of doublets at δ 5.443 (*J*=15.4, 7.1 Hz) and δ  
498 5.445 (*J*=15.3, 7.2 Hz), and quartets at δ 4.036 (*J*=6.7 Hz) and 4.031 (*J*=6.7 Hz), indicating the  
499 presence of substructure –CH=CH-CH(OH)- having *E*-configuration. At this stage, the active  
500 compounds were assumed to be four isomeric hydroxylated C18 *trans*-monounsaturated fatty acids.  
501 The positions of the double bonds and hydroxyl groups were determined by the EI-MS  
502 fragmentation pattern. The EI-MS of the bistrimethylsilyl derivatives of the mixture showed four  
503 distinct fragment ion peaks at *m/z* 227, 329, 199 and 357, which corresponded to the fragment ions  
504 of CH<sub>3</sub>(CH<sub>2</sub>)<sub>6</sub>CH=CHCHOTMS, TMSOCHCH=CH(CH<sub>2</sub>)<sub>6</sub>COOTMS,

505  $\text{CH}_3(\text{CH}_2)_4\text{CH}=\text{CHCHOTMS}$ , and  $\text{TMSOCHCH}=\text{CH}(\text{CH}_2)_8\text{COOTMS}$ , respectively, from cleavage  
506 at the allylic positions adjacent to the hydroxyl groups [20,21]. From these spectroscopic data, the  
507 active compounds were assigned to be (*E*)-9-hydroxyoctadec-10-enoic acid (**1**), (*E*)-10-  
508 hydroxyoctadec-8-enoic acid (**2**), (*E*)-11-hydroxyoctadec-12-enoic acid (**3**), and (*E*)-12-  
509 hydroxyoctadec-10-enoic acid (**4**) (Fig. 1).

510

### 511 3.2. Synthesis of the hydroxy monounsaturated fatty acids **1~4**

512

513 To confirm the structure elucidation as well as to obtain pure samples for evaluation of the  
514 algicidal activity of each acid and its related compounds, we synthesized each of the acids **1~4** by  
515 unambiguous routes.

516 (*E*)-9-Hydroxyoctadec-10-enoic acid (**1**) and (*E*)-12-hydroxyoctadec-10-enoic acid (**4**) are  
517 regioisomeric in the hydroxyl group position; thus both could be obtained from the same  
518 intermediate, (*Z*)-octadec-10-enoate (**7**), using a selenium dioxide allylic oxidation (Scheme 1).  
519 Alkylation of the lithium acetylide of 10-undecynoic acid with 1-bromoheptane followed by methyl  
520 esterification gave C18 acetylenic acid ester **6** in 27% yield. Partial hydrogenation of the triple bond  
521 of **6** with Lindlar's catalyst followed by selenium dioxide oxidation [19] of the resulting (*Z*)-olefin **7**  
522 afforded an equimolar mixture of alcohols **8** and **9** in 74% combined yield along with a trace of the  
523 9,12-dihydroxylated compound (14% yield). After separation of the regioisomeric monoalcohols by  
524 silica gel chromatography, each methyl ester was hydrolyzed to obtain (*E*)-9-hydroxyoctadec-10-  
525 enoic acid (**1**) and (*E*)-12-hydroxyoctadec-10-enoic acid (**4**). Compound **4** was also synthesized by a  
526 Birch reduction of alkynol **10**, which was obtained by acetylenic addition of 10-undecynoic acid to  
527 heptanal.

528 Syntheses of 11- and 10-hydroxyoctadecenoic acids (**3** and **2**) were achieved via addition  
529 reactions of alkenyl aluminum reagents (Scheme 2 and 3). Aldehyde **11**, prepared by a Kornblum  
530 oxidation of methyl 11-bromoundecanoate using a reported procedure [22], was reacted with alkenyl  
531 aluminum **12** prepared *in situ* from heptyne and DIBAL to give (*E*)-11-hydroxy-12-octadecanoate  
532 (**13**) in 28% yield. One of the target compounds (*E*)-11-hydroxyoctadec-12-enoic acid (**3**) was  
533 obtained by alkaline hydrolysis of **13** (Scheme 2). In this strategy, synthesis of another  
534 hydroxyoctadecenoic acid, **2**, required alkyne **15** as the source of alkenyl aluminum, which was  
535 prepared through an acetylene zipper reaction of commercially available 2-nonyn-1-ol according to a  
536 reported procedure [23]. After THP protection of the hydroxyl group of **14**, alkyne **15** was reacted  
537 with DIBAL to generate alkenyl aluminum, which was then trapped with nonanal to afford the *trans*-  
538 allylic alcohol **16** in 44% yield. The secondary hydroxyl group was protected as the acetate, and then  
539 the primary hydroxyl group was oxidized to furnish (*E*)-10-hydroxyoctadec-8-enoic acid (**2**) after  
540 hydrolytic removal of the acetyl group (Scheme 3).

541

### 542 3.3. Verification of the proposed structures of **1~4** and stereochemistry

543

544 Chemical shift values of selected carbons of the natural products **f5** and synthesized compounds  
545 **1~4** are listed with chemical shift difference values ( $\Delta$ ) in Table 1. The chemical shift values of all  
546 the carbons of **f5** exactly matched those of the corresponding carbons of synthesized compounds **1~4**  
547 within a 0.36-ppm difference with the exception of the carboxyl carbons. The slight change in the  
548 chemical shift values of the carboxyl carbons between the natural products and synthesized  
549 compounds might be due to a considerable difference in the concentration of the sample solutions  
550 prepared for NMR measurements. Indeed, 10-fold dilution of the NMR sample solution of **3** from 30  
551 mg/mL to 3 mg/mL resulted in a 1.59-ppm upfield shift in the carboxyl carbon signal.

552 Recorded specific rotation value of **f5** was close to zero ( $-0.67^\circ$ ). Esterification of **f5** with *p*-  
553 bromophenacyl bromide ( $K_2CO_3$ , acetone, rt) gave two separable fractions on silica gel TLC  
554 (hexane:EtOAc=2:1), named as **f5a** (Rf 0.50) and **f5b** (Rf 0.43), the former being a mixture of the *p*-  
555 bromophenacyl esters of **3** (**3a**, Rf 0.49) and **4** (**4a**, Rf 0.49), and the latter being the esters of **1** (**1a**,  
556 Rf 0.46) and **2** (**2a**, Rf 0.43). HPLC analysis of **f5b** using a chiral column, Chiralpak AD-H (solvent,  
557 2-propanol:hexane=15:85; flow rate, 0.5 mL/min) showed two pairs of peaks of almost equal  
558 intensities corresponding to the respective enantiomers of **1a** ( $t_R$  35.7 and 37.7 min) and **2a** ( $t_R$  40.7  
559 and 45.0 min), indicating **1** and **2** were isolated as racemates (Supplementary data, Fig. 23). On the  
560 other hand, **f5a** showed two peaks at  $t_R$  41.6 min and 45.0 min in an area ratio of 1:3. In the same  
561 HPLC conditions, synthesized ( $\pm$ )-**3a** was separated into two peaks at  $t_R$  41.6 min and 44.8 min,  
562 suggesting the isolated **3** was a racemate (Supplementary data, Fig. 24). However, ( $\pm$ )-**4a** was unable  
563 to separate by this chiral column and appeared as a single peak at nearly 45.0 min (46.1 min).  
564 Finally, separation of ( $\pm$ )-**4a** was achieved by using Chiralpak IA (solvent, MeOH, flow rate 0.5  
565 mL/min) and analysis of **f5a** revealed that the isolated compound **4** was a racemate (Supplementary  
566 data, Fig. 25).

567

### 568 3.4. Algicidal activity of hydroxylated *trans*-monounsaturated fatty acids **1~4** and their derivatives

569

570 Each of the synthesized hydroxy acids **1~4** as well as their synthetic intermediates **10** and **20**  
571 were evaluated for algicidal activity against *C. antiqua* (Fig. 2). For comparison, autoxidation  
572 products of oleic acid, (*E*)-8-hydroxyoctadec-9-enoic acid (**21**) and (*E*)-11-hydroxyoctadec-9-enoic  
573 acid (**22**) [24], their oxidized derivatives, diol **23** and ketone **24** (Scheme 4), (*Z*)-12-hydroxyoctadec-  
574 9-enoic acid (ricinoleic acid), and (*E*)-octadec-9-enoic acid (elaidic acid) were tested for algicidal  
575 activity. All the compounds isolated from *T. jejuensis* except for compound **1** showed complete  
576 toxicity to the phytoplankton at a concentration of 20  $\mu$ g/mL. Among the compounds tested,



577 compound **2** had the highest activity. The autoxidation products of oleic acid (**21** and **22**) and 8,11-  
578 dihydroxy derivative **23** also showed high activity. Oxidation of the hydroxyl group of **22** as ketone  
579 **24** maintained the activity, whereas elaidic acid, which lacks the 11-OH of **22**, had no activity at  
580 concentrations less than 80  $\mu\text{g/mL}$ . Ricinoleic acid having *cis*-double bond with a hydroxyl group at  
581 the homoallylic position displayed the same level of the activity as the *trans*-allylic alcohols. Taken  
582 together, presence of oxygen functional group(s) such as hydroxyl and carbonyl group is necessary  
583 for the activity, but the positions of the hydroxyl group and the geometry of the double bond are less  
584 important. Reduction of the carboxyl group to alcohol **20** caused somewhat decrease in activity  
585 compared with carboxylic acid **2**, but still maintained a moderate level of activity, indicating that the  
586 carboxyl group may be replaced with other polar functional groups. Compound **10** having triple  
587 bond had the same level of activity as **4**. Fig. 3. shows the cell of *C. antiqua* treated with 5  $\mu\text{g/mL}$  of  
588 compound **10** (A) and compound **2** (B) after 0.5- and 4-hour incubations. Interestingly, this  
589 propargylic alcohol **10** caused acute lysis of planktonic cells within 30 min (Fig. 3, A), at which  
590 period no other allylic alcohols affected the planktonic cells (Fig. 3, B).

591 (*E*)-9-Hydroxyoctadec-10-enoic acid (**1**) and (*E*)-10-hydroxyoctadec-8-enoic acid (**2**) have  
592 previously been isolated as the biotransformation products of oleic acid by *Pseudomonas* sp.  
593 [25,26,27,28,29]. The oxidation of unsaturated fatty acids proceeds via three different pathways;  
594 autoxidation, photo-oxidation and enzymatic oxidation such as that of lipoxygenases. Autoxidation  
595 of oleic acid involves allylic oxidation and allylic rearrangement of the resulting hydroperoxide, and  
596 is characterized by the formation of both *cis* and *trans* isomers of 8-hydroxyoctadec-9-enoic acid (8-  
597  $\text{OH}\Delta_{9,10}$ ) and 11-hydroxyoctadec-9-enoic acid (11- $\text{OH}\Delta_{9,10}$ ), and the *trans* isomers of 9- $\text{OH}\Delta_{10,11}$  (**1**)  
598 and 10- $\text{OH}\Delta_{8,9}$  (**2**) [24]. Photo-oxidation of oleic acid involves concerted ene reactions with a singlet  
599 oxygen, in which the oxidation proceeds at one end of the double bond to predominantly produce  
600 *trans*-9- $\text{OH}\Delta_{10,11}$  (**1**) and *trans*-10- $\text{OH}\Delta_{8,9}$  (**2**) [30]. (*E*)-11-hydroxyoctadec-12-enoic acid (**3**) and (*E*)-  
601 12-Hydroxyoctadec-10-enoic acid (**4**) may arise from *cis*-vaccenic acid by the same mechanism as  
602 that for **1** and **2**. Since oleic acid is widely distributed in nature, hydroxy acids **1** and **2** have been  
603 isolated from several plants and microorganisms; in some cases, both compounds were co-isolated  
604 from the same natural source. Compounds **1** and **2** isolated from stroma of the timothy plant  
605 *Epichloe typhina* showed antifungal activity against plant-pathogenic *Cladosporium herbarum* [31],  
606 and those isolated from the medicinal plant *Alternanthera brasiliana* and its endophytic bacteria had  
607 antimicrobial activity against some human pathogenic bacteria [32]. These hydroxy acids have also  
608 been found in macroalgae. Compound **2** isolated from the red alga *Gracilaria verrucosa* is reported  
609 to have moderate anti-inflammatory activity [33] and compound **1** isolated from the green alga  
610 *Caulerpa racemosa* exhibited potent protein tyrosine phosphatase 1B (PTP1B) inhibitory activity  
611 [34]. In contrast, (*E*)-11-hydroxyoctadec-12-enoic acid (**3**) and (*E*)-12-hydroxyoctadec-10-enoic acid  
612 (**4**) derived from *cis*-vaccenic acid have rarely been found in nature. Compound **3** was isolated from

613 the green alga *Ulva fasciata* Delile and shown to have moderate and weak antibacterial activity  
614 against *Streptomyces aureus* and *Escherichia coli*, respectively [35]. Compounds **1**~**4** have been  
615 detected in particulate matter and sediment samples collected in the northwestern Mediterranean Sea  
616 in GC/EIMS [36]. Nevertheless, to our knowledge, this is the first isolation of (*E*)-12-  
617 hydroxyoctadec-10-enoic acid (**4**) from living organisms. It has also been reported that the hydroxy  
618 lipids are the photo-oxidation products of oleic and *cis*-vaccenic acids generated in senescent  
619 phytoplanktonic cells [36]. Thereafter, Rontani et al. [37] investigated the origin of the *cis*-vaccenic  
620 acid photo-oxidation products in marine environment and concluded that heterotrophic bacteria that  
621 are attached to senescent phytoplanktonic cells most likely constitute the source of *cis*-vaccenic acid  
622 oxidation products **3** and **4** detected in the particulate matter samples.

623 Although the exact ratio of the four compounds was not determined, a GC/EI-MS spectrum of  
624 the mixture fraction **f5** displayed two peaks at  $t_R$  18.14 min and  $t_R$  18.21 min in a ratio of 59:41, the  
625 former being attributed to a mixture of compounds **3** and **4** and the latter to a mixture of compounds  
626 **1** and **2** (Supplementary data). It is interesting that the hydroxy fatty acids derived from *cis*-vaccenic  
627 acid are dominant over those from oleic acid in this alga.

628

#### 629 4. Conclusions

630

631 We isolated a highly algicidal fraction **f5** comprising four C18 hydroxy unsaturated fatty acids,  
632 (*E*)-9-hydroxyoctadec-10-enoic acid (**1**), (*E*)-10-hydroxyoctadec-8-enoic acid (**2**), (*E*)-11-  
633 hydroxyoctadec-12-enoic acid (**3**) and (*E*)-12-hydroxyoctadec-10-enoic acid (**4**), from a methanol  
634 extract of *T. jejuensis*. Their structures were confirmed by comparison of their spectral data with  
635 those of synthesized compounds. Among them, compound **2** was found to have the highest algicidal  
636 activity, showing >95% mortality against *C. antiqua* at a concentration of 5 µg/mL after 24 h. We  
637 also found that propargylic derivative **10** had high acute toxicity to the phytoplankton. Further  
638 detailed biological activity study to evaluate the effectiveness of these hydroxy lipids as anti-red tide  
639 agents and to obtain an insight on the mode of action are in progress.

640

#### 641 Declaration of Competing Interest

642

643 The authors declare no conflict of interest.

644

#### 645 Acknowledgments

646

647 We are grateful to the members of Laboratory of Marine Food Hygiene, Faculty of Fisheries,  
648 Nagasaki University in 2016 for collecting and providing us *T. jejuensis*. We also thank Dr. Tatsuya

649 Oda (Faculty of Fisheries, Nagasaki University) for providing the cultured strain of *Chattonella*  
650 *antiqua*. This work was the result of using research equipment shared in MEXT Project for  
651 promoting public utilization of advanced research infrastructure (Program for supporting  
652 introduction of the new sharing system) Grant Number JPMXS0422500320.

653

654 Appendix A. Supplementary data

655

656 Algicidal screening data of 17 seaweed extracts, NMR and Mass spectra of fraction **f5** and  
657 synthesized compounds used for bioassay are available as supplementary materials. Supplementary  
658 data to this article can be found online at

659

660 References

661

662 [1] J. J. Gallardo-Rodríguez, A. Astuya-Villalón A. Llanos-Rivera, V. Avello-Fontalba, V. Ulloa-  
663 Jofré, A critical review on control methods for harmful algal blooms, *Reviews in Aquaculture* 11  
664 (2019) 661-664. DOI: 10.1111/raq.12251

665 [2] J. G. Jeong, H. J. Jin, C. H. Sohn, K. H. Suh, Y. K. Hong, Algicidal activity of the seaweed  
666 *Corallina pilulifera* against red tide microalgae, *J. Appl. Phycol.* 12 (2000) 37-43.

667 [3] S. Nakai, Y. Inoue, M. Hosomi, A. Murakami, *Myriophyllum spicatum*-released allelopathic  
668 polyphenols inhibiting growth of blue-green algae *Microcystis aeruginosa*, *Water Res.* 34 (2000)  
669 3026-3032. [https://doi.org/10.1016/S0043-1354\(00\)00039-7](https://doi.org/10.1016/S0043-1354(00)00039-7)

670 [4] for a review see, Y. Sun, S. Dong, W. Zhou, L. Guo, G.-L. Guo, X. Zhang, A comprehensive  
671 review of secondary metabolites with antialgal activity from marine macroalgae against red tide  
672 microalgae, *J. Coast. Res.* 93(S1) (2019) 475-488.

673 [5] H. Kakisawa, F. Asari, T. Kusumi, T. Toma, T. Sakurai, T. Oohusa, Y. Hara, M. Chiharai, An  
674 allelopathic fatty acid from the brown alga *Cladosiphon okamuranus*, *Phytochem.* 27 (1988) 731-  
675 735. [https://doi.org/10.1016/0031-9422\(88\)84084-6](https://doi.org/10.1016/0031-9422(88)84084-6)

676 [6] I. Z. Chiang, W. Y. Huang, J. T. Wu, Allelochemicals of *Botryococcus braunii* (Chlorophyceae),  
677 *J. Phycol.* 40 (2004) 474-480. <https://doi.org/10.1111/j.1529-8817.2004.03096.x>

678 [7] M. A. Alamsjah, S. Hirao, F. Ishibashi, Y. Fujita, Isolation and structure determination of  
679 algicidal compounds from *Ulva fasciata*, *Biosci. Biotechnol. Biochem.* 69 (2005) 2186-2192.  
680 <https://doi.org/10.1271/bbb.69.2186>

681 [8] Q. H. Luyen, J. Y. Cho, J. S. Choi, J. Y. Kang, N. G. Park, Y. K. Hong, Isolation of algal spore  
682 lytic C17 fatty acid from the crustose coralline seaweed *Lithophyllum yessoense*. *J. Appl. Phycol.* 21  
683 (2009) 423-427. DOI 10.1007/s10811-008-9387-4

684 [9] R. Wang, Y. Wang, X. Tang, Identification of the toxic compounds produced by *Sargassum*

685 *thunbergii* to red tide microalgae, Chin. J. Oceanol. Limnol. 30 (2012) 778-785. DOI:  
686 10.1007/s00343-012-1294-5

687 [10] S. Hirao, K. Tara, K. Kuwano, J. Tanaka, F. Ishibashi, Algicidal Activity of Glycerolipids from  
688 brown alga *Ishige sinicola* toward red tide microalgae, Biosci. Biotechnol. Biochem. 76 (2012) 72-  
689 374. <https://doi.org/10.1271/bbb.110645>

690 [11] Y. Y. Sun, H. Wang, G. L. Guo, Y. F. Pu, B. L. Yan, C. H. Wang, Isolation, purification, and  
691 identification of antialgal substances in green alga *Ulva prolifera* for antialgal activity against the  
692 common harmful red tide microalgae, Environ. Sci. Pollut. Res. 23 (2016) 1149-1459. DOI  
693 10.1007/s11356-015-5377-7

694 [12] Y. Y. Sun, S. Dong, G. Guo, L. Guo, Y. Pu, Antialgal activity of glycolipids derived  
695 from a green macroalgae *Ulva prolifera* on six species of red tide microalgae, IOP Conf. Series:  
696 Materials Science and Engineering 484 (2019) 012057. DOI:10.1088/1757-899X/484/1/012057

697 [13] J. Y. Kim, M. A. Alamsjah, A. Hamada, Y. Fujita, F. Ishibashi, Algicidal diterpenes from the  
698 brown alga *Dictyota dichotoma*, Biosci. Biotechnol. Biochem. 70 (2006) 2571-2574.  
699 <https://doi.org/10.1271/bbb.60281>

700 [14] H. Lu, H. Xie, Y. Gong, Q. Wang, Y. Yang, Secondary metabolites from the seaweed  
701 *Gracilaria lemaneiformis* and their allelopathic effects on *Skeletonema costatum*, Biochem. Syst.  
702 Ecol. 39 (2011) 397-400. <https://doi.org/10.1016/j.bse.2011.05.015>

703 [15] Y. Y. Sun, K. Meng, Z. X. Su, G. L. Guo, Y. F. Pu, C. H. Wang, Isolation and purification of  
704 antialgal compounds from the red alga *Gracilaria lemaneiformis* for activity against common  
705 harmful red tide microalgae, Environ. Sci. Pollut. Res. 24 (2017) 4964-4972. DOI 10.1007/s11356-  
706 016-8256-y

707 [16] F. Ishibashi, S. Sato, K. Sakai, S. Hirao, K. Kuwano, Algicidal sesquiterpene hydroquinones  
708 from the brown Alga *Dictyopteria undulata*, Biosci. Biotechnol. Biochem. 77 (2013) 1120-1122.  
709 <https://doi.org/10.1271/bbb.130018>

710 [17] Y. Y. Sun, W. J. Zhou, H. Wang, G. L. Guo, Z. X. Su, Y. F. Pu, Antialgal compounds with  
711 antialgal activity against the common red tide microalgae from a green algae *Ulva pertusa*, Ecotox.  
712 Environ. Safe. 157 (2018) 61-66. <https://doi.org/10.1016/j.ecoenv.2018.03.051>

713 [18] C. Litchfield, R. D. Harlow, A. F. Isbell, R. Reiser, *Cis-trans* isomerization of oleic acid by  
714 nitrous acid, J. Am. Oil. Chem. 42 (1965) 73-78.

715 [19] G. Knothe, D. Weisleder, M. O. Bagby, R. E. Peterson, Hydroxy fatty acids through  
716 hydroxylation of oleic acid with selenium dioxide/*tert*-butylhydroperoxide, J. Am. Oil. Chem. 70  
717 (1993) 401-404.

718 [20] J. F. Rontani, Photodegradation of unsaturated fatty acids in senescent cells of phytoplankton:  
719 photoproduct structural identification and mechanistic aspects, J. Photochem. Photobiol. A 114  
720 (1998) 37-44. [https://doi.org/10.1016/S1010-6030\(98\)00201-9](https://doi.org/10.1016/S1010-6030(98)00201-9)

721 [21] R. Kleiman, G. F. Spencer, Gas chromatography-mass spectrometry of methyl esters of  
722 unsaturated oxygenated fatty acids, *J. Am. Oil. Chem.* 50 (1973) 31-38.

723 [22] S. Ravi, D. Padmanabhan, V. R. Mamdapur, Macrocyclic musk compounds: Synthetic  
724 approaches to key intermediates for exaltolide, exaltone and dilactones, *J. Indian. Inst. Sci.* 81 (2001)  
725 299-312.

726 [23] S. R. Abrams, A. C. Shaw, Triple-bond isomerizations: 2- to 9-decyn-1-ol, *Org. Synth.* 66  
727 (1988) 127-129. DOI:10.15227/orgsyn.066.0127

728 [24] E. N. Frankel, W. E. Neff, W. K. Rohwedder, B. P. S. Khambay, R. F. Garwood, B. C. L.  
729 Weedon, Analysis of autoxidized fats by gas chromatography-mass spectrometry: I. Methyl oleate,  
730 *Lipids* 12 (1977) 901-907.

731 [25] A. Guerrero, I. Casals, M. Basquets, Y. Leon, A. Manresa, Oxidation of oleic acid to (*E*)-10-  
732 hydroperoxy-8-octadecenoic and (*E*)-10-hydroxy-8-octadecenoic acids by *Pseudomonas* sp. 42A2,  
733 *Biochim. Biophys. Acta.* 1347 (1997) 75-81.

734 [26] J. Bastida, C. de Andrés, J. Culleré, M. Busquets, A. Manresa, Biotransformation of oleic acid  
735 into 10-hydroxy-8*E*-octadecenoic acid by *Pseudomonas* sp. 42A2, *Biotechnol. Lett.* 21 (1999) 1031-  
736 1035.

737 [27] E. Rodríguez, M. J. Espuny, A. Manresa, A. Guerrero A, Identification of (*E*)-11-hydroxy-9-  
738 octadecenoic acid and (*E*)-9-hydroxy-10-octadecenoic acid by biotransformation of oleic acid by  
739 *Pseudomonas* sp. 32T3, *J. Am. Oil. Sci.* 78 (2001) 593-597.

740 [28] J. Culleré, O. Durany, M. Busquets, A. Manresa, Biotransformation of oleic acid into (*E*)-10-  
741 hydroxy-8-octadecenoic acid and (*E*)-7,10-dihydroxy-8-octadecenoic acid by *Pseudomonas* sp.  
742 42A2 in an immobilized system, *Biotechnol. Lett.* 23 (2001) 215-219.

743 [29] E. Martínez, M. Hamberg, M. Busquets, P. Díaz, A. Manresa, E. H. Oliw, Biochemical  
744 characterization of the oxygenation of unsaturated fatty acids by the dioxygenase and hydroperoxide  
745 isomerase of *Pseudomonas aeruginosa* 42A2, *J. Biol. Chem.* 285 (2010) 9339-9345. DOI  
746 10.1074/jbc.M109.078147

747 [30] E. N. Frankel, W. E. Neff, T. R. Bessler, Analysis of autoxidized fats by gas chromatography-  
748 mass spectrometry: V. Photosensitized oxidation, *Lipids*, 14 (1979) 961-967

749 [31] H. Koshino, S. Togiya, T. Yoshihara, S. Sakamura, T. Shimanuki, T. Sato, A. Tajimi, Four  
750 fungitoxic C-18 hydroxy unsaturated fatty acids from stroma of *Epichloe typhina*, *Tetrahedron Lett.*  
751 28 (1987) 73-76.

752 [32] M. A. Trapp, M. Kai, A. Mithöfer, E. Rodrigues-Filho, Antibiotic oxylipins from *Alternanthera*  
753 *brasiliensis* and its endophytic bacteria, *Phytochem.* 110 (2015) 72-82.  
754 <https://doi.org/10.1016/j.phytochem.2014.11.005>

755 [33] H. T. Dang, H. J. Lee, E. S. Yoo, P. B. Shinde, Y. M. Lee, J. Hong, D. K. Kim, J. H. Jung, Anti-  
756 inflammatory Constituents of the Red Alga *Gracilaria verrucosa* and Their Synthetic Analogues, *J*

757 Nat. Prod. 71 (2008) 232-240. <https://doi.org/10.1021/np070452q>  
758 [34] D. Q. Liu, J. Li, H. Y. Zhang, M. T. Feng, H. Yang, P. Yang, K. Lin, Y. W. Guo, S. C. Mao,  
759 The fatty acids of the green alga *Caulerpa racemose* and their bioactivities, Chinese Journal of  
760 Marine Drugs 32 (2013) 13-20. DOI:10.13400/j.cnki.cjmd.2013.06.006  
761 [35] G. S. E. Abou-ElWafa, M. Shaaban, K. A. Shaaban, M. E. E. El-Naggar, H. Laatsch H, Three  
762 New Unsaturated Fatty Acids from the Marine Green Alga *Ulva fasciata* Delile, Z. Natureforsch. 64b  
763 (2009) 1199-1207. <https://doi.org/10.1515/znb-2009-1014>  
764 [36] D. Marchand, J. F. Rontani, Characterization of photo-oxidation and autoxidation products of  
765 phytoplanktonic monounsaturated fatty acids in marine particulate matter and recent sediments, Org.  
766 Geochem. 32 (2001) 287-304. [https://doi.org/10.1016/S0146-6380\(00\)00175-3](https://doi.org/10.1016/S0146-6380(00)00175-3)  
767 [37] J. F. Rontani, M. Koblížek, B. Beker, P. Bonin, Z. S. Kolber, On the origin of *cis*-vaccenic acid  
768 photodegradation products in the marine environment, Lipids 38 (2003) 1085-1092.  
769 <https://doi.org/10.1007/s11745-006-1164-z>  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786

**Table 1.** Chemical shift values ( $\delta_C$ , ppm) of the selected carbons of fraction **f5** and compounds **1~4** (125 MHz in CDCl<sub>3</sub>)

<sup>13</sup> C atom	fraction/compound					chemical shift difference
	<b>f5</b>	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>	
<u>C</u> H <sub>3</sub>	14.05			14.03		0.02
	14.10				14.07	0.03
	14.10	14.08				0.02
	14.11		14.09			0.02
<u>C</u> H <sub>2</sub> -CH=CH-CH(OH)	31.36			31.34		0.02
	31.82				31.80	0.02
	31.84	31.82				0.02
	31.87		31.86			0.01
CH <sub>2</sub> - <u>C</u> H=CH-CH(OH)	131.97		131.97			0.00
	132.20				132.19	0.01
	132.28			132.29		0.01
	132.31	132.34				0.03
CH <sub>2</sub> -CH= <u>C</u> H-CH(OH)	132.93	132.85				0.05
	132.94			132.89		0.05
	132.97				132.94	0.03
	133.16		133.12			0.04
CH <sub>2</sub> -CH=CH- <u>C</u> H(OH)	73.20	73.23				0.03
	73.25			73.27		0.02
	73.26		73.27			0.01
	73.30				73.30	0.00
<u>C</u> H <sub>2</sub> -COOH	33.54		33.88			0.34
	33.57			33.93		0.36
	33.59				33.95	0.36
	33.59	33.95				0.36
CH <sub>2</sub> - <u>C</u> OOH	177.09		179.08			1.99
	177.12				179.22	2.11
	177.16	179.22				2.06
	177.18			179.23		2.05






Fig. 1. Structures of compound 1~4






**Scheme 1.** Synthesis of (*E*)-hydroxyoctadec-10-enoic acid (**1**) and (*E*)-12-hydroxyoctadec-10-enoic acid (**4**)

*Reagents:* (a) BuLi, HMPA-THF, 0 °C, 2 h, then 1-bromoheptane, r.t., 18 h (27%); (b) TMSCHN<sub>2</sub>, DCM-MeOH, rt (100%); (c) H<sub>2</sub>, 5% Pd/Ca-Pb, EtOAc, rt (96%); (d) SeO<sub>2</sub>, TBHP, DCM, rt, 50 h (74%); (e) NaOH, MeOH, reflux, **1** (89%), **2** (85%); (f) BuLi, THF, rt, 20 min, then heptanal, -78 °C to rt, 1.5 h (48%); (g) Li, NH<sub>3</sub>, *t*-BuOH, THF, -78 °C, 2 h (38%); (h) 4-bromophenacyl bromide, K<sub>2</sub>CO<sub>3</sub>, acetone, rt, **1a** (66%), **4a** (43%).




**Scheme 2.** Synthesis of (*E*)-11-hydroxyoctadec-12-enoic acid (**3**)

*Reagents:* (a) hexane-toluene,  $-10\text{ }^\circ\text{C}$ , 1 h (28%); (b) NaOH, MeOH, rt, **3** (72%); (c) 4-bromophenacyl bromide,  $\text{K}_2\text{CO}_3$ , acetone (52%).




**Scheme 3.** (*E*)-10-hydroxyoctadec-8-enoic acid (**2**) and (*E*)-1,10-dihydroxyoctadec-8-ene (**20**)

*Reagents:* (a) DHP, *p*-TsOH, DCM, rt, 18 h (69%); (b) DIBAL, hexane-toluene, 60 °C, 2 h then nonanal, -60 °C, 2 h (44%); (c) Ac<sub>2</sub>O, pyridine, rt 40 h; (d) *p*-TsOH, EtOH, rt, 25 min (33%, 2 steps); (e) PDC, DMF, rt 17 h (45%); (f) NaOH, MeOH, rt, (36%); (g) PPTS, MeOH, rt, 20 h (46%); (h) 4-bromophenacyl bromide, K<sub>2</sub>CO<sub>3</sub>, acetone (59%).




**Fig. 2.** Algicidal activity [mortality (%)] of compound 1~4 and its related compounds at concentrations of 80, 20 and 5 µg/mL for 24 h against *C. antiqua*. Values are the mean ± SD from three independent experiments.



**Scheme 4.** Syntheses of (*E*)-8-hydroxyoctadec-9-enoic acid (**21**), (*E*)-11-hydroxyoctadec-9-enoic acid (**22**), (*E*)-8,11-dihydroxyoctadec-9-enoic acid (**23**), and (*E*)-11-oxooctadec-9-enoic acid (**24**).

*Reagents:* (a)  $\text{SeO}_2$ , *t*-BuOOH,  $\text{CH}_2\text{Cl}_2$ , rt, 72 h; (b) NaOH, MeOH; (c) Dess-Martin periodinane,  $\text{CH}_2\text{Cl}_2$ , rt, 18 h (63%)



**Fig. 3.** The cell of *Chattonella antiqua* treated with compound **10** (A) and compound **2** (B) at a concentration of 5 μg/mL each, and untreated cells (C) just after treatment (0 h) and after 0.5- and 4-hour incubations. Arrowheads indicate debris of dead cells of *C. antiqua* cells. Bar indicates 100 μm.